




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 209/18, A61K 31/40, C07D 401/06, 403/06, 417/06, 405/06, 409/10, 405/10, 401/10, 417/10, 413/06, 409/06		A1	(11) International Publication Number: WO 99/35130
(21) International Application Number: PCT/IB98/02065		(43) International Publication Date: 15 July 1999 (15.07.99)	
(22) International Filing Date: 18 December 1998 (18.12.98)		(74) Agents: SPIEGEL, Allen, J. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).	
(30) Priority Data: PCT/IB98/00003 5 January 1998 (05.01.98) CA		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for JP only): PFIZER PHARMACEUTICALS INC. [JP/JP]; Mitsui Building, 1-1, Nishi-shinjuku 2-chome, Shinjuku-ku, Tokyo 163 (JP).		<p>Published With international search report.</p>	
(71) Applicant (for all designated States except JP US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): NAKAO, Kazunari [JP/JP]; 29-1, Shinbayashi, Shinbayashi-cho, Chiryu-shi, Aichi-ken 472-0017 (JP). STEVENS, Rodney, William [AU/JP]; 3-4-26, Seishiro-cho, Handa-shi, Aichi-ken 475-0917 (JP). KAWAMURA, Kiyoshi [JP/JP]; 40-1, Daimonsaki, Inuyama, Inuyama-shi, Aichi-ken 484-0081 (JP). UCHIDA, Chikara [JP/JP]; 118-401, Miyaji-cho, Handa-shi, Aichi-ken 475-0902 (JP). KOIKE, Hiroki [JP/JP]; 1-100, Souga-cho, Handa-shi, Aichi-ken			

(54) Title: **2,3-SUBSTITUTED INDOLE COMPOUNDS AS COX-2 INHIBITORS**

(I)

(57) Abstract

This invention provides a compound of formula (I) or the pharmaceutically acceptable salts thereof wherein Z is OH, C₁₋₆ alkoxy, -NR²R³ or heterocycle; Q is selected from the following: (a) an optionally substituted phenyl, (b) an optionally substituted 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), (c) an optionally substituted 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, (d) an optionally substituted C₃₋₇ cycloalkyl and (e) an optionally substituted benzofused heterocycle; R¹ is hydrogen, C₁₋₄ alkyl or halo; R² and R³ are independently hydrogen, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy or CN; X is independently selected from H, halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino and CN; and n is 0, 1, 2, 3 and 4. This invention also provides a pharmaceutical composition useful for the treatment of a medical condition in which prostaglandins are implicated as pathogens.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

2,3-SUBSTITUTED INDOLE COMPOUNDS AS COX-2 INHIBITORS

Technical Field

This invention relates to novel 2,3-substituted indoles as pharmaceutical agents.

5 This invention specifically relates to compounds, compositions and methods for the treatment or alleviation of pain and inflammation and other inflammation-associated disorders, such as arthritis.

Background Art

Nonsteroidal antiinflammatory drugs (NSAIDs) are widely used in treating pain

10 and the signs and symptoms of arthritis because of their analgesic and anti-inflammatory activity. It is accepted that common NSAIDs work by blocking the activity of cyclooxygenase (COX), also known as prostaglandin G/H synthase (PGHS), the enzyme that converts arachidonic acid into prostanoids. Prostaglandins, especially prostaglandin E₂ (PGE₂), which is the predominant eicosanoid detected in

15 inflammation conditions, are mediators of pain, fever and other symptoms associated with inflammation. Inhibition of the biosynthesis of prostaglandins has been a therapeutic target of anti-inflammatory drug discovery. The therapeutic use of conventional NSAIDs is, however, limited due to drug associated side effects, including life threatening ulceration and renal toxicity. An alternative to NSAIDs is

20 the use of corticosteroids, however, long term therapy can also result in severe side effects.

Recently, two forms of COX were identified, a constitutive isoform (COX-1) and an inducible isoform (COX-2) of which expression is upregulated at sites of inflammation (Vane, J. R.; Mitchell, J. A.; Appleton, I.; Tomlinson, A.; Bishop-Bailey,

25 D.; Croxtoll, J.; Willoughby, D. A. *Proc. Natl. Acad. Sci. USA*, 1994, 91, 2046). COX-1 is thought to play a physiological role and to be responsible for gastrointestinal and renal protection. On the other hand, COX-2 appears to play a pathological role and to be the predominant isoform present in inflammation conditions. A pathological role for prostaglandins has been implicated in a number of human disease


30 states including rheumatoid and osteoarthritis, pyrexia, asthma, bone resorption, cardiovascular diseases, nephrotoxicity, atherosclerosis, hypotension, shock, pain,

cancer, and Alzheimer disease. The NSAIDs currently on market inhibit both isoforms of COX with little variation for selectivity, explaining their beneficial (inhibition of COX-2) and deleterious effects (inhibition of COX-1). It is believed that compounds that would selectively inhibit the biosynthesis of prostaglandins by
 5 intervention of the induction phase of the inducible enzyme cyclooxygenase-2 and/or by intervention of the activity of the enzyme cyclooxygenase-2 on arachidonic acid would provide alternate therapy to the use of NSAIDs or corticosteroids in that such compounds would exert anti-inflammatory effects without the adverse side effects associated with COX-1 inhibition.

10 A variety of indole compounds are known and are disclosed in several patent applications. The International Publication Numbers WO 96/32379 discloses N-substituted indole compounds as cGMP-PDE Inhibitors. The International Publication Numbers WO 96/37467, WO 96/37469, UK Patent Publication GB 2283745 A and US Publication Number 5510368 disclose 2-methyl-N-substituted
 15 indole compounds as cyclooxygenase-2 Inhibitors. Also, a variety of indole compounds are disclosed as agents for controlling underwater fouling organisms in European Patent Publication Number 0 556 949 A2 by Konya, Kazumi et.al. Specifically, the International Publication Numbers WO 97/09308 discloses indole compounds as neuropeptide receptor antagonists. Besides, in Sci. Pharm. 64, 577
 20 (1996), a process for preparing a 2-ester-substituted indoline is disclosed.

Brief Disclosure of the Invention


The present invention provides a compound of the following formula:



(I)

25 or the pharmaceutically acceptable salts thereof wherein

Z is OH, C₁₋₆ alkoxy, -NR²R³ or a group of the formula (II) or (III):



wherein r is 1, 2, 3 or 4, Y is a direct bond, O, S or NR^4 , and W is OH or $\text{---NR}^2\text{R}^3$;

Q is selected from the following:

- 5 (a) phenyl optionally substituted with one, two or three substituents independently selected from
 - (a-1) halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(\text{C}_{1-4}$ alkyl)amino, C_{1-4} alkylamino, CN, HO- (C_{1-4}) alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $\text{---NH}_2\text{S(O)}_2\text{NR}^2\text{R}^3$, acetyl, ---COOH , $\text{---C(O)O---C}_{1-4}$ alkyl, C_{1-4} alkylsulfonylamino and C_{3-7} cycloalkyl,
 - (a-2) aryl or $\text{---O---(CH}_2\text{)}_n\text{---aryl}$, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(\text{C}_{1-4}$ alkyl)amino, C_{1-4} alkylamino and CN,
 - (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(\text{C}_{1-4}$ alkyl)amino, C_{1-4} alkylamino and CN,
 - (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(\text{C}_{1-4}$ alkyl)amino, C_{1-4} alkylamino and CN,
- (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently

selected from the above group (a-1), (a-2), (a-3) and (a-4),

- 5 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- 10 (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and
- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R¹ is hydrogen, C₁₋₄ alkyl or halo;

15 R² and R³ are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

R⁴ is hydrogen or C₁₋₄ alkyl;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, 20 aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; and

n is 0, 1, 2, 3 or 4.

25 The indole compounds of the present invention exhibit inhibition of COX activity. Preferably compounds of this invention exhibit inhibitory activity against COX-2, with more preferable compounds having COX-2 selectivity.

Accordingly, the present invention also provides a pharmaceutical composition, useful for the treatment of a medical condition in which prostaglandins are implicated as pathogens, which comprises a compound of the formula (I) and the pharmaceutically acceptable salts thereof.

30 Further, the present invention provides a method for the treatment of a medical condition in which prostaglandins are implicated as pathogens, in a mammalian subject,

which comprises administering to said subject a therapeutically effective amount of said pharmaceutical composition.


The medical conditions in which prostaglandins are implicated as pathogens, include the relief of pain, fever and inflammation of a variety of conditions including
5 rheumatic fever, symptoms associated with influenza or other viral infections, common cold, low back and neck pain, dysmenorrhea, headache, toothache, sprains and strains, myositis, neuralgia, synovitis, arthritis including rheumatoid arthritis, degenerative joint disease (osteoarthritis), gout, ankylosing spondylitis, systemic lupus erythematosus and juvenile arthritis, bursitis, burns, injuries following surgical and
10 dental procedures.

The compounds and pharmaceutical composition of this invention may inhibit cellular neoplastic transformations and metastatic tumor growth and thus may be used in the treatment and/or prevention of cancers in the colon, breast, skin, esophagus, stomach, urinary bladder, lung and liver. The compounds and pharmaceutical
15 composition of this invention were used in the treatment and/or prevention of cyclooxygenase-mediated proliferation disorders such as which occur in diabetic retinopathy and tumor angiogenesis.

The compounds and pharmaceutical composition of this invention may inhibit prostanoic-induced smooth muscle contraction by preventing the synthesis of
20 contractile prostanoids, and thus may be of use in the treatment of dysmenorrhea, premature labor, asthma and eosinophil related disorders and in the treatment of neurodegenerative diseases such as Alzheimer's and Parkinson's disease, and for the treatment of bone loss (treatment of osteoarthritis), stroke, seizures, migraine, multiple sclerosis, AIDS and encephalopathy.

By virtue of the COX-2 activity and/or specificity for COX-2 over COX-1, such
25 compounds will prove useful as an alternative to conventional NSAIDs particularly where such NSAIDs may be contra-indicated such as in patients with ulcers (such as peptic ulcers and gastric ulcers), gastritis, regional enteritis, ulcerative colitis, diverticulitis or with a recurrent history of GI lesions, GI bleeding, coagulation
30 disorders including anemia such as hypoprothrombinemia, haemophilia and other bleeding problems; kidney disease; prior to surgery of taking of anticoagulants.

Also, the present invention provides a compound of formula 7-VI:



7-VI

wherein **B** is a suitable protecting group;

5 **Q** is selected from the following:

(a) phenyl optionally substituted with one, two or three substituents independently selected from

(a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,

(a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄


alkyl)amino, C₁₋₄ alkylamino and CN,

- (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and
- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R² and R³ are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; **R⁵** is C₁₋₆ alkyl; and **n** is 0, 1, 2, 3 or 4.

Also, the present invention provides a compound of formula 7-V:



wherein **B** is a suitable protecting group;

Q is selected from the following:

- 5 (a) phenyl optionally substituted with one, two or three substituents independently selected from
- (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, 10 -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,
- (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di- 15 (C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN, 20
- (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN, 25
- 30 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- (c) a 5-membered monocyclic aromatic group containing one heteroatom

- selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- 5 (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and
- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);
- 10 **R² and R³** are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;
- X** is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; **R⁵** is C₁₋₆ alkyl; and
- 15 **n** is 0, 1, 2, 3 or 4.

Detailed Disclosure of the Invention

- 20 As used herein, "halo" is fluoro, chloro, bromo or iodo.
- As used herein, the term "C₁₋₄ alkyl" means straight or branched chain saturated radicals of 1 to 4 carbon atoms, including, but not limited to methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *sec*-butyl, *tert*-butyl, and the like.
- As used herein, an example of "propyl" is *n*-propyl and isopropyl.
- 25 As used herein, an example of "butyl" is *n*-butyl, isobutyl, *sec*-butyl and *tert*-butyl.
- As used herein, an example of "alkoxy" is methoxy, ethoxy, *n*-propoxy, isopropoxy, *n*-butoxy, isobutoxy, *sec*-butoxy, *tert*-butoxy, and the like.
- As used herein, an example of "alkylthio" is methylthio, ethylthio, *n*-propylthio, isopropylthio, *n*-butylthio, isobutylthio, *sec*-butylthio, *tert*-butylthio, and the like.
- 30 As used herein, an example of "di-(C₁₋₄ alkyl)amino" is dimethylamino,

diethylamino, dipropylamino, N-methyl-N-ethylamino, N-methyl-N-propylamino, N-methyl-N-butylamino, N-ethyl-N-propylamino, and the like.

As used herein, an example of "C₁₋₄ alkylamino" is methylamino, ethylamino, n-propylamino, isopropylamino, n-butylamino, isobutylamino, *sec*-butylamino, *tert*-butylamino, and the like.

As used herein, an example of "HO-(C₁₋₄)alkyl" is hydroxymethyl, hydroxyethyl (e.g., 1-hydroxyethyl and 2-hydroxyethyl), hydroxypropyl (e.g., 1-hydroxypropyl, 2-hydroxypropyl and 3-hydroxypropyl)

As used herein, an example of "C₁₋₄ alkoxy-C₁₋₄ alkyl" is methoxymethyl, methoxyethyl, methoxypropyl, methoxybutyl, ethoxymethyl, ethoxyethyl, ethoxypropyl, and the like.

As used herein, the term "halo-substituted alkyl" refers to an alkyl radical as described above substituted with one or more halogens included, but not limited to, chloromethyl, dichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trichloroethyl, and the like.

As used herein, an example of "halo-substituted alkoxy" is chloromethoxy, dichloromethoxy, fluoromethoxy, difluoromethoxy, trifluoromethoxy, 2,2,2-trichloroethoxy, and the like.

As used herein, the term "C₃₋₇ cycloalkyl" means carbocyclic radicals, of 3 to 7 carbon atoms, including, but not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and the like.

As used herein, an example of "aryl" is phenyl and naphthyl.

As used herein, a 5-membered monocyclic aromatic group usually has one heteroatom selected from O, S and N in the ring. In addition to said heteroatom, the monocyclic aromatic group may optionally have up to three N atoms in the ring. For example, the 5-membered monocyclic aromatic group includes thienyl, furyl, thiazolyl (e.g., 1,3-thiazolyl, 1,2-thiazolyl), imidazolyl, pyrrolyl, oxazolyl (e.g., 1,3-oxazolyl, 1,2-oxazolyl, isoxazolyl), pyrazolyl, tetrazolyl, triazolyl (e.g., 1,2,3-triazolyl, 1,2,4-triazolyl), oxadiazolyl (e.g., 1,2,3-oxadiazolyl), thiadiazolyl (e.g., 1,3,4-thiadiazolyl, 1,2,3-thiadiazolyl) and the like.

As used herein, an example of a 6-membered monocyclic aromatic group

includes pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl (e.g., 1,3,5-triazinyl), tetrazinyl and the like.

As used herein, an example of a benzo-fused heterocycle includes quinolyl, isoquinolyl, cinnolyl, quinoxalyl, benzoimidazolyl, benzothiazolyl, benzoxazolyl, 5 benzofuranyl, benzothiophenyl, indolyl, isoindolyl, 1H-indazolyl, quinazolinyl, phthalazinyl and the like.

As used herein, an example of (ethyl)(ethoxy)pyridyl includes 3-ethoxy-4-ethyl-2-pyridyl, 4-ethoxy-3-ethyl-2-pyridyl and the like.

As used herein, an example of (chloro)(ethyl)pyridyl includes 3-chloro-4-ethyl-2-10 pyridyl, 4-chloro-3-ethyl-2-pyridyl and the like.

As used herein, an example of (fluoro)(ethyl)phenyl includes 3-fluoro-4-ethyl-2-pyridyl, 4-fluoro-3-ethyl-2-pyridyl and the like.

Preferred compounds of this invention are those of the formula (I) wherein Z is OH, C₁₋₆ alkoxy, dimethylamino, methylamino, amino, N-methoxy-N-15 methylamino, 2-cyanoethylamino, 2-hydroxyethylamino, pyrrolidinyl, piperidino, piperazinyl, N-methyl-piperazinyl, morpholino, methoxyamino, piperazynyl, aminopyrrolidinyl or aminoethylamino.

Further preferred compounds of this invention are those of the formula (I) wherein

20 Z is OH or C₁₋₆ alkoxy; and

Q is selected from the following:

(a) phenyl optionally substituted with one, two or three substituents independently selected from

25 (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,

30 (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently

selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

5 (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

10 (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

15 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),

20 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);

25 (d) C₃₋₇ cycloalkyl selected from cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl, and the said cycloalkyl being optionally substituted with one substituent selected from OH, methyl, ethyl, propyl, F, Cl and CF₃; and

30 (e) a benzo-fused heterocycle selected from quinolyl, isoquinolyl, cinnolinyl, quinoxalinyl, benzoimidazolyl, benzothiazolyl, benzoxazolyl, benzofuranyl, benzothiophenyl and indolyl, and the benzo-fused heterocycle being optionally substituted with one, two, or three substituents independently selected from the group (a-1).

Further preferred compounds of this invention are those of the formula (I) wherein

Q is selected from the following:

- 5 (a) phenyl optionally substituted with one, two or three substituents independently selected from
- (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, 10 -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,
- (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di- 15 (C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN, 20
- (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN, 25
- (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4), 30
- (c) a 5-membered monocyclic aromatic group containing one heteroatom

selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);

- (d) cyclopropyl, cyclobutyl and cyclohexyl; and
- (e) quinolyl or isoquinolyl, and said quinolyl or isoquinolyl being optionally substituted with one substituent selected from the group halo, C₁₋₄ alkyl, NH₂, OH, C₁₋₄ alkoxy and C₁₋₄ halo-substituted alkyl.

Further preferred compounds of this invention are those of the formula (I) wherein

Z is OH, C₁₋₆ alkoxy;

Q is selected from the following:

- (a) phenyl optionally substituted with one, two or three substituents independently selected from

(a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,

(a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

- (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4); and
- (e) isoquinolyl;

R¹ is hydrogen or C₁₋₄ alkyl;

R² and R³ are independently H or methyl;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOR⁴, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; and

n is 0, 1, 2, or 3.

Further preferred compounds of this invention are those of the formula (I) wherein

Z is OH, methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, iso-butoxy or tert-butoxy;

Q is selected from the following:

- (a) phenyl optionally substituted with one or two substituents independently selected from

- (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, -COOH, C₁₋₄ alkylsulfonylamino, nitro, C₁₋₄ alkylsulfonyl and cyano,
- 5 (a-2) phenyl or benzyloxy, and the phenyl or phenyl moiety of benzyloxy being optionally substituted with one substituent selected from C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, halo, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy and NH₂,
- 10 (a-3) 5-membered monocyclic aromatic group selected from imidazolyl, thiazolyl, furyl, thienyl, pyrrolyl, tetrazolyl, triazolyl, oxazolyl, isoxazolyl, thiadiazolyl and pyrazolyl, and the 5-membered monocyclic aromatic group optionally being substituted with one substituent selected from C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, halo, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy and NH₂,
- 15 (a-4) 6-membered monocyclic aromatic group selected from pyridyl, pyrazinyl, pyrimidinyl and pyridazinyl, and the 6-membered monocyclic aromatic group optionally being substituted with one substituent selected from C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, halo, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy and NH₂,
- 20 (b) a 6-membered monocyclic aromatic group selected from pyridyl, pyrazinyl, pyrimidinyl and pyridazinyl, and said monocyclic aromatic group being optionally substituted with one or two substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- 25 (c) a 5-membered monocyclic aromatic group selected from imidazolyl, thiazolyl, furyl, thienyl, pyrrolyl, tetrazolyl, triazolyl, oxazolyl, isoxazolyl, thiadiazolyl and pyrazolyl, and said monocyclic aromatic group being optionally substituted with one or two substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- R¹ is hydrogen, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl or tert-butyl;
- 30 X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino,

C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl and aminosulfonyl; and

n is 0, 1, 2, or 3.

Further preferred compounds of this invention are those of the formula (I)
5 wherein

Z is OH, methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, iso-butoxy or tert-butoxy;

Q is selected from the following:

- 10 (a) phenyl optionally substituted with one or two substituents independently selected from
- (a-1) fluoro, chloro, bromo, iodo, methyl, ethyl, propyl, butyl, CH₂F, CHF₂, CF₃, methoxy, ethoxy, n-propoxy, n-butoxy, isopropoxy, CH₂F-O-, CHF₂-O-, CF₃-O-, methylthio, ethylthio, hydroxymethyl, methoxymethyl, methoxyethyl, ethoxymethyl, hydroxy, nitro, 15 methylsulfonyl, cyano, (HO)(H₃C)₂C-, acetyl and methylsulfonylamino,
- (a-2) phenyl or benzyloxy, and the phenyl or phenyl moiety of benzyloxy being optionally substituted with one substituent selected from methyl, ethyl, propyl, CF₃, F, Cl, OH, methoxy, ethoxy and NH₂,
- 20 (a-3) 5-membered monocyclic aromatic group selected from furyl, thienyl and pyrrolyl, and the 5-membered monocyclic aromatic group optionally being substituted with one substituent selected from methyl, ethyl, propyl, CF₃, F, Cl, OH, methoxy, ethoxy and NH₂,
- (a-4) pyridyl optionally substituted with one substituent selected from 25 methyl, ethyl, propyl, CF₃, F, Cl, OH, methoxy, ethoxy and NH₂,
- (b) pyridyl optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-1), (a-2), (a-3) and (a-4),
- 30 (c) imidazolyl, thiazolyl, furyl, thienyl, isoxazolyl, 1,2,3-thiadiazolyl or pyrrolyl, and said imidazolyl, thiazolyl, furyl, thienyl, isoxazolyl, 1,2,3-thiadiazolyl or pyrrolyl being optionally substituted with one or two

substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);

R^1 is hydrogen, methyl, ethyl, n-propyl, iso-propyl;

X is independently selected from fluoro, chloro, bromo, methyl, ethyl, propyl, butyl,
5 CH_2F , CHF_2 , CF_3 , methoxy, CF_3O or ethoxy; and

n is 0, 1 or 2.

Further preferred compounds of this invention are those of the formula (I) wherein

Z is OH, ethoxy or methoxy; Q is phenyl, chlorophenyl, fluorophenyl, bromophenyl,
10 methylphenyl, methoxyphenyl, (furyl)phenyl, trifluoromethylphenyl, trifluoromethoxyphenyl, pyridyl, methylpyridyl, ethylpyridyl, propylpyridyl, dimethylpyridyl, chloropyridyl, fluoropyridyl, trifluoromethylpyridyl, methoxypyridyl, (ethyl)(ethoxy)pyridyl, (chloro)(ethyl)pyridyl, thiazolyl, methylthiazolyl, furyl, methoxymethylfuryl, isoquinolyl, cyclohexyl, methoxyphenyl, (fluoro)(ethyl)pyridyl,
15 dimethylpyridyl or (ethoxy)(ethyl)pyridyl;

R^1 is hydrogen; X is fluoro, chloro, methyl, ethyl, isopropyl, tert-butyl, CF_3 or methoxy; and n is 1 or 2.

Further preferred compounds of this invention are those of the formula (I) wherein

20 Z is OH, ethoxy or methoxy; Q is phenyl, chlorophenyl, pyridyl, methylpyridyl, ethylpyridyl, propylpyridyl or chloropyridyl; R^1 is hydrogen; X is fluoro, chloro, methyl or CF_3 ; and n is 1 or 2.

Preferred individual compounds of this invention are:

ethyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
25 (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid, sodium salt;
[6-chloro-2-(2-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]acetic acid;
30 [6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate;

- [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
5 [2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3,4-dichlorobenzoyl)-1H-indol-3-yl]acetic acid;
(2-benzoyl-4-chloro-1H-indol-3-yl)acetic acid;
10 [5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
[2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
15 [5-methoxy-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
(2-benzoyl-7-chloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-4,5-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-4,6-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
20 *dl*-2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
less polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
more polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
25 methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
30 [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;

- methyl [5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
(2-benzoyl-6-chloro-1H-indol-3-yl)-*N,N*-dimethylacetamide;
5 (2-benzoyl-6-chloro-1H-indol-3-yl)-*N*-methylacetamide;
(2-benzoyl-6-chloro-1H-indol-3-yl)acetamide;
(2-benzoyl-6-chloro-1H-indol-3-yl)-*N*-methoxy-*N*-methylacetamide;
2-(2-benzoyl-6-chloro-1H-indol-3-yl)-1-piperidino-1-ethanone;
2-(2-benzoyl-6-chloro-1H-indol-3-yl)-1-(4-methyl-1-piperazinyl)-1-ethanone;
10 (2-benzoyl-6-chloro-1H-indol-3-yl)-*N*-(2-cyanoethyl)acetamide;
(2-benzoyl-6-chloro-1H-indol-3-yl)-*N*-(2-hydroxyethyl)acetamide;
2-(2-benzoyl-6-chloro-1H-indol-3-yl)-1-morpholino-1-ethanone;
[2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2-furylcarbonyl)-1H-indol-3-yl]acetic acid;
15 [6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 methyl [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
25 [5-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 methyl [2-(4-*tert*-butylpyridine-2-carbonyl)-6-chloro-1H-indol-3-yl]acetate;
[2-(4-*tert*-butylpyridine-2-carbonyl)-6-chloro-1H-indol-3-yl]acetic acid;

- methyl [2-(4-*tert*-butylpyridine-2-carbonyl)-5-chloro-1H-indol-3-yl]acetate;
[2-(4-*tert*-butylpyridine-2-carbonyl)-5-chloro-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 methyl [5-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
10 [5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
15 methyl [5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
20 [5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(pyridine-3-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(pyridine-3-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(pyridine-4-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(pyridine-4-carbonyl)-1H-indol-3-yl]acetic acid;
25 methyl [6-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[5-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(3,4-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
30 [5-chloro-2-(3,4-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;

- [5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetate;
5 [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 methyl [5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-ethoxy-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
15 [6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 methyl [6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
25 [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 methyl [6-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- methyl [5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-ethyl-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-ethyl-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 methyl [6-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetate;
10 [2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetate;
[2-(4-methyl-2-pyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
15 methyl [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetate;
[2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
methyl [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
20 [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl (2-benzoyl-1H-indol-3-yl)acetate;
(2-benzoyl-1H-indol-3-yl)acetic acid;
25 methyl [2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetate;
[2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-methyl-1H-indol-3-yl]acetic acid;
methyl [6-methoxy-2-(4-chlorobenzoyl)-1H-indol-3-yl] acetate;
[6-methoxy-2-(4-chlorobenzoyl)-1H-indol-3-yl] acetic acid;
30 [2-(4-chlorobenzoyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-5-ethyl-1H-indol-3-yl]acetate;

- [2-(4-chlorobenzoyl)-5-ethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-5-methoxy-1H-indol-3-yl]acetate;
[2-(4-chlorobenzoyl)-5-methoxy-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-5-isopropyl-1H-indol-3-yl]acetate;
5 [2-(4-chlorobenzoyl)-5-isopropyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-5-trifluoromethoxy-1H-indol-3-yl] acetate;
[2-(4-chlorobenzoyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
10 methyl [6-chloro-2-(2-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(2-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-benzyloxybenzoyl)-1H-indol-3-yl]acetate;
15 [6-chloro-2-(3-benzyloxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-hydroxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-hydroxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-benzyloxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-benzyloxybenzoyl)-1H-indol-3-yl]acetic acid;
20 methyl [6-chloro-2-(4-hydroxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-hydroxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-isopropoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-isopropoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-phenylbenzoyl)-1H-indol-3-yl]acetate;
25 [6-chloro-2-(4-phenylbenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
30 methyl [5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;

- methyl [6-chloro-2-(4-nitrobenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-nitrobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[(4-methylsulfonyl)benzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[(4-methylsulfonyl)benzoyl]-1H-indol-3-yl]acetic acid;
5 methyl [6-chloro-2-[4-(methylsulfonylamino)benzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(methylsulfonylamino)benzoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2,4-dichlorobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chloro-3-fluorobenzoyl)-1H-indol-3-yl]acetate;
10 [6-chloro-2-(4-chloro-3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-cyanobenzoyl)-1H-indol-3-yl]acetate;
methyl [6-chloro-2-[4-bromobenzoyl]-1H-indol-3-yl]acetate;
methyl [6-chloro-2-[4-(2-thienyl)benzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(2-thienyl)benzoyl]-1H-indol-3-yl]acetic acid;
15 methyl [6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(3-pyridyl)benzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(3-pyridyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(2-thiazolyl)benzoyl]-1H-indol-3-yl]acetate;
20 [6-chloro-2-[4-(2-thiazolyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-bromobenzoyl)-1H-indol-3-yl]acetate;
methyl [6-chloro-2-[3-(2-furyl)benzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[3-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl *dl*-2-[6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]propionate;
25 *dl*-2-[2-(4-chlorobenzoyl)-6-chloro-1H-indol-3-yl]propionic acid;
methyl [5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
30 methyl [5-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetic acid;

- methyl [6-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methyl-1,2,3-thiadiazole-5-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methyl-1,2,3-thiadiazole-5-carbonyl)-1H-indol-3-yl]acetic acid;
5 methyl [6-chloro-2-(4-methyl-1,2,3-thiadiazole-5-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methyl-1,2,3-thiadiazole-5-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
10 [6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2-thienyl)carbonylindol-3-yl]acetic acid;
methyl [6-chloro-2-[3-(1-hydroxy-1-methylethyl)-2-furoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[3-(1-hydroxy-1-methylethyl)-2-furoyl]-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetate;
15 [6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetate;
methyl [5-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 methyl [5-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
25 [5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(1-methylpyrrole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(1-methylpyrrole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetic acid;
30 methyl [5-chloro-2-(thiazole-5-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(thiazole-5-carbonyl)-1H-indol-3-yl]acetic acid;

- methyl [6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-[3-(ethoxycarbonyl)isoxazole-5-carbonyl]-1H-indol-3-yl]acetate;
[5-chloro-2-[3-(carboxy)isoxazole-5-carbonyl]-1H-indol-3-yl]acetic acid;
- 5 methyl [6-chloro-2-cyclopropanecarbonyl-1H-indol-3-yl]acetate;
[6-chloro-2-cyclopropanecarbonyl-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-cyclobutanecarbonyl-1H-indol-3-yl]acetate;
[6-chloro-2-cyclobutanecarbonyl-1H-indol-3-yl]acetic acid;
methyl [5-(*tert*-butyl)-2-(4-chlorobenzoyl)-1H-indol-3-yl] acetate;
- 10 [5-(*tert*-butyl)-2-(4-chlorobenzoyl)-1H-indol-3-yl] acetic acid;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-*N,N*-dimethylacetamide;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-*N*-methylacetamide;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-*N*-(2-hydroxyethyl)acetamid
e;
- 15 [5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-*N*-methoxyacetamide;
2-[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-1-piperazinyl-1-ethanone;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-*N*-(2-aminoethyl)acetamide;
2-[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-1-(3-amino-1-pyrrolidinyl
)-1-ethanone;
- 20 [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
methyl [6-chloro-5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
- 25 [6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2-nitrobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2,4-dimethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-difluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2,5-dimethoxybenzoyl)-1H-indol-3-yl]acetic acid;
- 30 methyl [5-acetyl-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate;
[5-acetyl-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;

- methyl [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetate;
methyl [6-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-fluoro-2-(4-chlorobenzoyl)-1H-indol-3-yl] acetate;
5 [6-fluoro-2-(4-chlorobenzoyl) -1H-indol-3-yl]acetic acid;
[2-(4-methylpyridine-2-carbonyl)-5-methylthio-1H-indol-3-yl]acetic acid;
[2-(4-methylpyridine-2-carbonyl)-5-methylthio-1H-indol-3-yl]acetic acid, and a salt thereof.

Preferred individual compounds of this invention are:

- 10 ethyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
(2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid, sodium salt;
[6-chloro-2-(2-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
15 [6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
20 [6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
25 [6-chloro-2-(3,4-dichlorobenzoyl)-1H-indol-3-yl]acetic acid;
(2-benzoyl-4-chloro-1H-indol-3-yl)acetic acid;
[5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
30 [2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
[2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;

- [5-methoxy-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
(2-benzoyl-7-chloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-4,5-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-4,6-dichloro-1H-indol-3-yl)acetic acid;
5 (2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
dl-2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
less polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
more polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 [6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
15 methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 methyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
[2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2-furylcarbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
25 [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 [5-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- [5-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-*tert*-butylpyridine-2-carbonyl)-6-chloro-1H-indol-3-yl]acetic acid;
[2-(4-*tert*-butylpyridine-2-carbonyl)-5-chloro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 [5-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
10 [5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-4-carbonyl)-1H-indol-3-yl]acetic acid;
15 [6-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3,4-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-ethoxy-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
25 [6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- [5-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-ethyl-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 [6-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
[5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-methyl-2-pyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
10 [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
[6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
(2-benzoyl-1H-indol-3-yl)acetic acid;
15 [2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-methyl-1H-indol-3-yl]acetic acid;
[6-methoxy-2-(4-chlorobenzoyl)-1H-indol-3-yl] acetic acid;
[2-(4-chlorobenzoyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-ethyl-1H-indol-3-yl]acetic acid;
20 [2-(4-chlorobenzoyl)-5-methoxy-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-isopropyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
25 [6-chloro-2-(3-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-benzyloxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-hydroxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-benzyloxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-hydroxybenzoyl)-1H-indol-3-yl]acetic acid;
30 [6-chloro-2-(4-isopropoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-phenylbenzoyl)-1H-indol-3-yl]acetic acid;

- [6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-nitrobenzoyl)-1H-indol-3-yl]acetic acid;
5 [6-chloro-2-[(4-methylsulfonyl)benzoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-[4-(methylsulfonylamino)benzoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2,4-dichlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chloro-3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
10 methyl [6-chloro-2-(4-cyanobenzoyl)-1H-indol-3-yl]acetate;
methyl [6-chloro-2-[4-bromobenzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(2-thienyl)benzoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-[4-(3-pyridyl)benzoyl]-1H-indol-3-yl]acetic acid;
15 [6-chloro-2-[4-(2-thiazolyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-bromobenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-[3-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
dl-2-[2-(4-chlorobenzoyl)-6-chloro-1H-indol-3-yl]propionic acid;
[5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
20 [6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methyl-1,2,3-thiadiazole-5-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methyl-1,2,3-thiadiazole-5-carbonyl)-1H-indol-3-yl]acetic acid;
25 [5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2-thienyl)carbonylindol-3-yl]acetic acid;
[6-chloro-2-[3-(1-hydroxy-1-methylethyl)-2-furoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetic acid;
30 [6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetate;

- [5-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 [5-chloro-2-(1-methylpyrrole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(thiazole-5-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-[3-(carboxy)isoxazole-5-carbonyl]-1H-indol-3-yl]acetic acid;
10 [6-chloro-2-cyclopropanecarbonyl-1H-indol-3-yl]acetic acid;
[6-chloro-2-cyclobutanecarbonyl-1H-indol-3-yl]acetic acid;
[5-(*tert*-butyl)-2-(4-chlorobenzoyl)-1H-indol-3-yl] acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
[6-chloro-5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
15 methyl [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-ethyl-3-fluoro-pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2-nitrobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2,4-dimethoxybenzoyl)-1H-indol-3-yl]acetic acid;
20 [6-chloro-2-(4-difluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2,5-dimethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[5-acetyl-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid, and a salt thereof.

Preferred individual compounds of this invention are:

- ethyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
25 (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
[6-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate;
30 [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;

- [6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
5 [6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
(2-benzoyl-4-chloro-1H-indol-3-yl)acetic acid;
[5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
10 [2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
[2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
(2-benzoyl-4,5-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-4,6-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
15 *dl*-2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
less polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
more polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
25 [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
30 [6-chloro-2-(2-furylcarbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]acetic acid;

- methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 methyl [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
10 [6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-*tert*-butylpyridine-2-carbonyl)-6-chloro-1H-indol-3-yl]acetate;
[2-(4-*tert*-butylpyridine-2-carbonyl)-6-chloro-1H-indol-3-yl]acetic acid;
15 methyl [6-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
20 [5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
25 methyl [5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
30 [5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;

- [5-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
5 [6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 methyl [6-chloro-2-(3,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
15 [6-chloro-2-(3-ethoxy-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 methyl [5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
25 [5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 methyl [5-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- methyl [6-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 methyl [5-ethyl-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-ethyl-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
10 [5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
15 methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetate;
[2-(4-methyl-2-pyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
methyl [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
20 [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetate;
[2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetic acid;
25 [2-(4-chlorobenzoyl)-5-methyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-5-trifluoromethoxy-1H-indol-3-yl] acetate;
[2-(4-chlorobenzoyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
30 methyl [6-chloro-2-(3-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-methoxybenzoyl)-1H-indol-3-yl]acetic acid;

- methyl [6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
5 methyl [5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
10 [5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetic acid;
15 methyl [5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetate;
20 [6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetate;
25 [5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
30 [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

methyl [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetate, and a salt thereof.

Preferred individual compounds of this invention are:

- ethyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
(2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
- 5 [6-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
- 10 [6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
- 15 [6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
(2-benzoyl-4-chloro-1H-indol-3-yl)acetic acid;
[5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
- 20 [2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
[2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
(2-benzoyl-4,5-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-4,6-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
- 25 *dl*-2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
less polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
more polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 30 methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- [6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 [6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
[6-chloro-2-(2-furylcarbonyl)-1H-indol-3-yl]acetic acid;
10 [6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
15 [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-*tert*-butylpyridine-2-carbonyl)-6-chloro-1H-indol-3-yl]acetic acid;
20 [6-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
25 [5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 [6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- [5-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-ethoxy-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 [6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
15 [5-ethyl-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
[5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 [2-(4-methyl-2-pyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
[6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetic acid;
25 [2-(4-chlorobenzoyl)-5-methyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
30 [5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;

- [6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetic acid;
5 [5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 [5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
15 2-{6-chloro-2-[(4-ethyl-3-fluoro-2-pyridinyl)carbonyl]-1H-indol-3-yl}acetic acid;
methyl [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetate, and a salt thereof.

Preferred individual compounds of this invention are:

- (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
[6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]acetic acid;
20 [6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
25 [2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
30 [5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;

- [2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
(2-benzoyl-4,5-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-4,6-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
- 5 [6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 10 [5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 15 methyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
[6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
- 20 [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 25 methyl [6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
- 30 [5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;

- [6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
5 [5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 methyl [5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetate;
15 [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-ethoxy-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 methyl [6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
25 [6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 methyl [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- methyl [5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 methyl [2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetate;
10 [2-(4-methyl-2-pyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
methyl [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
15 methyl [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-methyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
20 methyl [6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
25 [5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
30 methyl [6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;

- methyl [5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 methyl [6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetate;
10 [5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
15 methyl [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetate, and a salt thereof.

Preferred individual compounds of this invention are:

- (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
[6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
20 methyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
25 [2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
30 [2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
[2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;

- (2-benzoyl-4,5-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-4,6-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 [6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
15 [6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
25 [6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 [5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-ethoxy-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 [5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 [5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
[5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-methyl-2-pyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
15 [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
20 [5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
25 methyl [5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetic acid;
30 [6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;

- [6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetate;
[6-chloro-5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
5 [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid,
and a salt thereof.

Preferred individual compounds of this invention are:

- (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
[6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
10 [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
15 [6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 [5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
25 [6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 methyl [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;

- methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 methyl [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
10 [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
15 [6-chloro-5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
20 [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
25 [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid (cj-020,099);
methyl [6-chloro-5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate,
and a salt thereof.

Most preferred individual compounds are:

- (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
30 [6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;

- [5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
- 5 [6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 10 [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 15 [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 20 [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 25 [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid,
and a salt thereof.


- 30 Preferred pharmaceutical compositions of this invention comprise those compounds of the formula (I), wherein the compound is as defined above.

Most preferred individual compounds to be contained in the pharmaceutical compositions are:

- (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
- [6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
- 5 [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
- [5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
- [5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
- [2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
- (2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
- 10 [6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- [6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- [6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- [5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 15 [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- [6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- [6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 20 [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
- [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
- [5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- [5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 25 [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- [5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
- [6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
- [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
- [6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 30 [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

[5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
 [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid,
 and a salt thereof.

Also, the present invention provides a process for preparing a compound of the
 5 formula:



7-IV

Q is selected from the following:

- (a) phenyl optionally substituted with one, two or three substituents independently selected from
 - 10 (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,
 - 15 (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
 - 20 (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
 - 25 (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋

4 alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

- 5 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- 10 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- 15 (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and
- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

20 **R² and R³** are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;


X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; and

25

n is 0, 1, 2, 3 or 4,


which process comprises the steps of:

- i) reacting a compound of the formula:



7-II

wherein B is a suitable protecting group; R^5 is C_{1-6} alkyl; X and n are as defined above, with a compound of the formula:



5 wherein E is halo and Q are as defined above, with a first base and a suitable solvent;

ii) reacting the product of step i) with a second base.

iii) reacting the product of step ii) with an acid.

Preferred process of the above mentioned process is a process, wherein said first
10 base is potassium carbonate, potassium bicarbonate, sodium bicarbonate, sodium carbonate or cesium carbonate.

Preferred process of the above mentioned process is a process, wherein said first base is potassium carbonate.

Preferred process of the above mentioned process is a process, wherein said
15 second base is aqueous sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium pentoxide (followed by water), sodium methoxide (followed by water) or potassium *t*-butoxide (followed by water).

Preferred process of the above mentioned process is a process, wherein said second base is sodium hydroxide.

20 Preferred process of the above mentioned process is a process, wherein said acid is aqueous hydrochloric acid, hydrobromic acid, sulfuric acid or ammonium chloride.


Preferred process of the above mentioned process is a process, wherein said acid is aqueous hydrochloric acid.

Preferred process of the above mentioned process is a process, wherein said
25 solvent is *N,N*-dimethylacetamide, *N,N*-dimethylformamide, methyl ethyl ketone, acetone, or tetrahydrofuran.

Preferred process of the above mentioned process is a process, wherein said

solvent is *N,N*-dimethylaetamide.

Also, the present invention provides a process for prepaing a compound of the formula:



7-IV

5 wherein

Q is selected from the following:

- (a) phenyl optionally substituted with one, two or three substituents independently selected from
 - (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,
 - (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
 - (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
 - (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-

substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino and CN,


- (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- (d) C_{3-7} cycloalkyl optionally substituted with one or two substituents independently selected from OH, C_{1-4} alkyl, halo and halo-substituted C_{1-4} alkyl; and
- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R^2 and R^3 are independently H, OH, C_{1-4} alkoxy, C_{1-4} alkyl or C_{1-4} alkyl substituted with halo, OH, C_{1-4} alkoxy, NH_2 or CN;

- X is independently selected from halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino, CN, HO- (C_{1-4}) alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $-NH_2S(O)_2NR^2NR^3$, acetyl, $-COOH$, $-C(O)O-C_{1-4}$ alkyl, C_{1-4} alkylsulfonylamino and C_{3-7} cycloalkyl; and

- n is 0, 1, 2, 3 or 4,

which process comprises reacting a compound of the formula:



7-VII

wherein R^5 is C_{1-6} alkyl; Q, X and n are as defined as before, with a base in a suitable solvent.


Preferred process of the above mentioned process is a process, wherein said base is sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate,
5 sodium bicarbonate, potassium bicarbonate, sodium *t*-pentoxide, sodium methoxide, sodium ethoxide or potassium *t*-butoxide.

Preferred process of the above mentioned process is a process, wherein said base is sodium hydroxide.

Preferred process of the above mentioned process is a process, wherein said
10 solvent is an aqueous mixture of methanol, ethanol, isopropyl alcohol or tetrahydrofuran.

Preferred process of the above mentioned process is a process, wherein said solvent is methanol containing water.

Also, the present invention provides a process for prepaing a compound of the
15 formula:



7-VII

wherein

Q is selected from the following:

- 20 (a) phenyl optionally substituted with one, two or three substituents independently selected from
- (a-1) halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino, CN, HO- (C_{1-4}) alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $-NH_2S(O)_2NR^2R^3$, acetyl,
25 -COOH, -C(O)O- C_{1-4} alkyl, C_{1-4} alkylsulfonylamino and C_{3-7} cycloalkyl,

- 5 (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 10 (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 15 (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 20 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- 25 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- 30 (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and
- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);


R² and R³ are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted

with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; R⁵ is C₁₋₆ alkyl; and

n is 0, 1, 2, 3 or 4,

which process comprises reacting a compound of the formula:



7-VI

10 wherein B, Q, X, n and R⁵ are as defined above with a base in a suitable solvent.


Preferred process of the above mentioned process is a process, wherein said base is 1,8-diazabicyclo[5.4.0]undec-7-ene, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,1,3,3-tetramethylguanidine, sodium *t*-pentoxide, sodium methoxide, or potassium *t*-butoxide.

15 Preferred process of the above mentioned process is a process, wherein said base is 1,8-diazabicyclo[5.4.0]undec-7-ene or potassium *t*-butoxide.

Preferred process of the above mentioned process is a process, wherein said solvent is *N,N*-dimethylacetamide, *N,N*-dimethylformamide, methyl ethyl ketone, acetone, or tetrahydrofuran.

20 Preferred process of the above mentioned process is a process, wherein said solvent is *N,N*-dimethylacetamide.

Also, the present invention provides a process for prepaing a compound of the formula:



wherein **B** is a suitable protecting group;

Q is selected from the following:

- 5 (a) phenyl optionally substituted with one, two or three substituents independently selected from
 - (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,
 - (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
 - (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
 - (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

- (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- 5 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- 10 (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and
- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);
- 15


R² and R³ are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; R⁵ is C₁₋₆ alkyl; and

20

n is 0, 1, 2, 3 or 4,

which process comprises reacting a compound of the formula:



7-V

wherein B, Q, X, n and R⁵ are as defined above, with a base in the presence of a solvent.


Preferred process of the above mentioned process is a process, wherein said base is potassium carbonate, potassium bicarbonate, sodium bicarbonate, sodium carbonate,
5 or cesium carbonate.

Preferred process of the above mentioned process is a process, wherein said base is potassium carbonate.

Preferred process of the above mentioned process is a process, wherein said solvent is *N,N*-dimethylacetamide, *N,N*-dimethylformamide, methyl ethyl ketone,
10 acetone, or tetrahydrofuran.

Preferred process of the above mentioned process is a process, wherein said solvent is *N,N*-dimethylacetamide.

Also, the present invention provides a process for preparing a compound of the formula:



15

wherein B is a suitable protecting group;

Q is selected from the following:

(a) phenyl optionally substituted with one, two or three substituents
20 independently selected from

(a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl,
25 -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,


- 5 (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 10 (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 15 (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 20 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- 25 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- 30 (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and
- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R² and R³ are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted


with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; R⁵ is C₁₋₆ alkyl; and n is 0, 1, 2, 3 or 4,

which comprises reacting a compound of the formula:



wherein B, X, n and R⁵ are as defined above, with a compound of the formula:



wherein E is halo and Q are as defined above, with a base in the presence of a solvent.


Preferred process of the above mentioned process is a process, wherein said base is potassium carbonate, potassium bicarbonate, sodium bicarbonate, sodium carbonate, or cesium carbonate.

Preferred process of the above mentioned process is a process, wherein said base is potassium carbonate.

Preferred process of the above mentioned process is a process, wherein said solvent is *N,N*-dimethylacetamide, *N,N*-dimethylformamide, methyl ethyl ketone, acetone or tetrahydrofuran.

Preferred process of the above mentioned process is a process, wherein said solvent is *N,N*-dimethylacetamide.

Also, the present invention provides a process for preparing a compound of the formula:



wherein

Q is selected from the following:

- (a) phenyl optionally substituted with one, two or three substituents independently selected from
 - (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,
 - (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
 - (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
 - (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being

optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),

- (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and
- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R¹ is hydrogen, C₁₋₄ alkyl or halo;


- 15 R² and R³ are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

R⁵ is C₁₋₆ alkyl;

- X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; and

n is 0, 1, 2, 3 or 4,

which process comprises treating a compound of the formula (X):




(X)

25

wherein R¹, R⁵, X, Q and n are as defined herein before, and B is a suitable protecting group, in the presence of a suitable base to obtain a compound of the formula (XII).

Also, the present invention provides a process for preparing a compound of the formula:



wherein

5 Q is selected from the following:

(a) phenyl optionally substituted with one, two or three substituents independently selected from

(a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,

(a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

- (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- 5 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- 10 (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and
- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);
- 15

R¹ is hydrogen, C₁₋₄ alkyl or halo;


R² and R³ are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

R⁵ is C₁₋₆ alkyl;

- 20 X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; and


- 25 n is 0, 1, 2, 3 or 4,

which process comprises reacting a compound of the formula (IX):



(IX)

wherein R^1 , R^5 , X, and n are as defined above, and B is a suitable protecting group, with a compound of the formula (XI):



(XI)

- 5 wherein E is halo and Q is as defined as before, in the presence of a suitable base at a temperature of $-40\text{ }^{\circ}\text{C}$ to $200\text{ }^{\circ}\text{C}$ to obtain a compound of the formula (XII).

Preferred process of the above mentioned process is a process, wherein the reaction is carried out at a temperature of $0\text{ }^{\circ}\text{C}$ to $100\text{ }^{\circ}\text{C}$

- 10 Preferred process of the above mentioned process is a process, wherein the suitable base is potassium carbonate, cesium carbonate, sodium carbonate, sodium tert-butoxide, potassium tert-butoxide, sodium hydride, potassium hydride or potassium fluoride.

- Preferred process of the above mentioned process is a process, wherein the reaction is firstly carried out in the presence of a base for 2 minutes to a day; and then, another base is added to the reaction mixture.

Preferred process of the above mentioned process is a process, wherein the reaction is firstly carried out for 30 minutes to 8 hours.

- 20 Preferred process of the above mentioned process is a process, wherein the suitable protecting group is methoxycarbonyl, ethoxycarbonyl, tert-butoxycarbonyl, benzyloxycarbonyl, phenylsulfonyl, p-toluenesulfonyl, methanesulfonyl or trifluoromethanesulfonyl.

- Preferred process of the above mentioned process is a process, wherein the suitable protecting group is phenylsulfonyl, p-toluenesulfonyl, methanesulfonyl or trifluoromethanesulfonyl.
- 25

Preferred process of the above mentioned process is a process, wherein the first base is selected from sodium *tert*-butoxide, potassium *tert*-butoxide, sodium carbonate, potassium carbonate, cesium carbonate, sodium hydride, potassium hydride, sodium carbonate, potassium carbonate, cesium carbonate, potassium fluoride, 1,8-diazabicyclo[5.4.0]undec-7-ene, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,4-diazabicyclo[2.2.2]octane, pyridine, pyrrolidine, triethylamine, diisopropylamine, diisopropylethylamine and diethylisopropylamine; and the second base is selected from sodium *tert*-butoxide, potassium *tert*-butoxide, sodium carbonate, potassium carbonate, cesium carbonate, sodium hydride, potassium hydride, sodium carbonate, potassium carbonate, cesium carbonate, potassium fluoride, 1,8-diazabicyclo[5.4.0]undec-7-ene, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,4-diazabicyclo[2.2.2]octane, pyridine, pyrrolidine, triethylamine, diisopropylamine, diisopropylethylamine and diethylisopropylamine.

Preferred process of the above mentioned process is a process, wherein the first base is selected from potassium carbonate, cesium carbonate, sodium hydride and potassium fluoride; and the second base is selected from 1,8-diazabicyclo[5.4.0]undec-7-ene, cesium carbonate, pyrrolidine, diisopropylamine, triethylamine, diethylisopropylamine and diisopropylethylamine.


Preferred process of the above mentioned process is a process, wherein the first base is potassium carbonate, cesium carbonate or potassium fluoride; and the second base is 1,8-diazabicyclo[5.4.0]undec-7-ene, potassium *tert*-butoxide or cesium carbonate.

Preferred process of the above mentioned process is a process, wherein the combination of the first base and the second base (first base/second base) is selected from potassium carbonate/1,8-diazabicyclo[5.4.0]undec-7-ene, potassium carbonate/cesium carbonate, cesium carbonate/potassium *tert*-butoxide, cesium carbonate/1,8-diazabicyclo[5.4.0]undec-7-ene and potassium fluoride/1,8-diazabicyclo[5.4.0]undec-7-ene and potassium fluoride/cesium carbonate.

Preferred process of the above mentioned process is a process, wherein the combination of the first base and the second base (first base/second base) is selected

from potassium carbonate/1,8-diazabicyclo[5.4.0]undec-7-ene, potassium carbonate/cesium carbonate and cesium carbonate/potassium tert-butoxide.

Also, the present invention provides a process for preparing a compound of the formula:



wherein

Q is selected from the following:

- (a) phenyl optionally substituted with one, two or three substituents independently selected from

(a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,

(a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-

substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

- 5 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- 10 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- 15 (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and
- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R¹ is hydrogen, C₁₋₄ alkyl or halo;


20 R² and R³ are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; and

25

n is 0, 1, 2, 3 or 4,


which process comprises treating a compound of the formula (X):



(X)

wherein R^1 , R^5 , X, Q and n are as defined here before, with a suitable base under hydrolyzing conditions to obtain the compound of formula (VIII).

This invention also provides a process for preparing a compound of the formula
5 (VIII):



(VIII)

wherein

Q is selected from the following:

- (a) phenyl optionally substituted with one, two or three substituents
10 independently selected from
 - (a-1) halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino, CN, HO- (C_{1-4}) alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $-NH_2S(O)_2NR^2R^3$, acetyl,
15 $-COOH$, $-C(O)O-C_{1-4}$ alkyl, C_{1-4} alkylsulfonylamino and C_{3-7} cycloalkyl,
 - (a-2) aryl or $-O-(CH_2)_n$ -aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino and CN,
 - (a-3) 5-membered monocyclic aromatic group optionally substituted with
20 one, two or three substituents independently selected from halo, C_1 .

4 alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

5 (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

10 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),

15 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);

20 (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and

(e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R¹ is hydrogen, C₁₋₄ alkyl or halo;


25 **R² and R³** are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; and

30

n is 0, 1, 2, 3 or 4,

which process comprises hydrolyzing a compound of the formula (XII):




wherein R^5 is C_{1-6} alkyl, R^1 , X, Q and n are as defined herein before.

5

General Synthesis


A compound of general formula (I) may be prepared by any synthetic procedure applicable to structure-related compounds known to those skilled in the art. The following representative examples as described hereinafter are illustrative and are not meant to limit the scope of the invention in anyway. Unless otherwise stated, Q, X, Z,

10 R^1 , and n are as defined above.



Scheme 1:

15 In one embodiment, for example, a compound of the formula (VI) may be prepared according to the reaction sequences depicted in Scheme 1. (Compound (VI) corresponds to a compound (I) wherein R^1 is H, and Z is OH.)




Scheme 1

In brief, a compound of formula (III) is subjected to oxidative homolytic malonylation (for leading references see J. M. Muchowski et al; Can. J. Chem., **70**, 1838, 1992 and E. Baciocchi et al; J. Org. Chem., **58**, 7610, 1993). In one example, a compound of the formula (III) is reacted with a suitable malonyl radical generated from a compound of formula $\text{C(R}^4\text{)H(CO}_2\text{R}^5\text{)}_2$, wherein R^4 is hydrogen or halogen, preferably chloro, and R^5 is C_{1-6} alkyl, and a manganese(III) agent, preferably manganese (III) triacetate. The manganese(III) agent is usually used in stoichiometric amounts but, alternatively, may be made catalytic by use of a suitable reoxidizing agent such as sodium persulfate, usually in the presence of a co-catalyst such as, a silver(I) salt such as silver nitrate. A preferred reaction solvent is acetic acid; however, acetic acid-acetic anhydride or other protic solvents such as propionic acid can be used. The reaction is preferably conducted in the presence of sodium acetate or potassium acetate, but, may be conducted in solvent alone. Reaction temperatures are generally in the range of room temperature (e.g., 25°C) to reflux temperature of solvent, preferably 60°C .

to 100 °C, but if necessary, lower or higher temperature can be employed. Reaction times are, in general, from one hour to a day, preferably from 4 to 16 hours, however shorter or longer reaction times, if necessary, can be employed. In the immediate instance, the α -acetoxo compounds of formula (IV) is usually obtained as the major product. Compounds of formula (IV) can readily be transformed to compounds of formula (V) by reduction with a suitable reducing agent, for example, a trialkylsilane, sodium α -(dimethylamino)naphthalenide, lithium in liquid ammonia, sodium naphthalenide, preferably triethylsilane in a suitable protic solvent, notably, trifluoroacetic acid. Alternatively, the reaction can be conducted in a reaction inert co-solvent such as dichloromethane or 1,2-dichloroethane. Reaction temperatures are generally in the range of room temperature to reflux temperature of solvent, preferably 15 to 100 °C, but if necessary, lower or higher temperature can be employed. Reaction times are, in general, from several minutes to a day, preferably from 20 minutes to 5 hours, however shorter or longer reaction times, if necessary, can be employed. Alternatively, a compound of formula (V) may be obtained directly from a compound of formula (III) from a malonyl radical generated from (i) a suitable monohalomalonate, preferably, bromomalonate, mediated by aerial oxidation of a trialkylborane such as triethylborane (see B. Giese; In Radicals in organic synthesis: formation of carbon-carbon bonds. Pergamon Press, Oxford. pp. 86-89, 1986, and P. G. Allies and P. B. Brindley; J. Chem. Soc. (B), 1126, 1960) or, (ii) a malonic ester in the presence of a cerium(IV) salt such as cerium (IV) ammonium nitrate (for example, see E. Baciocchi et al; Tetrahedron Lett, 2763, 1986). A compound of formula (V) may be readily transformed to a compound of formula (VI) by subjection to standard saponification / decarboxylation conditions.

Scheme 2:

Alternatively, as depicted in Scheme 2, a compound of the formula (VIII) (a compound (I) wherein Z is OH), wherein R¹ is C₁₋₄ alkyl, may be prepared in an analogous manner to that of a compound of formula (VI) employing appropriate reaction conditions as described by illustration herein above from a suitable monoalkylmalonate, wherein R¹ is C₁₋₄ alkyl, W is hydrogen or a halogen, preferably bromide, and R⁵ is C₁₋₆ alkyl, from a compound of formula (III).

[R¹ is not hydrogen]


[W = H or halo]

[R⁵ = C₁₋₆alkyl]**Scheme 2**

In Scheme 2, for example, the oxidant is manganese (III) agent such as manganese (III) triacetate, or Cerium (IV) agent such as ammonium Cerium (IV) nitrate and Cerium (IV) sulfate.

Scheme 3:

In another embodiment, a compound of formula (VIII) is readily accessible from the appropriate 2-aminocinnamic acid ester (IX) wherein B is a suitable protecting group, for example, methoxycarbonyl, ethoxycarbonyl, tert-butoxycarbonyl (Boc), benzyloxycarbonyl, phenylsulfonyl, p-toluenesulfonyl, methanesulfonyl, trifluoromethanesulfonyl, methanesulfonyl or trifluoromethanesulfonyl (preferably phenylsulfonyl, p-toluenesulfonyl, methanesulfonyl or trifluoromethanesulfonyl).



Scheme 3


In Scheme 3, the requisite 2-aminocinnamic acid ester (IX) is reacted with a compound of formula (XI), wherein Q is as defined above and E is halogen, preferably, iodo, bromo or chloro, in the presence of a suitable base. A suitable base is, for example, an alkali or alkaline earth metal alkoxide, carbonate, fluoride or hydride, such as sodium *tert*-butoxide, potassium *tert*-butoxide, sodium carbonate, potassium carbonate, cesium carbonate, sodium hydride, potassium fluoride or potassium hydride. Preferred reaction inert solvents include, but are not limited to, acetone, methyl ethyl ketone, acetonitrile, N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), dimethylsulfoxide (DMSO), dioxane or tetrahydrofuran (THF). Reaction temperatures are preferably in the range of -40 °C to reflux temperature of solvent (for example 200 °C), usually in the range of 0 °C to 100 °C, but if necessary, lower or higher temperature can be employed. Reaction time is in general from 2 minutes to a

day, preferably from 30 minutes to 8 hours, however shorter or longer reaction times, if necessary, can be employed. When the reaction is, for example, conducted at room temperature (e.g., 25 °C) the intermediate indoline (X) can be isolated. Reaction at higher temperatures (e.g., 40 to 100 °C) can result in formation of indole (XII).

- 5 Usually the intermediate indoline (X) is not isolated but either (i) hydrolyzed with committant formation of the indole ring directly to a compound of formula (VIII) under standard conditions known to those skilled in the art, or (ii) transformed to a compound of formula (XII) by using a suitable base, for example, an alkali or alkaline earth metal carbonate such as sodium carbonate, potassium carbonate or cesium carbonate, or an
10 organic base such as 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), 1,5-diazabicyclo[4.3.0]non-5-ene (DBN), 1,4-diazabicyclo[2.2.2]octane (DABCO), pyridine, pyrrolidine, triethylamine, diisopropylamine, diisopropylethylamine, diethylisopropylamine, Hunig's base, potassium tert-butoxide, sodium tert-butoxide, or the like, or a suitable oxidant such as cerium (IV) ammonium nitrate (CAN),
15 manganese(IV) oxide, manganese(III) triacetate, copper (II) acetate / air, chloranil, 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ), N-methylmorpholine-N-oxide, or the like (for example, see H. Dumoulin et al; J. Heterocycl. Chem., **32**, 1703, 1995; H. Rapoport et al; Tetrahedron Lett., 5053, 1991; P. Martin et al; Helv. Chim. Acta, **77**, 111, 1994; Y. Kikugawa et al, J. Chem. Soc. Perkins Trans 1, **7**, 1401, 1984; A.
20 Goti et al; Tetrahedron Lett., 6567, 1996; L. S. Liebeskind et al; J. Org. Chem, **61**, 2594, 1996). Preferred reaction inert solvents include, but are not limited to, acetone, methyl ethyl ketone, acetonitrile, dioxane or tetrahydrofuran (THF). Reaction temperatures are preferably in the range of 0 °C to reflux temperature of solvent, usually in the range of 15 to 60 °C, but if necessary, lower or higher temperature can be
25 employed. Reaction time is in general from several minutes to a day, preferably from 30 minutes to 8 hours, however shorter or longer reaction times, if necessary, can be employed. A compound of formula (XII) may be readily hydrolyzed to a compound of formula (VIII) under standard conditions.

Scheme 4:

- 30 In another embodiment, a compound of formula (VIII), wherein Q, X, R¹ and n are as defined above, may be prepared as illustrated in Scheme 4.



Scheme 4

For example, treatment of a compound of formula (XIII), wherein R¹, R⁵, X and n are as defined above, with a trialkyltin hydride, e.g., tributyltin hydride usually in the presence of a radical initiator such as, 2,2'-azobisisobutyronitrile (AIBN), affords the intermediate 2-stannylindole (XIV) via an intramolecular radical cyclization as described in J. Am. Chem. Soc., **116**, 3127, (1994); T. Fukuyama et al. The intermediate (XIV) generated *in situ* is subsequently treated with an acyl halide, wherein Q and E are as defined above, in the presence of a suitable palladium catalyst according to Stille's procedure (for example see. J. K. Stille et al; J. Am. Chem. Soc., **109**, 813, 5478, (1987) and J. Am. Chem. Soc., **106**, 4833, (1984)) to afford indole (XII) which may be hydrolyzed to a compound of formula (VIII) by conventional


procedure.

Examples of the palladium catalyst are
 tetrakis(triphenylphosphine)palladium(0),
 dichlorobis(triphenylphosphine)palladium(II), bis(dibenzylideneacetone)palladium(0),
 5 benzyl(chloro)bis(triphenylphosphine)palladium(II),
 bis(acetonitrile)dichloropalladium(II).

Scheme 5:

In another embodiment, a compound of formula (VIII), wherein Q, X, R¹ and n are as defined above, may be prepared as illustrated in Scheme 5.

10




Scheme 5

For example, treatment of a compound (XV), wherein R¹, X and n are as defined above, is reacted with a compound of formula Q-C(O)-A affords a compound
 15 of formula (VIII), or a compound of formula (XVI) (for example see U.Pindur et al.,
Liebigs Ann. Chem., 601 (1991) and C.J.Moody et al., *J.Chem.Soc.Perkin Trans.I*,
 3249 (1988)) which may be hydrolyzed to a compound of formula (VIII) by
 conventional procedure (for example see E.B.Fray et al., *Tetrahedron*, **49**, 439 (1993)
 and U.Pindur et al., *J.Heterocycl.Chem.*, **29**, 145 (1992)). In a compound of formula
 20 A-C(O)-Q, A is defined such that the compound of A-C(O)-Q is, for example, an acyl
 halide, carboxylic acid, carboxylic acid anhydride, a mixed carboxylic sulfonic

anhydride, or the like. The reaction may be conducted in the presence or absence of catalyst, preferably in the presence of catalyst such as, boron trifluoride-diethyl ether, tin(IV) chloride, aluminum chloride, ferric chloride, zinc chloride, iodine, iron, or the like. Preferred reaction inert solvents include, but are not limited to, diethyl ether, 5 dichloromethane, 1,2-dichloroethane, carbon disulfide, nitrobenzene or nitromethane. Reaction temperatures are preferably in the range of -78 to 210 °C, usually in the range of -10 °C to reflux temperature of solvent, but if necessary, lower or higher temperature can be employed. Reaction time is in general from several minutes to a day, preferably from 30 minutes to 8 hours, however shorter or longer reaction times, if 10 necessary, can be employed.

Scheme 6:

Acetic acid compounds of formulae (VI) and (VIII) as described in the aforementioned schemes may be readily transformed to the corresponding amide, compounds of formulae (XVII) and (XVIII), or ester, compound of formula (XII), by 15 any conventional method known to those skilled in the art.




[R⁵ = C₁₋₆alkyl]

Scheme 6

As depicted in Scheme 6, compounds of formulae (XVII) and (XVIII) can be readily prepared by treating the requisite acetic acid compounds of formulae (VI) and (VIII) with an appropriate amine, wherein R², R³, Y and r are as described herein before, in the presence of a suitable coupling reagent such as, but not limited to, 1-(dimethylaminopropyl)-3-ethylcarbodiimide (WSC), N,N'-dicyclohexylcarbodiimide (DCC), carbonyldiimidazole, diethylphosphorocyanidate (DEPC), or the like. Preferred reaction inert solvents include, but are not limited to, acetone, acetonitrile, dichloromethane, 1,2-dichloroethane, N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), dimethylsulfoxide (DMSO), dioxane, tetrahydrofuran (THF) or pyridine. Reaction temperatures are preferably in the range of -40 to 150 °C, usually in the range of 15 °C to reflux temperature of solvent, but if

necessary, lower or higher temperature can be employed. Reaction time is in general from several minutes to a day, preferably from 30 minutes to 8 hours, however shorter or longer reaction times, if necessary, can be employed. The compounds of formulae (VI) and (VIII) can also be readily transformed to the corresponding ester by conventional methods.

Scheme 7



(wherein B is a suitable protecting group, R⁵ is C₁₋₆ alkyl, E is halo, Q, X and n is as defined above.)

In Scheme 7, the starting material of formula 7-I may be prepared according to methods familiar to those of ordinary skill in the art, including one or more synthetic procedures described in R. W. Carling, P. D. Leeson, K. Moore, J. D. Smith, C. R. Moyes, *J. Med. Chem.*, 1993, pages 3397-3408.

The compound of formula 7-II is prepared from a compound of formula 7-I by treatment with a base and an electrophile in a suitable solvent. Suitable bases include such as triethylamine, diisopropylethylamine, or pyridine optionally substituted by 1 to 3 (C₁-C₄)alkyl groups, preferably pyridine. Suitable electrophiles include methanesulfonyl chloride or anhydride, or phenylsulfonyl chloride wherein the phenyl moiety of said phenylsulfonyl optionally includes 1 or 2 substituents selected from halo, nitro, and (C₁-C₄)alkyl. Suitable solvents include dichloromethane, dichloroethane,

methyl *t*-butyl ether, diisopropyl ether or toluene, preferably dichloromethane. The temperature of the aforesaid reaction may range from about 0 °C to about 50 °C, preferably about room temperature (20-25 °C) for a period of about 1 to 30 hours, preferably about 18 hours.

5 The compound of formula 7-IV is prepared from a compound of formula 7-II by treatment with a first base and an alkylating agent of the formula 7-III in the presence of a solvent followed by reaction with a second base followed by reaction with an acid.. Suitable first bases include potassium carbonate, potassium bicarbonate, sodium bicarbonate, sodium carbonate or cesium carbonate, preferably
10 potassium carbonate. Suitable solvents include *N,N*-dimethylacetamide, *N,N*-dimethylformamide, methyl ethyl ketone, acetone or tetrahydrofuran, preferably *N,N*-dimethylacetamide. The aforesaid reaction is performed at a temperature ranging from about 0 °C to about 100 °C, preferably room temperature (20-25 °C), for a period of time of about 10 minutes to 5 hours, typically 15 minutes. Suitable second
15 bases include an aqueous solution of a base such as sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium *t*-pentoxide (followed by water), sodium methoxide (followed by water) or potassium *t*-butoxide (followed by water), preferably sodium hydroxide. The reaction with the second base is performed at a temperature ranging from about 20 °C to about 120 °C, preferably 100 °C, for a
20 period of time of about 1 hour to 24 hours, typically 8 hours. Suitable acids include aqueous hydrochloric acid, hydrobromic acid, sulfuric acid or ammonium chloride, preferably hydrochloric acid. The reaction with the acid is performed at a temperature ranging from about 0 °C to about 50 °C, preferably about 20 °C to about 25 °C, for a period of time of about 1/2 hour to about 6 hours, typically about 1 hour.

25 Alternatively, the conversion of the compound of formula 7-II to a compound of formula 7-IV can be accomplished stepwise. The compound of formula 7-V may be prepared from a compound of formula 7-II by treatment with a base and an alkylating agent of formula 7-III in the presence of a solvent. Suitable bases include potassium carbonate, potassium bicarbonate, sodium bicarbonate, sodium carbonate, or
30 cesium carbonate, preferably potassium carbonate. Suitable solvents include *N,N*-dimethylacetamide, *N,N*-dimethylformamide, methyl ethyl ketone, acetone or

tetrahydrofuran, preferably *N,N*-dimethylacetamide. The temperature for the aforesaid reaction may range from about 0 °C to about 50 °C, preferably room temperature (20-25 °C), for a period of time of about 10 minutes to 40 minutes, typically 30 minutes.

- 5 The compound of formula 7-VI is prepared from a compound of formula 7-V by reaction with a base in the presence of a solvent. Suitable bases include potassium carbonate, potassium bicarbonate, sodium bicarbonate, sodium carbonate or cesium carbonate, preferably potassium carbonate. Suitable solvents include *N,N*-dimethylacetamide, *N,N*-dimethylformamide, methyl ethyl ketone, acetone or
10 tetrahydrofuran, preferably *N,N*-dimethylacetamide. The temperature for the aforesaid reaction may range from about 0 °C to about 50 °C, preferably room temperature (20-25 °C), for a period of time of about 1 hour to 6 hours, preferably 4 hours.

- The compound of formula 7-VII is prepared from a compound of formula 7-VI
15 by reaction with a base in a suitable solvent. Suitable bases include 1,8-diazabicyclo[5.4.0]undec-7-ene, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,1,3,3-tetramethylguanidine, sodium *t*-pentoxide, sodium methoxide or potassium *t*-butoxide, preferably 1,8-diazabicyclo[5.4.0]undec-7-ene methoxide or potassium *t*-butoxide. Suitable solvents include *N,N*-dimethylacetamide, *N,N*-dimethylformamide, methyl
20 ethyl ketone, acetone or tetrahydrofuran, preferably *N,N*-dimethylacetamide. The temperature for the aforesaid reaction may range from about 0 °C to 100 °C, preferably room temperature (20-25 °C), for a period of 30 minutes to 5 hours, preferably 1 hour.

- The compound of formula 7-IV is prepared from a compound of formula 7-VII
25 by treatment with a base in a suitable solvent. Suitable bases include sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate, sodium *t*-pentoxide, sodium methoxide, sodium ethoxide or potassium *t*-butoxide, preferably sodium hydroxide. Suitable solvents include an aqueous mixture of methanol, ethanol, isopropyl alcohol or tetrahydrofuran,
30 preferably methanol, containing water. The temperature of the aforesaid reaction may range from about 10 °C to 100 °C, preferably room temperature (20-25 °C), for a

period of 12 to 48 hours, preferably 24 hours, to provide the carboxylate salt of compound of formula 7-IV which can then be treated with an acid to provide the compound of formula 7-IV.

The compound of formula 7-VI has asymmetric atoms and therefore exist in
5 different enantiomeric and diastereomeric forms. Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods known to those skilled in the art, for example, by chromatography or fractional crystallization. The use of all such isomers, including diastereoisomer mixtures and pure enantiomers, are considered to be part of the present
10 invention.

The starting materials in the aforementioned general syntheses may be obtained by conventional methods known to those skilled in the art. The preparation of such starting materials is described within the accompanying non-limiting examples which are provided for the purpose of illustration only. Alternatively, requisite starting
15 materials may be obtained by analogous procedures, or modifications thereof, to those described hereinafter.

The products which are addressed in the aforementioned general syntheses and illustrated in the experimental examples described herein after may be isolated by standard methods and purification can be achieved by conventional means known to
20 those skilled in the art, such as distillation, crystallization or chromatography techniques.

Certain compounds described herein contain one or more asymmetric centers and are capable of existing in various stereoisomeric forms. The present invention contemplates all such possible stereoisomers as well as their racemic and resolved,
25 enantiomerically pure forms and pharmaceutically acceptable salts thereof.

Certain compounds of the present invention are capable of forming addition salts with inorganic or organic acids. The pharmaceutically acceptable acid salts of the compounds of formula (I) are those which form non-toxic addition salts, such as, but not limited to, the hydrochloride, hydrobromide, sulfate or bisulfate, acetate,
30 benzoate, besylate, citrate, fumarate, glucuronate, hippurate, lactate, tartrate, saccharate, succinate, maleate, methanesulfonate, *p*-toluenesulfonate, phosphate and pamoate (i.e.,

4,4'-methylene-bis-(3-hydroxy-2-naphthoate)) salts. The pharmaceutically acceptable acid salts may be prepared by conventional techniques.

Certain compounds of the present invention are capable of forming pharmaceutically acceptable non-toxic cations. Pharmaceutically acceptable non-toxic cations of compounds of formula (I) may be prepared by conventional techniques by, for example, contacting said compound with a stoichiometric amount of an appropriate alkali or alkaline earth metal (sodium, potassium, calcium and magnesium) hydroxide or alkoxide in water or an appropriate organic solvent such as ethanol, isopropanol, mixtures thereof, or the like.

Also included within the scope of this invention are bioprecursors (also called pro-drugs) of the compounds of the formula (I). A bioprecursor of a compound of the formula (I) is a chemical derivative thereof which is readily converted back into the parent compound of the formula (I) in biological systems. In particular, a bioprecursor of a compound of the formula (I) is converted back to the parent compound of the formula (I) after the bioprecursor has been administered to, and absorbed by, a mammalian subject, e.g., a human subject. When the compounds of the formula (I) of this invention may form solvates such as hydrates, such solvates are included within the scope of this invention.

An example of prodrug of the compound of formula (I) is a compound of the formula (I), wherein the 1st position of indole ring is substituted with a group selected from hydroxymethyl, $-C(O)-C_{1-4}$ alkyl, $-C(O)-(NH_2)CH-(C_{1-4}$ alkyl), $-C(O)$ -phenyl, $-CH_2NHC(O)$ -aryl, $-CH_2-C_{1-4}$ alkyl-O-C(O)- C_{1-4} alkyl, $-C_{1-4}$ alkyl-pyridyl, $-C(O)CH_2NR_2$ and $-CH_2N(C_{1-4}$ alkyl)₂.

Another example of prodrug of the compound of formula (I) is a compound of the formula (I), wherein the carboxyl group is substituted with a group selected from C_{1-4} alkyl, $-CH_2-C_{1-4}$ alkyl-O-C(O)- C_{1-4} alkyl, $-CH_2-C_{1-4}$ alkyl-O-C(O)-N(C_{1-4} alkyl)₂, $-CH_2C(O)-N(C_{1-4}$ alkyl)₂, $-CH_2-C_{1-4}$ alkyl-O-C(O)-O- C_{1-4} alkyl, ethyl-OH and $-CH_2CO_2H$.

The compounds of the formula (I) of this invention can be administered via either the oral, parenteral or topical routes to mammals. In general, these compounds are most desirably administered to humans in doses ranging from 0.01 mg to 100 mg

per kg of body weight per day, although variations will necessarily occur depending upon the weight, sex and condition of the subject being treated, the disease state being treated and the particular route of administration chosen. However, a dosage level that is in the range of from 0.01 mg to 10 mg per kg of body weight per day, single or
5 divided dosage is most desirably employed in humans for the treatment of abovementioned diseases.

The compounds of the present invention may be administered alone or in combination with pharmaceutically acceptable carriers or diluents by either of the above routes previously indicated, and such administration can be carried out in single
10 or multiple doses. More particularly, the novel therapeutic agents of the invention can be administered in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, lozenges, trochees, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs,
15 syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various nontoxic organic solvents, etc. Moreover, oral pharmaceutical compositions can be suitably sweetened and/or flavored. In general, the therapeutically-effective compounds of this invention are present in such dosage forms at concentration levels ranging 5% to 70% by weight, preferably 10% to 50% by
20 weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dipotassium phosphate and glycine may be employed along with various disintegrants such as starch and preferably corn, potato or tapioca starch, alginic acid and certain complex silicates,
25 together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatine capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight
30 polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or

flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various combinations thereof.

For parenteral administration, solutions of a compound of the present invention
5 in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably buffered (preferably pH>8) if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intra-articular, intra-muscular and subcutaneous injection purposes. The preparation of all these
10 solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well-known to those skilled in the art. Additionally, it is also possible to administer the compounds of the present invention topically when treating inflammatory conditions of the skin and this may preferably be done by way of creams, jellies, gels, pastes, ointments and the like, in accordance with standard
15 pharmaceutical practice.

The compounds of formula (I) may also be administered in the form of suppositories for rectal or vaginal administration of the active ingredient. These compositions can be prepared by mixing the active ingredient with a suitable non-irritating excipient which is solid at room temperature (for example, 10 °C to 32 °C)
20 but liquid at the rectal temperature and will melt in the rectum or vagina to release the active ingredient. Such materials are polyethylene glycols, cocoa butter, suppository and wax.

For buccal administration, the composition may take the form of tablets or lozenges formulated in conventional manner.

25 *Combination with Other Drugs:*

Compounds of Formula I would be useful for, but not limited to, the treatment of inflammation in a subject, and for treatment of other inflammation-associated disorders, such as, as an analgesic in the treatment of pain and headaches, or as an antipyretic for the treatment of fever. For example, combinations of the invention would be useful
30 to treat arthritis, including but not limited to rheumatoid arthritis, spondyloarthropathies, gouty arthritis, osteoarthritis, systemic lupus erythematosus and juvenile arthritis.

Such combinations of the invention would be useful in the treatment of asthma, bronchitis, inmenstrual cramps, tendinitis, bursitis, and skin related conditions such as psoriasis, eczema, burns and dermatitis. Combinations of the invention also would be useful to treat gastrointestinal conditions such as inflammatory bowel disease.

5 Crohn's disease, gastritis, irritable bowel syndrome and ulcerative colitis and for the prevention of colorectal cancer. Combinations of the invention would be useful in creating inflammation in such diseases as vascular diseases, migraine headaches, periarteritis nodosa, thyroiditis, aplastic anemia, Hodgkin's disease, sclerodoma, rheumatic fever, type I diabetes, myasthenia gravis, multiple sclercsis, sarcoidosis,

10 nephrotic syndrome, Behcet's syndrome, polymyositis, gingivitis, hypersensitivity, Conjunctivitis, swelling occurring after injury, myocardial ischemia, and the like. The combinations would also be useful for the treatment of certain central nervous system disorders such as Alzheimer's disease and dimentia. The combinations of the invention are useful as anti-inflammatory agents, such as for the treatment of arthritis,

15 with the additional benefit of having significantly less harmful side effects. These compositions would also be useful in the treatment of allergic rhinitis, respiratory distress syndrome, endotoxin shock syndrome, atherosclerosis and central nervous system damage resulting from stroke, ischemia and trauma.

Compounds of formula (I) will be useful as a partial or complete substitute for

20 conventional NSAID's in preparations wherein they are presently co-administered with other agents or ingredients. Thus, the invention encompasses pharmaceutical compositions for treating COX-2 mediated diseases as defined above comprising a non-toxic therapeutically effective amount of the compound of formula (I) and one or more ingredients such as another pain reliever including acetaminophen or phenacetin;

25 a potentiator including caffeine; an H₂-antagonist, aluminom or magnesium hydroxide, simethicone, a decongestant including phenylephrine, phenylproanolamine, psuedophedrine, oxymetazoline, ephinephrine, naphazoline, xylometazoline, propylhexedrine, or levodesoxyephedrine; an antiitussive including codeine, hydrocodone, caramiphen, carbetapentane, or dexamethorphan; a prostaglandin

30 including misoprostol, enprostil, rioprostil, ornoprotol or rosaprostol; a diuretic; a sedating or non-sedating antihistamine; anticancer agents such as angiostatin and

endostatin; anti-Alzheimers such as Donepezil and Tacrine hydrochloride; and TNF alpha inhibitors such as Etanercept.

These cyclooxygenase inhibitors can further be used in combination with a nitric oxide inhibitors disclosed in WO 96/28145.

- 5 Also, the invention encompasses pharmaceutical compositions for treating COX-2 mediated diseases as defined above comprising a non-toxic therapeutically effective amount of the compound of formula (I) and one or more anti-ulcer agent and/or prostaglandins, which are disclosed in WO 97/11701.

10 The useful prostaglandins include misoprostol, plus-minus methyl 11 α , 16-dihydroxy-16-methyl-9-oxoprost 13E-en-1-oate; enisoprost and methyl-7-[2B-[6-(1-cyclopenten-1-yl)-4-hydroxy-4-methyl-1E, 5E-hexadienyl]-3 α -hydroxy-5-oxo 1R, 1 α -cyclopentyl]-4Z-heptenoate. Prostaglandins within the scope of the invention also include arbutoprost, enprostil, rioprostol, nocolprost, mexiprostil, ornoprostol, dimoxaprost, tioprostanide and rosaprostol.

- 15 The present compounds may also be used in co-therapies, partially or completely, in place of other conventional antiinflammatories, such as together with steroids, 5-lipoxygenase inhibitors, LTB₄ antagonists and LTA₄ hydrolase inhibitor's.

An example of LTB₄ is disclosed in WO97/29774. Suitable LTB₄ inhibitors include, among others, ebselen, Bayer Bay-x-1005, Ciba Geigy compound CGS-20 25019C, Leo Denmark compound ETH-615, Lilly compound LY-293111, Ono compound ONO-4057, Terumo compound TMK-688, Lilly compounds LY-213024, 264086 and 292728, Ono compound ONO-LB457, Searle compound SC-S3228, calcitrol, Lilly compounds LY-210073, LY223982, LY233469, and LY255283, Ono compound ONO-LB-448, Searle compounds SC-41930, SC-50605 and SC-51146, and 25 SK&F compound SKF-104493. Preferably, the LTB₄ inhibitors are selected from ebselen, Bayer Bay-x-1005, Ciba Geigy compound CGS-25019C, Leo Denmark compound ETH-61S, Lilly compound LY-293111, Ono compound ONO-4057 and Terumo compound TMK-688.

- 30 An example of 5-LO inhibitors is disclosed in WO97/29776. Suitable 5-LO inhibitors include, among others, masoprocol, tenidap, zileuton, pranlukast, tepoxalin, rilopirox, flezelastine hydrochloride, enazadrem phosphate and bunaprolast.

An example of LTA₄ hydrolase inhibitors is disclosed in WO97/29774. Suitable LTA₄ hydrolase inhibitors include, among others, Rhone-Poulenc Rorer RP-64966.

The administration of the present invention may be for either prevention or treatment purposes. The methods and compositions used herein may be used alone or in conjunction with additional therapies known to those skilled in the art in the prevention or treatment of angiogenesis. Alternatively, the methods and compositions described herein may be used as adjunct therapy. By way of example, the cyclooxygenase-2 inhibitor may be administered alone or in conjunction with other antineoplastic agents or other growth inhibiting agents or other drugs or nutrients.

There are large numbers of antineoplastic agents available in commercial use, in clinical evaluation and in pre-clinical development, which could be selected for treatment of angiogenesis by combination drug chemotherapy. Such antineoplastic agents fall into several major categories, namely, antibiotic-type agents, alkylating agents, antimetabolite agents, hormonal agents, immunological agents, interferon-type agents and a category of miscellaneous agents. Alternatively, other anti-neoplastic agents, such as metallomatrix proteases inhibitors (MMP), such as MMP-13 inhibitors including batistat, marimastat. Agouron Pharmaceuticals AG-3340, and Roche R0-32-3555, or alpha,beta,inhibitors may be used.

A first family of antineoplastic agents which may be used in combination with a selective cyclooxygenase-2 inhibitor consists of antimetabolite-type antineoplastic agents. Suitable antimetabolite antineoplastic agents may be selected from the group consisting of 5-FU-fibrinogen, acanthifolic acid, aminothiadiazole, brequinar sodium, carmofur, Ciba-Geigy CGP-30694, cyclopentyl cytosine, cytarabine phosphate stearate, cytarabine conjugates, Lilly DATHF, Merrel Dow DDFC, dezaguanine, dideoxycytidine, dideoxyguanosine, didox, Yoshitomi DMDC, doxifluridine, Wellcome EHNA, Merck & Co. EX-015, fazarabine, floxuridine, fludarabine phosphate, 5-fluorouracil, N-(2'-furanidyl)-5-fluorouracil, Daiichi Seiyaku F0-152, isopropyl pyrrolizine, Lilly LY-188011, Lilly LY-264618, methobenzaprim, methotrexate, Wellcome MZPES, norspermidine, NCI NSC-127716, NCI NSC-264880, NCI NSC-39661, NCI NSC-612567, Warner-Lambert PALA, pentostatin,

piritrexim, plicamycin, Asahi Chemical PL-AC, Takeda TAC-788, thioguanine, tiazofurin, Erbamont TIF, trimetrexate, tyrosine kinase inhibitors, tyrosine protein kinase inhibitors, Taiho UFT and uricytin.

A second family of antineoplastic agents which may be used in combination with a selective cyclooxygenase-2 inhibitor consists of alkylating-type antineoplastic agents. Suitable alkylating-type antineoplastic agents may be selected from the group consisting of Shionogi 254-S, aldo-phosphamide analogues, altretamine, anaxirone, Boehringer Mannheim BBR-2207, bestabucil, budotitane, Wakunaga CA-102, carboplatin, carmustine, Chinoin-139, Chinoin-153, chlorambucil, cisplatin, cyclophosphamide, American Cyanamid CL-286558, Sanofi CY-233, cyplatate, Degussa D-19-384, Sumimoto DACHP(My₂), diphenylspiromustine, diplatinum cytostatic. Erba distamycin derivatives, Chugai DWA-2114R, ITI E09, elmustine, Erbamont FCE-24517, estramustine phosphate sodium, fotemustine, Unimed G-6-M, Chinoin GYKI-17230, hepsul-fam, ifosfamide, iproplatin, lomustine, mafosfamide, mitolactol, Nippon Kayaku NK-121, NCI NSC-264395, NCI NSC-342215, oxaliplatin, Upjohn PCNU, prednimustine, Proter PTT-II9, ranimustine, semustine, SmithKline SK&F-I01772, Yakult Honsha SN-22, spiromus-tine, Tanabe Seiyaku TA-077, tauromustine, temozolomide, teroxirone, tetraplatin and trimelamol.

A third family of antineoplastic agents which may be used in combination with a selective cyclooxygenase-2 inhibitor consists of antibiotic-type antineoplastic agents. Suitable antibiotic-type antineoplastic agents may be selected from the group consisting of Taiho 4181-A, aclarubicin, actinomycin D, actinoplanone, Erbamont ADR-456, aeroplysinin derivative, Ajinomoto AN-201-II. Ajinomoto AN-3, Nippon Soda anisomycins, anthracycline, azino-mycin-A, bisucaberin, Bristol-Myers BL-6859, Bristol-Myers BMY-25067, Bristol-Myers BMY-25551, Bristol-Myers BMY-26605, Bristol-Myers BMY-27557, Bristol-Myers BMY-28438, bleomycin sulfate, bryostatins, Taiho C-1027, caliche mycin, chromoximycin, dactinomycin, daunorubicin, Kyowa Hakko DC-102, Kyowa Hakko DC-79, Kyowa Hakko DC-88A, Kyowa Hakko DC89-A1, Kyowa Hakko DC92-B, ditrisarubicin B, Shionogi DOB-41, doxorubicin, doxorubicin-fibrinogen, elsamicin-A, epirubicin, erbstatin, esorubicin, esperamicin-A1, esperamicin-Alb. Erbamont FCE-21954, Fujisawa FK-973, fostriecin, Fujisawa FR-

900482, glidobactin, gregatin-A, grincamycin, herbimycin, idarubicin, illudins, kazusamycin, kesarirhodins, Kyowa Hakko KM-5539, Kirin Brewery KRN-8602, Kyowa Hakko KT-5432, Kyowa Hakko KT-5594, Kyowa Hakko KT-6149, American Cyanamid LL-D49194, Meiji Seika ME 2303, menogaril, mitomycin, mitoxantrone, 5 SmithKline M-TAG, neoenactin, Nippon Kayaku NK-313, Nippon Kayaku NKT-OI, SRI International NSC-357704, oxalysine, oxaunomycin, peplomycin, pilatin, pirarubicin, porothramycin, pyrindamycin A, Tobishi RA-I, rapamycin, rhizoxin, rodorubicin, sibanomicin, siwenmycin, Sumitomo SM-5887, Snow Brand SN-706, Snow Brand SN-07, sorangicin-A, sparsomycin, SS Pharmaceutical SS-21020, SS 10 Pharmaceutical SS-7313B, SS Pharmaceutical SS-9816B, steffimycin B, Taiho 4181-2, talisomycin, Takeda TAN-868A, terpentecin, thiazine, tricrozarin A, Upjohn U-73975, Kyowa Hakko UCN-I0028A, Fujisawa WF-3405, Yoshitomi Y-2S024 and zorubicin.

A fourth family of antineoplastic agents which may be used in combination with the selective cyclooxygenase-2 inhibitor consists of a miscellaneous family of 15 antineoplastic agents selected from the group consisting of alpha-carotene, alpha-difluoromethyl-arginine, acitretin, Biotec AD-5, Kyorin AHC-52, alstonine, amonafide, amphetinile, amsacrine, Angiostat, ankinomycin, anti-neoplaston AIO, antineoplaston A2, antineoplaston A3, antineoplaston A5, antineoplaston AS2-1, Henkel APD, aphidicolin glycinate, asparaginase, Avarol, baccharin, batracylin, benfluron, benzotript, 20 Ipsen-Beaufour BIM-23015, bisantrene, Bristo-Myers BMY-40481, Vestar boron-IO, bromofosfamide, Wellcome BW-502, Wellcome BW-773, caracemide, carmethizole hydrochloride, Ajinomoto CDAF, chlorsulfaquinoxalone, Chemes CHX-2053, Chemex CHX-100, Warner-Lambert CI-921, Warner-Lambert CI-937, Warner-Lambert CI-941, Warner-Lambert CI-958, clanfenur, claviridenone, ICN compound 25 1259, ICN compound 4711, Contracan, Yakult Honsha CPT-11, crisnatol, curaderm, cytochalasin B, cytarabine, cytosytin, Merz D-609, DABIS maleate, dacarbazine, datelliptinium, didemnin-B, dihaematoporphyrin ether, dihydrolenperone, dinaline, distamycin, Toyo Pharmar DM-341, Toyo Pharmar DM-75, Daiichi Seiyaku DN-9693, elliprabin, elliptinium acetate, Tsumura EPMTTC, ergotamine, etoposide, etretinate, 30 fenretinide, Fujisawa FR-57704, gallium nitrate, genkwadaphnin, Chugai GLA-43, Glaxo GR-63178, grifolan NMF-5N, hexadecylphosphocholine, Green Cross H0-221,

homoharringtonine, hydroxyurea, BTG ICRF-187, ilmofofine, isoglutamine, isotretinoin. Otsuka JI-36, Ramot K-477, Otsuak K-76COONa, Kureha Chemical K-AM, MECT Corp KI-8110, American Cyanamid L-623, leukoregulin, Ionidamine, Lundbeck LU-23-112, Lilly LY-186641, NCI (US) MAP, marycin, Merrel Dow MDL-5 27048, Medco MEDR-340, merbarone, merocyanine derivatives, methylanilinoacridine, Molecular Genetics MGI-136, minactivin, mitonafide, mitoquidone, mopidamol, motretinide, Zenyaku Kogyo MST-16, N-(retinoyl)amino acids, Nisshin Flour Milling N-021, N-acylated-dehydroalanines, nafazatrom, Taisho NCU-190, nocodazole derivative, Normosang, NCI NSC-145813, NCI NSC-361456, NCI NSC-604782, NCI 10 NSC-95580, octreotide, Ono ON0-112, oquizanocine, Akzo Org-10172, pancratistatin, pazelliptine, Warner-Lambert PD-111707, Warner-Lambert PD-115934, Warner-Lambert PD-131141, Pierre Fabre PE-1001, ICRT peptide D, piroxantrone, polyhaematoporphyrin, polypreic acid, Efamol porphyrin, probimane, procarbazine, proglurnide, Invitron protease nexin I, Tobishi RA-700, razoxane, Sapporo Breweries 15 RBS, restrictin-P, retelliptine, retinoic acid, Rhone-Poulenc RP-49532, Rhone-Poulenc RP-56976, SmithKline SK&F-104864, Sumitomo SM-108, Kuraray SMANCS, SeaPharm SP-10094, spatol, spirocyclopropane derivatives, spirogermanium, Unimed, SS Pharmaceutical SS-554, strypoldinone, Stypoldione, Suntory SUN 0237, Suntory SUN 2071, superoxide dismutase, Toyama T-506, Toyama T-680, taxol, Teijin TEI-20 0303, teniposide, thaliblastine, Eastman Kodak TJB-29, tocotrienol, Topostin, Teijin TT-82, kyowa Hakko UCN-01, Kyowa Hakko UCN-1028, ukrain, Eastman Kodak USB-006, vinblastine sulfate, vincristine, vindesine, vinestramide, vinorelbine, vintriptol, vinzolidine, withanolides and Yamanouchi YM-534.

Examples of radioprotective agents which may be used in the combination 25 chemotherapy of this invention are AD-5, adchnon, amifostine analogues, detox, dimesna, I-102, MN-159, N-acylated-dehydroalanines, TGF-Genentech, tiprotimod, amifostine, WR-151327, FUT-187, ketoprofen transdermal, naburnetone, superoxide dismutase (Chiron) and superoxide dismutase Enzon.

Methods for preparation of the antineoplastic agents described above may be 30 found in the literature. Methods for preparation of doxorubicin, for example, are described in U.S. Patents No. 3,590,028 and No. 4,012,448. Methods for preparing

metallomatrix protease inhibitors are described in EP 780386, W097/20824, W096/15096. Methods for preparing SOD mimics are described in EP 524,101. Methods for preparing alpha,beta, inhibitors are described in W097/08174.

In addition, the selective COX-2 inhibitor may be administered in conjunction
5 with other antiinflammatory agents for maximum safety and efficacy, including NSAID's, selective COX-1 inhibitors and inhibitors of the leukotriene pathway, including 5-lipoxygenase inhibitors. Examples of NSAID's include indomethacin, naproxen, ibuprofen, salicylic acid derivatives such as aspirin, diclofenac, ketorolac, piroxicam, meloxicam, mefenamic acid, sulindac, tolmetin sodium, zomepirac,
10 fenoprofen, phenylbutazone, oxyphenbutazone, nimesulide, zaltoprofen and letodolac.

Method for assessing biological activities:

The activity of the compounds of the formula (I) of the present invention was demonstrated by the following assays.

In vitro assays

15 *Human cell based COX-1 assay*

Human peripheral blood obtained from healthy volunteers was diluted to 1/10 volume with 3.8% sodium citrate solution. The platelet-rich plasma immediately obtained was washed with 0.14 M sodium chloride containing 12 mM Tris-HCl (pH 7.4) and 1.2 mM EDTA. Platelets were then washed with platelet buffer (Hanks
20 buffer (Ca free) containing 0.2% BSA and 20 mM Hepes). Finally, the human washed platelets (HWP) were suspended in platelet buffer at the concentration of 2.85×10^8 cells/ml and stored at room temperature until use. The HWP suspension (70 μ l aliquots, final 2.0×10^7 cells/ml) was placed in a 96-well U bottom plate and 10 μ l aliquots of 12.6 mM CaCl_2 added. Platelets were incubated with A23187 (final 10
25 μ M, Sigma) with test compound (0.1 - 100 μ M) dissolved in DMSO (final concentration; less than 0.01%) at 37 °C for 15 min. The reaction was stopped by addition of EDTA (final 7.7 mM) and TxB_2 in the supernatant quantitated by using a radioimmunoassay kit (Amersham) according to the manufacturer's procedure.

Human cell based COX-2 assay

30 Inhibition of COX-2 activity after induction of COX-2 by hIL-1 β

The human cell based COX-2 assay was carried out as previously described

(Moore *et al.*, *Inflam. Res.*, **45**, 54, 1996). Confluent human umbilical vein endothelial cells (HUVECs, Morinaga) in a 96-well U bottom plate were washed with 100 μ l of RPMI1640 containing 2% FCS and incubated with hIL-1 β (final concentration 300 U/ml, R & D Systems) at 37 °C for 24 hr. After washing, the
5 activated HUVECs were stimulated with A23187 (final concentration 30 μ M) in Hanks buffer containing 0.2% BSA, 20 mM Hepes and test compound (0.1 nM - 100 μ M) dissolved in DMSO (final concentration; less than 0.01%) at 37 °C for 15 min. 6-Keto-PGF $_{1\alpha}$, stable metabolite of PGI $_2$, in the supernatant was quantitated after adequate dilution by using a radioimmunoassay kit (Amersham) according to the
10 manufacturer's procedure.

Inhibition of COX-2 during the induction phase

Confluent human umbilical vein endothelial cells (HUVECs, Morinaga) in a 96-well U bottom plate were washed with 100 μ l of RPMI1640 containing 2% FCS and test compound (0.1 nM - 100 μ M) dissolved in DMSO (final concentration; less
15 than 0.01%), and incubated with hIL-1 β (final concentration 300 U/ml, R & D Systems) at 37 °C for 24 hr. After washing, the HUVECs were stimulated with A23187 (final concentration 30 μ M) in Hanks buffer containing 0.2% BSA and 20 mM Hepes at 37 °C for 15 min. 6-Keto-PGF $_{1\alpha}$, a stable metabolite of PGI $_2$, in the supernatant was quantitated after adequate dilution by using a radioimmunoassay kit
20 (Amersham) according to the manufacturer's procedure.

In vivo assays

Carrageenan induced foot edema in rats

Male Sprague-Dawley rats (5 weeks old, Charles River Japan) were fasted overnight. A line was drawn using a marker above the ankle on the right hind paw
25 and the paw volume (V0) was measured by water displacement using a plethysmometer (Muromachi). Animals were given orally either vehicle (0.1% methyl cellulose or 5% Tween 80) or a test compound (2.5 ml per 100 g body weight). One hour later, the animals were then injected intradermally with λ -carrageenan (0.1 ml of 1% w/v suspension in saline, Zushikagaku) into right hind paw (Winter *et al.*, *Proc.*
30 *Soc. Exp. Biol. Med.*, **111**, 544, 1962; Lombardino *et al.*, *Arzneim. Forsch.*, **25**, 1629, 1975) and three hours later, the paw volume (V3) was measured and the increase in

volume (V3-V0) calculated. Since maximum inhibition attainable with classical NSAIDs is 60-70%, ED30 values were calculated.

Gastric ulceration in rats

The gastric ulcerogenicity of test compound was assessed by a modification of the conventional method (Ezer *et al.*, *J. Pharm. Pharmacol.*, **28**, 655, 1976; Cashin *et al.*, *J. Pharm. Pharmacol.*, **29**, 330 - 336, 1977). Male Sprague-Dawley rats (5 weeks old, Charles River Japan), fasted overnight, were given orally either vehicle (0.1% methyl cellulose or 5% Tween 80) or a test compound (1 ml per 100 g body weight). Six hours after, the animals were sacrificed by cervical dislocation. The stomachs were removed and inflated with 1% formalin solution (10 ml). Stomachs were opened by cutting along the greater curvature. From the number of rats that showed at least one gastric ulcer or haemorrhaging erosion (including ecchymosis), the incidence of ulceration was calculated. Animals did not have access to either food or water during the experiment.

Data Analysis

Statistical program packages, SYSTAT (SYSTAT, INC.) and StatView (Abacus Concepts, Inc.) for Macintosh were used. Differences between test compound treated group and control group were tested for using ANOVA. The IC₅₀ (ED₃₀) values were calculated from the equation for the log-linear regression line of concentration (dose) versus percent inhibition.

Some compounds prepared in the Working Examples as described herein after were tested by these methods, and showed IC₅₀ values of 0.001 μ M to 10 μ M with respect to inhibition of COX-2.

Also, the above-mentioned most preferred compounds were tested by these methods, and showed IC₅₀ values of 0.001 μ M to 0.5 μ M with respect to inhibition of COX-2.

COX-2 selectivity can be determined by ratio in terms of IC₅₀ value of COX-1 inhibition to COX-2 inhibition. In general, it can be said that a compound showing a COX-1/COX-2 inhibition ratio of more than 2 has good COX-2 selectivity.

Some compounds prepared in Examples showed COX-1/COX-2 inhibition ratio of more than 10.

The following examples contain detailed descriptions of the methods of the preparation of compounds of formula (I). These detailed descriptions fall within the scope of the invention and serve to exemplify the above described general synthetic procedures which form part of the invention. These detailed descriptions are presented for illustrative purposes only and are not intended to restrict the scope of the present invention.

EXAMPLES

The invention is illustrated in the following non-limiting examples in which, unless stated otherwise: all operations were carried out at room or ambient temperature, that is, in the range of 18-25 °C; evaporation of solvent was carried out using a rotary evaporator under reduced pressure with a bath of up to 60 °C; reactions were monitored by thin layer chromatography (tlc) and reaction times are given for illustration only; melting points (m.p.) given are uncorrected (polymorphism may result in different melting points); structure and purity of all isolated compounds were assured by at least one of the following techniques: tlc (Merck silica gel 60 F-254 precoated plates), mass spectrometry, nuclear magnetic resonance (NMR) or microanalysis. Yields are given for illustrative purposes only. Flash column chromatography was carried out using Merck silica gel 60 (230-400 mesh ASTM). Low-resolution mass spectral data (EI) were obtained on a Automass 120 (JEOL) mass spectrometer. Low-resolution mass spectral data (ESI) were obtained on a Quattro II (Micromass) mass spectrometer. NMR data was determined at 270 MHz (JEOL JNM-LA 270 spectrometer) using deuterated chloroform (99.8% D) or dimethylsulfoxide (99.9% D) as solvent unless indicated otherwise, relative to tetramethylsilane (TMS) as internal standard in parts per million (ppm); conventional abbreviations used are: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, br = broad, etc.

EXAMPLE 1

ETHYL (2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)ACETATE

STEP 1. Ethyl *trans*-4-chloro-2-nitrocinnamate

To a suspension of sodium hydride (60% w/w dispersion in mineral oil, 4.4 g, 0.11 mol) in THF (150 ml) was added dropwise a solution of triethyl phosphonoacetate (25.0 g, 0.11 mol) in THF (50 ml) at room temperature. After stirring for 1 h, a

solution of 4-chloro-2-nitrobenzaldehyde (19.0 g, 0.10 mol) in THF (50 ml) was added. After stirring for an additional 1 h, saturated aqueous ammonium chloride (50 ml) was added and the resulting mixture was extracted with ethyl acetate (300 ml x 2). The combined organic extracts were dried (MgSO₄) and concentrated to gave 27 g (quant.) of the title compound as brown solids.

¹H-NMR (CDCl₃) δ: 8.04 (1H, d, J=15.8 Hz), 8.03 (1H, d, J=1.8 Hz), 7.64-7.58 (2H, m), 6.36 (1H, d, J=15.8 Hz), 4.30 (2H, q, J=7.0 Hz), 1.35 (3H, t, J=7.0 Hz).

STEP 2. Ethyl *trans*-2-amino-4-chlorocinnamate

A mixture of ethyl *trans*-4-chloro-2-nitrocinnamate (step 1, 27.0 g, 0.11 mol) and sodium hydrosulfite (92 g, 0.53 mol) in THF-H₂O (1:1, 500 ml) was stirred at room temperature for 1h. Saturated aqueous sodium bicarbonate (300 ml) was then added and the mixture was extracted with ethyl acetate (300 ml x 2). The combined organic extracts were dried (MgSO₄) and concentrated to gave 16.7 g (67%) of the title compound as yellow solids.

¹H-NMR (CDCl₃) δ: 7.72 (1H, d, J=15.8 Hz), 7.27 (1H, d, J=8.4 Hz), 6.78-6.68 (2H, m), 6.31 (1H, d, J=15.8 Hz), 4.27 (2H, q, J=7.0 Hz), 1.33 (3H, t, J=7.0 Hz).

STEP 3. Ethyl *trans*-4-chloro-2-formamidocinnamate

A mixture of acetic anhydride (20 ml) and formic acid (10 ml) was heated at 60 °C for 2 h. After cooling to 0 °C, a solution of ethyl *trans*-2-amino-4-chlorocinnamate (step 2, 15.5 g, 0.069 mol) in THF (80 ml) was carefully added. The resulting mixture was allowed to warm to room temperature. After stirring overnight, the mixture was concentrated and the precipitates were collected by filtration. The solids were washed with hexane to give 9.6 g (55%) of the title compound.

¹H-NMR (CDCl₃) δ: 9.40-9.15 (1H, m), 8.51-8.40 (1H, m), 8.10-7.80 (2H, m), 7.60-7.47 (1H, m), 7.28-7.12 (1H, m), 6.40 (1H, d, J=15.8 Hz), 4.25 (2H, q, J=7.3 Hz), 1.34 (3H, t, J=7.3 Hz).

STEP 4. Ethyl *trans*-4-chloro-2-isocyanocinnamate

To a solution of triphenylphosphine (5.3 g, 20 mmol) in dichloromethane (80 ml) cooled to 0 °C was added dropwise a solution of triphosgene (2.0 g, 6.7 mmol) in dichloromethane (20 ml). The ice-bath was removed and the resulting mixture stirred

at room temperature for 10 min. The mixture was then cooled to 0 °C and a solution of ethyl *trans*-4-chloro-2-formamidocinnamate (step 3, 5.2 g, 0.020 mol) in dichloromethane (80 ml) was added. The mixture was allowed to warm to room temperature overnight, and then concentrated. The residue was partitioned between
5 water (80 ml) and ethyl acetate (100 ml), the aqueous layer separated and extracted with ethyl acetate (100 ml). The combined organic extracts were dried (MgSO₄), solvent removed by evaporation and the crude product was purified by flash column chromatography eluting with ethyl acetate/hexane (1:6) to afford 3.9 g (83 %) of the title compound as white solids.

10 ¹H-NMR (CDCl₃) δ: 7.89 (1H, d, J=16.1 Hz), 7.60 (1H, d, J=8.8 Hz), 7.45 (1H, d, J=1.8 Hz), 7.42 (1H, dd, J=1.8, 8.8 Hz), 6.52 (1H, d, J=16.1 Hz), 4.30 (2H, q, J=7.0 Hz), 1.35 (3H, t, J=7.0 Hz).

STEP 5. Ethyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate

A mixture of ethyl *trans*-4-chloro-2-isocyanocinnamate (step 4, 1.2 g, 5.1
15 mmol), tributyltin hydride (1.6 g, 5.6 mmol) and AIBN (43 mg, 0.26 mmol) in acetonitrile (30 ml) was heated at 100 °C. After 1h, tetrakis(triphenylphosphine)palladium (580 mg, 0.50 mmol) and benzoyl chloride (0.65 ml, 5.6 mmol) were added and the mixture was heated for a further 17 h. The mixture was cooled and poured into 2N aqueous HCl (50 ml) and extracted with diethyl ether
20 (80 ml x 2). The combined organic extracts were washed with saturated aqueous potassium fluoride (50 ml) and dried (MgSO₄). After removal of solvent, the crude product was purified by flash column chromatography eluting with ethyl acetate/hexane (1:5) to afford 0.43 g (25 %) of the title compound as white solids.

m.p.: 160-163 °C.

25 ¹H-NMR (CDCl₃) δ: 8.94 (1H, br s), 7.82-7.75 (2H, m), 7.67-7.47 (4H, m), 7.37 (1H, d, J=1.8 Hz), 7.13 (1H, dd, J=1.8, 8.4 Hz), 4.11 (2H, q, J=7.3 Hz), 3.78 (2H, s), 1.22 (3H, t, J=7.3 Hz).

EXAMPLE 2

(2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)ACETIC ACID

30 METHOD A

To a solution of ethyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate (Example 1,

380 mg, 0.11 mmol) in ethanol (15 ml) was added 2N aqueous KOH (5 ml). After heating at 80 °C for 1 h, the mixture was cooled and concentrated, and then 2N aqueous HCl (15 ml) added carefully. The mixture was extracted with diethyl ether (50 ml x 2), the combined organic extracts dried (MgSO₄) and concentrated. The residual solids were recrystallized from ethyl acetate/hexane to afford 60 mg (17%) of the title compound as pale yellow solids.

m.p.: 183-186 °C.

IR (KBr) ν : 1700, 1610, 1520, 1425, 1330, 1000 cm⁻¹.

¹H-NMR (DMSO-d₆) δ : 12.26 (1H, br s), 11.76 (1H, s), 7.77-7.66 (4H, m), 7.62-7.54 (2H, m), 7.48 (1H, d, J=1.8 Hz), 7.13 (1H, dd, J=1.8, 8.7 Hz), 3.80 (2H, s).

METHOD B

STEP 1. 6-Chloro-1-(phenylsulfonyl)indole

A mixture of 6-chloroindole (Y. Watanabe et al., *J. Org. Chem.*, **1990**, 55, 580, 36.2 g, 0.24 mol), tetrabutylammonium hydrogen sulfate (8.1 g, 0.024 mol) and 50 % aqueous KOH (160 ml) in benzene (500 ml) was stirred at room temperature for 10 min. The mixture was then cooled to 0 °C and a solution of benzenesulfonyl chloride in benzene (20 ml) was added. After stirring at room temperature for 3 h, the mixture was poured into water (200 ml), the organic layer separated and the aqueous layer extracted with diethyl ether (200 ml x 2). The combined organic extracts were washed with brine (200 ml), dried (MgSO₄) and concentrated. The residual solids were washed with ethanol (100 ml x 3) to give 58 g (83%) of the title compound as off-white solids.

¹H-NMR (CDCl₃) δ : 8.02 (1H, s), 7.92-7.85 (2H, m), 7.60-7.40 (5H, m), 7.21 (1H, dd, J=1.8, 8.4 Hz), 6.62 (1H, d, J=3.6 Hz).

STEP 2. 6-chloro-2-benzoyl-1-(phenylsulfonyl)indole

To a stirred solution of 6-chloro-1-(phenylsulfonyl)indole (Step 1, 12.58 g, 43.0 mmol) in THF (270 ml) cooled to - 78 °C was added dropwise *tert*-butyllithium (32 ml, 52.0 mmol, 1.64 M in *n*-pentane) with keeping the internal temperature below - 65 °C. After stirring for 30 min. at -78 °C, this solution was transferred via cannula to a solution of benzoyl chloride (6.0 ml, 52.0 mmol) in THF (30 ml) cooled to - 78 °C. The mixture was stirred for 1.5 h and then quenched with saturated ammonium

chloride (200 ml) at -78 °C and allowed to warm to room temperature. The aqueous layer was separated and neutralized with aqueous sodium carbonate, and then extracted with ethyl acetate (50 ml x 2). The combined organic extracts were washed with brine (50 ml), dried (MgSO₄), and concentrated. Crystallization of the residue from diethyl ether/hexane (1:3) afforded the title compound as white solids (14.4 g, 85%).

¹H-NMR(CDCl₃)δ: 8.20 - 8.16 (1H, m), 8.14 - 8.06 (2H, m), 8.01 - 7.93 (2H, m), 7.66 - 7.47 (7H, m), 7.29 (1H, dd, J=1.7, 8.5Hz), 6.89 (1H, J=0.7Hz).

STEP 3. 2-Benzoyl-6-chloroindole

A mixture of 2-benzoyl-6-chloro-1-(phenylsulfonyl)indole (step 2, 48 g, 0.12 mol) and potassium carbonate (80 g, 0.58 mol) in THF-MeOH-H₂O (4:2:1, 1100 ml) was heated at reflux temperature overnight. After removal of solvent, the residue was extracted with diethyl ether (300 ml x 2) and dried (MgSO₄). Removal of solvent gave the crude product as pale brown solids. Recrystallization from ethyl acetate afforded 20 g (65%) of the title compound as white solids.

m.p.: 206-207 °C.

¹H-NMR (CDCl₃) δ: 9.31 (1H, br s), 8.01-7.95 (2H m), 7.68-7.47 (5H, m), 7.17-7.12 (2H, m).

STEP 4. Diethyl α-acetoxy-(2-benzoyl-6-chloro-1H-indol-3-yl)malonate

A mixture of 2-benzoyl-6-chloroindole (step 3, 4.0 g, 16 mmol), manganese(III) acetate dihydrate (13 g, 48 mmol), diethyl malonate (14 g, 80 mmol) and sodium acetate (6.6 g, 80 mmol) in acetic acid (150 ml) was heated at 80 °C with stirring for 2h. Manganese(III) acetate dihydrate (3 g, 11 mmol) was added and heating was continued for an additional 2 h. The mixture was cooled and brine (200 ml) was added. The resulting mixture was extracted with diethyl ether (200 ml x 2) and the combined organic extracts dried (MgSO₄) and concentrated. The residue was purified by flash column chromatography eluting with ethyl acetate/hexane (1:3) to afford 5.2 g (69 %) of the title compound as yellow solids.

m.p.: 141-144 °C.

¹H-NMR (CDCl₃) δ: 8.84 (1H, br s), 7.90-7.81 (3H, m), 7.66-7.58 (1H, m), 7.51-7.42 (2H, m), 7.38 (1H, d, J=1.3 Hz), 7.15 (1H, dd, J=2.0, 8.7 Hz), 4.30-4.06 (4H, m), 1.70 (3H, s), 1.30-1.58 (6H, m).

STEP 5. Diethyl (2-benzoyl-6-chloro-1H-indol-3-yl)malonate

A mixture of diethyl α -acetoxy-(2-benzoyl-6-chloro-1H-indol-3-yl)malonate (step 4, 5.0 g, 11 mmol), trifluoroacetic acid (3.3 ml, 44 mmol) and triethylsilane (2.1 ml, 13 mmol) in dichloromethane (80 ml) was heated at reflux temperature for 12 h and then cooled and concentrated. The resulting residue was partitioned between saturated sodium bicarbonate (50 ml) and dichloromethane (80 ml). The aqueous layer was separated and extracted with dichloromethane (80 ml). The combined organic extracts were dried (MgSO_4) and solvent removed. Crude product was purified by flush column chromatography eluting with ethyl acetate/hexane (1:4) to afford 4.0 g (87 %) of the title compound as white solids.

$^1\text{H-NMR}$ (CDCl_3) δ : 8.93 (1H, br s), 7.81-7.70 (3H, m), 7.68-7.60 (1H, m), 7.55-7.48 (2H, m), 7.32 (1H, s), 7.12 (1H, dd, $J=1.8, 8.8$ Hz), 5.29 (1H, s), 4.26-4.09 (4H, m), 1.21 (6H, t, $J=7.1$ Hz).

STEP 6. (2-Benzoyl-6-chloro-1H-indol-3-yl)acetic acid

Diethyl (2-benzoyl-6-chloro-1H-indol-3-yl)malonate (step 5, 4.4 g, 11 mmol) in a mixture of ethanol (120 ml) and 2N aqueous NaOH (15 ml) was heated at reflux temperature for 1 h. The mixture was cooled and concentrated, and the residue carefully acidified with 2N aqueous HCl (30 ml). The mixture was extracted with diethyl ether (150 ml x 3), and the combined extracts were dried (MgSO_4) and concentrated. The residual solids were recrystallized from ethyl acetate/hexane to afford 1.1 g (30%) of the title compound as pale yellow solids.

m.p.: 183-186 °C.

IR (KBr) ν : 1700, 1610, 1520, 1425, 1330, 1000 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.26 (1H, br s), 11.76 (1H, s), 7.77-7.66 (4H, m), 7.62-7.54 (2H, m), 7.48 (1H, d, $J=1.8$ Hz), 7.13 (1H, dd, $J=1.8, 8.7$ Hz), 3.80 (2H, s).

EXAMPLE 3**(2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)ACETIC ACID, SODIUM SALT**

(2-Benzoyl-6-chloro-1H-indol-3-yl)acetic acid (Example 2, 480 mg, 1.5 mmol) in ethanol (10 ml) was treated with 2N aqueous NaOH (0.7 ml, 1.4 mmol) at room temperature for 30 min. and then concentrated. The residue was dissolved in water

(10 ml) and washed with diethyl ether (15 ml x 2). The aqueous layer was concentrated to afford 350 mg (68%) of the title compound as pale brown solids.

m.p.: 185-189 °C.

IR (KBr) ν : 1523, 1380, 1230, 1060, 1004, 918 cm^{-1} .

- 5 $^1\text{H-NMR}$ (DMSO-d_6) δ : 11.46 (1H, br s), 7.88-7.84 (2H, m), 7.66 (1H, d, $J=8.4$ Hz), 7.64-7.46 (3H, m), 7.39 (1H, d, $J=1.8$ Hz), 7.00 (1H, dd, $J=1.8, 8.4$ Hz), 3.32 (2H, s).

EXAMPLE 4

[6-CHLORO-2-(2-METHYLBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

STEP 1. 6-Chloro-2-(2-methylbenzoyl)-1-(phenylsulfonyl)indole

- 10 The title compound was prepared according to the procedure described in step 2 of Example 2 (Method B) from 6-chloro-1-(phenylsulfonyl)indole (step 1 of Example 2, Method B) and *o*-toluoyl chloride.

tlc: $R_f=0.3$ (ethyl acetate/hexane=1:10).

STEP 2. 6-Chloro-2-(2-methylbenzoyl)indole

- 15 The title compound was prepared according to the procedure described in step 3 of Example 2 (Method B) from 6-chloro-2-(2-methylbenzoyl)-1-(phenylsulfonyl)indole (step 1).

$^1\text{H-NMR}$ (CDCl_3) δ : 9.37 (1H, br s), 7.58 (2H, d, $J=8.91\text{Hz}$), 7.48 (1H, s), 7.42 (1H, dd, $J=1.49, 7.75\text{Hz}$), 7.34-7.27 (2H, m), 7.12 (2H, dd, $J=1.81, 8.56\text{Hz}$), 2.44 (3H, s).

- 20 STEP 3. Diethyl α -acetoxy-[6-chloro-2-(2-methylbenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 6-chloro-2-(2-methylbenzoyl)indole (step 2).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.52 (1H, br s), 7.78 (1H, d, $J=8.88\text{Hz}$), 7.48-7.13 (6H, m), 4.31-4.16 (4H, m), 2.52 (3H, s), 1.96 (3H, s), 1.22 (6H, t, 7.26Hz).

- 25 STEP 4. Diethyl [6-chloro-2-(2-methylbenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetoxy-[6-chloro-2-(2-methylbenzoyl)-1H-indol-3-yl]malonate (step 3).

- 30 $^1\text{H-NMR}$ (CDCl_3) δ : 9.0 (1H, br s), 7.73 (1H, d, $J=8.88\text{Hz}$), 7.73-7.09 (5H, m), 7.11 (1H, dd, $J=1.97, 8.75\text{Hz}$), 4.89 (1H, s), 4.20-4.08 (4H, m), 2.35 (3H, s), 1.19 (6H, t,

J=7.1Hz).

STEP 5. [6-Chloro-2-(2-methylbenzoyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [6-chloro-2-(2-methylbenzoyl)-1H-indol-3-yl]malonate (step 4).

m.p.: 150-152 °C.

IR (KBr) ν : 3321, 1717, 1624, 1602, 1568, 1531, 1431, 1319, 1249, 1230 cm^{-1} .

$^1\text{H-NMR}$ (DMSO- d_6) δ : 11.70 (1H, s), 7.69 (2H, d, 8.72Hz), 7.51-7.31 (3H, m), 7.10 (2H, dd, J=1.97, 8.56Hz), 3.57 (2H, s), 2.24 (3H, s).

10 EXAMPLE 5

[6-CHLORO-2-(3-METHYLBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

STEP 1. 6-Chloro-2-(3-methylbenzoyl)-1-(phenylsulfonyl)indole

The title compound was prepared according to the procedure described in step 2 of Example 2 (Method B) from 6-chloro-1-(phenylsulfonyl)indole (step 1 of Example 2, Method B) and *m*-toluoyl chloride.

tlc: R_f=0.3 (ethyl acetate/hexane=1:10).

STEP 2. 6-Chloro-2-(3-methylbenzoyl)indole

The title compound was prepared according to the procedure described in step 3 of Example 2 (Method B) from 6-chloro-2-(3-methylbenzoyl)-1-(phenylsulfonyl)indole (step 1).

$^1\text{H-NMR}$ (CDCl_3) δ : 9.37 (1H, s), 7.77 (2H, br s), 7.64 (1H, d, J=8.56Hz), 7.47-7.12 (5H, m), 2.47 (3H, s).

STEP 3. Diethyl α -acetoxy-[6-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 6-chloro-2-(3-methylbenzoyl)indole (step 2).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.80 (1H, br s), 7.83 (1H, d, J=8.88Hz), 7.67-7.32 (5H, m), 7.16 (1H, dd, J=1.81, 8.88Hz), 4.27-4.15 (4H, m), 2.39 (3H, s), 1.72 (3H, s), 1.29 (6H, t, J=7.26Hz).

STEP 4. Diethyl [6-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]malonate

30 The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetoxy-[6-chloro-2-(3-methylbenzoyl)-1H-

indol-3-yl]malonate (step 3).

¹H-NMR (CDCl₃) δ: 9.21 (1H, br s), 7.71 (1H, d, J=8.51Hz), 7.59-7.28 (5H, m), 7.10 (1H, dd, J=1.97, 8.72Hz), 5.27 (1H, s), 4.23-4.07 (4H, m), 2.40 (3H, s), 1.21 (6H, t, J=7.1Hz).

5 STEP 5. [6-Chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [6-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]malonate (step 4).

m.p.: 182-184 °C.

10 IR (KBr) ν: 3313, 1699, 1616, 1568, 1533, 1408, 1325, 1265, 1203 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.75 (1H, s), 7.70(1H, d, J=8.75), 7.55-7.46 (5H, m), 7.12 (1H, dd, J=1.97, 8.72Hz), 3.75 (2H, s), 2.39 (3H, s).

EXAMPLE 6

[6-CHLORO-2-(4-METHYLBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

15 STEP 1. 6-Chloro-2-(4-methylbenzoyl)-1-(phenylsulfonyl)indole

The title compound was prepared according to the procedure described in step 2 of Example 2 (Method B) from 6-chloro-1-(phenylsulfonyl)indole (step 1 of Example 2, Method B) and *p*-toluoyl chloride.

tlc: R_f=0.3 (ethyl acetate/hexane=1:10).

20 STEP 2. 6-Chloro-2-(4-methylbenzoyl)indole

The title compound was prepared according to the procedure described in step 3 of Example 2 (Method B) from 6-chloro-2-(4-methylbenzoyl)-1-(phenylsulfonyl)indole (step 1).

25 ¹H-NMR (CDCl₃) δ: 9.37 (1H, br s), 7.90 (2H, d, J=8.6Hz), 7.63 (1H, d, J=8.9Hz), 7.48 (1H, s), 7.34 (2H, d, J=8.6Hz), 7.16-7.10 (2H, m), 2.47 (3H, s).

STEP 3. Diethyl α-acetoxy-[6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 6-chloro-2-(4-methylbenzoyl)indole (step 2).

30 ¹H-NMR (CDCl₃) δ: 8.65 (1H, br s), 7.84 (1H, d, J=8.9Hz), 7.76 (2H, d, J=8.2Hz), 7.39-7.37 (1H, m), 7.27 (2H, d, J=8.2Hz), 7.16 (1H, dd, J=1.8, 8.2Hz), 4.36-4.16 (4H,

m), 2.44 (3H, s), 1.72 (3H, s), 1.34-1.22 (6H, m).

STEP 4. Diethyl [6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetoxo-[6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]malonate (step 3).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.83 (1H, br s), 7.76 (1H, d, $J=8.7\text{Hz}$), 7.71 (2H, d, $J=8.2\text{Hz}$), 7.39-7.36 (1H, m), 7.32 (2H, d, $J=8.2\text{Hz}$), 7.14 (1H, dd, $J=2.0, 8.7\text{Hz}$), 5.30 (1H, s), 4.26-4.14 (4H, m), 2.47 (3H, s), 1.22 (6H, t, $J=7.1\text{Hz}$).

STEP 5. [6-Chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]malonate (step 4).

m.p.: 182-184 °C.

IR (KBr) ν : 3321, 1705, 1616, 1602, 1566, 1529, 1431, 1323, 1257, 1230 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.72 (1H, s), 7.74-7.64 (3H, m), 7.49-7.45 (1H, m), 7.39 (2H, d, $J=8.1\text{Hz}$), 7.16-7.09 (1H, m), 3.81 (2H, s), 2.43 (3H, s).

EXAMPLE 7

[6-CHLORO-2-(3-CHLOROBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

STEP 1. 6-Chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole

To a stirred suspension of 6-chloroindole-2-carboxylic acid (H.N.Rydon and J.C.Tweddle, *J.Chem.Soc.*, 1955, 3499., 7.0g, 36 mmol) in thionyl chloride (30 ml) was added dropwise DMF (1 ml). After stirring for 30 min., the mixture was concentrated, and the residue was dissolved in dichloromethane (100 ml) and cooled to 0 °C. To the mixture was added *N,O*-dimethylhydroxylamine hydrochloride (7.0 g, 72 mmol) and pyridine (15 ml). After stirring 2h, the mixture was quenched with water (100 ml), and extracted with dichloromethane (150 ml x 2). The combined organic extracts were washed with 2N aqueous HCl (100 ml), saturated sodium bicarbonate (100 ml), brine (100 ml), and dried (MgSO_4). Removal of solvent afforded 8.2 g (96 %) of the title compound as yellow solids.

$^1\text{H-NMR}$ (CDCl_3) δ : 9.48 (1H, br s), 7.60 (1H, d, $J=8.6\text{Hz}$), 7.46-7.41 (1H, m), 7.22-7.18 (1H, m), 7.11 (1H, dd, $J=8.6, 1.8\text{Hz}$), 3.85 (3H, s), 3.44 (3H, s).

STEP 2. 6-Chloro-2-(3-chlorobenzoyl)indole

To a stirred solution of 6-chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (step 1, 610 mg, 2.56 mmol) and 3-bromochlorobenzene (1.47 g, 7.67 mmol) in THF (20 ml) at -78°C was added dropwise *n*-butyllithium (1.54M in hexane, 4.90 ml, 7.67 mmol). After stirring for 1h, the mixture was quenched with saturated ammonium chloride (20 ml), and extracted with ethyl acetate (50 ml x 2). The combined organic extracts were washed with 2N aqueous HCl (50 ml), saturated sodium bicarbonate (50 ml), brine (50 ml), and dried (MgSO₄). After removal of solvent, the crude product was purified by flash column chromatography eluting with ethyl acetate/hexane (1:10) to afford 565 mg (76 %) of the title compound as pale brown solids.

¹H-NMR (CDCl₃) δ: 9.31 (1H, br s), 7.94 (1H, t, J=1.9Hz), 7.85 (1H, dt, J=1.9, 7.6Hz), 7.65 (1H, d, J=8.4Hz), 7.63-7.58 (1H, m), 7.52-7.45 (2H, m), 7.18-7.12 (2H, m).

STEP 3. Diethyl α-acetoxy-[6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 6-chloro-2-(3-chlorobenzoyl)indole (step 2).

¹H-NMR (CDCl₃) δ: 8.88 (1H, br s), 7.87-7.74 (3H, m), 7.61-7.56 (1H, m), 7.43 (1H, d, J=7.6Hz), 7.41-7.38 (1H, m), 7.20-7.14 (1H, m), 4.30-4.14 (4H, m), 1.74 (3H, s), 1.20 (6H, t, J=8.2Hz).

STEP 4. Diethyl [6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α-acetoxy-[6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]malonate (step 3).

¹H-NMR (CDCl₃) δ: 8.87 (1H, br s), 7.79-7.74 (2H, m), 7.70-7.58 (2H, m), 7.47 (1H, t, J=8.1Hz), 7.39-7.37 (1H, m), 7.15 (1H, dd, J=1.8, 8.7Hz), 5.19 (1H, s), 4.27-4.10 (4H, m), 1.22 (6H, t, J=7.1Hz).

STEP 5. [6-Chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]malonate (step 4).

m.p.: 199-201 °C.

¹H-NMR (DMSO-d₆) δ: 12.25 (1H, br s), 11.81 (1H, s), 7.79-7.57 (5H, m), 7.48 (1H, d, J=1.8Hz), 7.14 (1H, dd, J=1.8, 8.7Hz), 3.81 (2H, s).

EXAMPLE 8

5 METHYL [6-CHLORO-2-(4-CHLOROBENZOYL)-1H-INDOL-3-YL]ACETATE

Method A

STEP 1. Methyl trans 4-chloro-2-(phenylsulfonylamino)cinnamate

To a solution of methyl *trans*-2-amino-4-chlorocinnamate (R.W.Carling et al., *J.Med.Chem.*, 1993, 36, 3397., 30.7 g, 0.15 mol) and pyridine (36 ml, 0.45 mol) in
10 dichloromethane (500 ml) was added benzenesulfonyl chloride (20 ml, 0.16 mol). After stirring for 20 h, methanol (50 ml) was added and the mixture was concentrated. The residual solids were dissolved in dichloromethane (700 ml) and washed with 2N aqueous HCl (150 ml), brine (150 ml) and dried (MgSO₄). After removal of solvent, the residual solids were recrystallized from ethanol to give 40 g (76 %) of the title
15 compound as pale yellow solids.

¹H-NMR (CDCl₃) δ: 7.77-7.71 (2H, m), 7.59-7.52 (1H, m), 7.48-7.35 (5H, m), 7.20 (1H, dd, J=2.0, 8.4 Hz), 6.85 (1H, br s), 6.15 (1H, d, J=15.8 Hz), 3.78 (3H, s).

STEP 2. methyl [6-chloro-2-(4-chlorobenzoyl)-1-(phenylsulfonyl)indolin-3-yl]acetate

A mixture of methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1,
20 1.1 g, 3.1 mmol), 4-chlorophenacylbromide (1.1 g, 4.6 mmol) and potassium carbonate (2.1 g, 15.4 mmol) in acetone (10 ml) was stirred at room temperature for 1.5 h. The mixture was filtered and the filtrate concentrated. The residual solids were recrystallized from ethyl acetate/ hexane to afford 0.91 g (59%) of the title compound as pale yellow solids.

25 ¹H-NMR (CDCl₃) δ: 7.97-7.94 (2H, m), 7.86-7.81 (2H, m), 7.65-7.58 (1H, m), 7.55-7.46 (5H, m), 7.02 (1H, dd, J=2.0, 8.2 Hz), 6.90 (1H, d, J=8.2 Hz), 5.99 (1H, d, J=9.7 Hz), 4.00-3.87 (1H, m), 3.41 (3H, s), 2.63 (1H, dd, J=6.3, 17.6 Hz), 2.51 (1H, dd, J=6.3, 17.6 Hz).

STEP 3. Methyl [6-Chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate

30 To a stirred solution of methyl [6-chloro-2-(4-chlorobenzoyl)-1-(phenylsulfonyl)indolin-3-yl]acetate (step 2, 50 mg, 0.10 mmol) in THF was added 1,8-

diazabicyclo[5.4.0]undec-7-ene (DBU, 30 μ l, 0.20 mmol). After stirring for 15h, the mixture was quenched with water (30 ml), and extracted with ethyl acetate (50 ml). The extract was washed with 2N aqueous HCl (30 ml), saturated sodium bicarbonate (30 ml), brine (30 ml), and dried (MgSO₄). Removal of solvent afforded 34 mg
5 (93 %) of the title compound as yellow solids.

MS (EI) m/z : 361 (M⁺).

¹H-NMR (CDCl₃) δ : 8.90 (1H, br s), 7.74 (2H, d, J=8.6Hz), 7.56 (1H, d, J=8.6Hz), 7.49 (2H, d, J=8.6Hz), 7.36 (1H, d, J=1.6Hz), 7.16 (1H, dd, J=1.6, 8.6Hz), 3.81 (2H, s), 3.66 (3H, s).

10 Method B

A mixture of methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Method A, 0.70 g, 2.0 mmol), 4-chlorophenacyl bromide (0.51 g, 2.2 mmol), and potassium carbonate (0.83 g, 6.0 mmol) in acetone (20 ml) was stirred at room temperature. After stirring for 2h, cesium carbonate (2.0 g, 6.0 mmol) was added and
15 the stirring was continued for an additional 4h. The mixture was concentrated and the residue was diluted in water (100 ml). The aqueous mixture was extracted with ethyl acetate (100 ml x 2). The combined organic extracts were washed with 2N aqueous HCl (100 ml), saturated sodium bicarbonate (100 ml), brine (100 ml), and dried (MgSO₄). After removal of solvent, the solids were recrystallized from ethanol to
20 afford 0.44 g (61 %) of the title compound.

MS and NMR spectra were identical with those of the compound prepared in step 3.

EXAMPLE 9

[6-CHLORO-2-(4-CHLOROBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

Method A

25 STEP 1. 6-Chloro-2-(4-chlorobenzoyl)indole

The title compound was prepared according to the procedure described in step 2 of Example 7 from 6-chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (Example 7, step 1) and 4-bromochlorobenzene.

¹H-NMR (DMSO-d₆) δ : 12.14 (1H, br s), 7.97 (2H, d, J=8.4Hz), 7.76 (1H, d, J=8.6Hz),
30 7.66 (2H, d, J=8.4Hz), 7.54-7.50 (1H, m), 7.20 (1H, s), 7.14 (1H, dd, J=2.0, 8.6Hz).

STEP 2. Diethyl α -acetoxy-[6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 6-chloro-2-(4-chlorobenzoyl)indole (step 1).

¹H-NMR (CDCl₃) δ: 8.83 (1H, br s), 7.87-7.77 (3H, m), 7.48-7.36 (3H, m), 7.17 (1H, dd, J=2.0, 8.9Hz), 4.28-4.14 (4H, m), 1.73 (3H, s), 1.20 (6H, t, J=7.1Hz).

5 **STEP 3. Diethyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]malonate**

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α-acetoxy-[6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]malonate (step 2).

¹H-NMR (CDCl₃) δ: 8.91 (1H, br s), 7.75 (2H, d, J=8.6Hz), 7.75-7.69 (1H, m), 7.50
10 (2H, d, J=8.6Hz), 7.29 (1H, d, J=1.8Hz), 7.13 (1H, dd, J=1.8, 8.7Hz), 5.23 (1H, s), 4.28-4.07 (4H, m), 1.23 (6H, t, J=7.2Hz).

STEP 4. [6-Chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]malonate (step 3).
15

m.p.: 189-190 °C.

IR (KBr) ν: 3309, 1699, 1616, 1525, 1431, 1325, 1255, 1226, 1091 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.78 (1H, s), 7.80-7.72 (3H, m), 7.65 (2H, d, J=8.6Hz), 7.47
(1H, d, J=1.8Hz), 7.13 (1H, dd, J=1.8, 8.7Hz), 3.83 (2H, s).

20 **Method B**

[6-Chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid

A mixture of methyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate (Example 8, 1.80 g) and 2N aqueous NaOH (7.5 ml) in MeOH-THF (10 ml-10 ml) was stirred at 80 °C for 1h. The mixture was cooled and concentrated. The residue was
25 dissolved in water (150 ml) and washed with diethyl ether (50 ml). The aqueous layer was acidified with 2N aqueous HCl (10 ml), and extracted with ethyl acetate (100 ml x 2). The combined organic extracts were washed with brine (50 ml), dried (MgSO₄), and concentrated. The residual solids were recrystallized from toluene to afford 1.58 g (91 %) of the title compound.

30 IR and NMR spectra were identical with those of the compound prepared by Method A.

EXAMPLE 10**[6-CHLORO-2-(3-FLUOROBENZOYL)-1H-INDOL-3-YL]ACETIC ACID****STEP 1. 6-Chloro-2-(3-fluorobenzoyl)indole**

The title compound was prepared according to the procedure described in step 2 of Example 7 from 6-chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (Example 7, step 1) and 3-bromofluorobenzene.

¹H-NMR (CDCl₃) δ: 9.28 (1H, br s), 7.79-7.75 (1H, m), 7.68-7.63 (2H, m), 7.56-7.48 (2H, m), 7.36-7.30 (1H, m), 7.17-7.14 (2H, m).

STEP 2. Diethyl α-acetoxy-[6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 6-chloro-2-(3-fluorobenzoyl)indole (step 1).

¹H-NMR (CDCl₃) δ: 9.15 (1H, br s), 8.83 (1H, d, J=8.72Hz), 7.66-7.27 (5H, m), 7.17 (1H, dd, J=2.00, 8.72Hz), 4.25-4.13 (4H, m), 1.75 (3H, s), 1.19 (6H, t, J=7.07Hz).

STEP 3. Diethyl [6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α-acetoxy-[6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]malonate (step 2).

¹H-NMR (CDCl₃) δ: 8.91 (1H, br s), 7.77-7.12 (7H, m), 5.21 (1H, s), 4.25-4.11 (4H, m), 1.22 (6H, t, J=7.07Hz).

STEP 4. [6-Chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]malonate (step 3).

m.p.: 278-281 °C.

IR (KBr) ν: 3385, 1697, 1638, 1583, 1541, 1508, 1420, 1400, 1315, 1261, 1236 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.79 (1H, s), 7.76-7.48 (7H, m), 7.13 (1H, dd, J=1.97, 8.56Hz), 3.80 (2H, s).

EXAMPLE 11**[6-CHLORO-2-(4-FLUOROBENZOYL)-1H-INDOL-3-YL]ACETIC ACID****STEP 1. 6-Chloro-2-(4-fluorobenzoyl)indole**

The title compound was prepared according to the procedure described in step 2 of Example 7 from 6-chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (Example 7, step 1) and 4-bromofluorobenzene.

¹H-NMR (CDCl₃) δ: 9.29 (1H, br s), 8.05-8.00 (2H, m), 7.64 (1H, d, 8.72Hz), 7.48-7.11 (5H, m).

STEP 2. Diethyl α-acetoxy-[6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 6-chloro-2-(4-fluorobenzoyl)indole (step 1).

¹H-NMR (CDCl₃) δ: 9.20 (1H, br s), 7.92-7.11 (7H, m), 4.25-4.14 (4H, m), 1.73 (3H, s), 1.20 (6H, t, J=7.10Hz).

STEP 3. Diethyl [6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α-acetoxy-[6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]malonate (step 2).

¹H-NMR (CDCl₃) δ: 9.01 (1H, br s), 7.87-7.69 (2H, m), 7.71 (2H, d, J=8.91 Hz), 7.27-7.10 (4H, m), 5.25 (1H, s), 4.26-4.11 (4H, m), 1.22 (6H, t, J=7.07Hz).

STEP 4. [6-Chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]malonate (step 3).

m.p.: 181-183 °C.

IR (KBr) ν: 3309, 1701, 1616, 1601, 1566, 1527, 1508, 1419, 1323, 1257, 1229 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.75 (1H, s), 7.87-7.82 (2H, m), 7.73 (1H, d, J=8.72), 7.47-7.36 (3H, m), 7.12 (1H, dd, J=1.84, 8.72Hz), 3.78 (2H, s).

EXAMPLE 12

[2-(3-BROMOBENZOYL)-6-CHLORO-1H-INDOL-3-YL]ACETIC ACID

STEP 1. 2-(3-Bromobenzoyl)-6-chloroindole

The title compound was prepared according to the procedure described in step 2 of Example 7 from 6-chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (Example 7, step 1) and 3-bromoiodobenzene.

¹H-NMR (CDCl₃) δ: 9.28 (1H, br s), 8.11-8.08 (1H, m), 7.93-7.87 (1H, m), 7.79-7.73 (1H, m), 7.65 (1H, d, J=8.6Hz), 7.50-7.38 (2H, m), 7.19-7.01 (2H, m).

STEP 2. Diethyl α-acetoxy-[2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 2-(3-bromobenzoyl)-6-chloroindole (step 1).

¹H-NMR (CDCl₃) δ: 8.73 (1H, br s), 7.93 (1H, t, J=1.8Hz), 7.85 (1H, d, J=8.6Hz), 7.82-7.72 (2H, m), 7.42-7.32 (2H, m), 7.18 (1H, dd, J=2.0, 8.6Hz), 4.32-4.16 (4H, m), 1.75 (3H, s), 1.20 (6H, t, J=7.1Hz).

STEP 3. Diethyl [2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α-acetoxy-[2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]malonate (step 2).

¹H-NMR (CDCl₃) δ: 8.88 (1H, br s), 7.92-7.90 (1H, m), 7.80-7.68 (3H, m), 7.45-7.36 (2H, m), 7.15 (1H, dd, J=1.8, 8.7Hz), 5.20 (1H, s), 4.30-4.10 (4H, m), 1.23 (6H, t, J=7.2Hz).

STEP 4. [2-(3-Bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]malonate (step 3).

m.p.: 215-218 °C.

IR (KBr) ν: 3369, 1710, 1604, 1558, 1533, 1423, 1319, 1253, 1228 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.81 (1H, s), 7.92-7.83 (2H, m), 7.78-7.70 (2H, m), 7.54 (1H, t, J=7.8Hz), 7.48 (1H, d, J=2.0Hz), 7.13 (1H, dd, J=2.0, 8.7Hz), 3.80 (2H, s).

EXAMPLE 13

[2-(4-BROMOBENZOYL)-6-CHLORO-1H-INDOL-3-YL]ACETIC ACID

STEP 1. 2-(4-Bromobenzoyl)-6-chloroindole

The title compound was prepared according to the procedure described in step 2 of Example 7 from 6-chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (Example 7, step 1) and 4-bromoiodobenzene.

¹H-NMR (DMSO-d₆) δ: 12.28 (1H, br s), 7.88 (2H, d, J=8.7Hz), 7.80 (2H, d, J=8.7Hz),

7.76 (1H, d, J=8.7Hz), 7.53-7.50 (1H, m), 7.22-7.19 (1H, m), 7.13 (1H, dd, J=2.0, 8.7Hz).

STEP 2. Diethyl α -acetoxy-[2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 2-(4-bromobenzoyl)-6-chloroindole (step 1).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.72 (1H, br s), 7.84 (1H, d, J=8.7Hz), 7.73 (2H, d, J=8.4Hz), 7.62 (2H, d, J=8.4Hz), 7.38 (1H, d, J=1.8Hz), 7.18 (1H, dd, J=1.8, 8.7Hz), 4.28-4.14 (4H, m), 1.73 (3H, s), 1.20 (6H, t, J=7.1Hz).

STEP 3. Diethyl [2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetoxy-[2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]malonate (step 2).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.86 (1H, br s), 7.75 (1H, d, J=8.4Hz), 7.67 (4H, s), 7.37-7.32 (1H, m), 7.18-7.12 (1H, m), 5.22 (1H, s), 4.26-4.10 (4H, m), 1.23 (6H, t, J=7.1Hz).

STEP 4. [2-(4-Bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]malonate (step 3).

m.p.: 199-201 °C.

IR (KBr) ν : 3300, 1699, 1618, 1587, 1525, 1433, 1406, 1325, 1255, 1226 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.78 (1H, s), 7.80 (2H, d, J=8.4Hz), 7.75 (1H, d, J=8.6Hz), 7.69 (2H, d, J=8.4Hz), 7.47 (1H, d, J=1.8Hz), 7.13 (1H, dd, J=1.8, 8.6Hz), 3.84 (2H, s).

EXAMPLE 14

[6-CHLORO-2-(3-TRIFLUOROMETHYLBENZOYL)-1H-INDOL-3-YL]ACETIC

ACID

STEP 1. 6-Chloro-2-(3-trifluoromethylbenzoyl)indole

The title compound was prepared according to the procedure described in step 2 of Example 7 from 6-chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (Example 7, step 1) and 3-bromobenzotrifluoride.

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.23 (1H, br s), 8.28-8.04 (3H, m), 7.89-7.75 (2H, m), 7.55-

7.51 (1H, m), 7.22-7.12 (2H, m).

STEP 2. Diethyl α -acetoxy-[6-chloro-2-(3-trifluoromethylbenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 6-chloro-2-(3-trifluoromethylbenzoyl)indole (step 1).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.82 (1H, br s), 8.25-7.80 (4H, m), 7.68-7.57 (1H, m), 7.43-7.41 (1H, m), 7.22-7.16 (1H, m), 4.36-4.20 (4H, m), 1.67 (3H, s), 1.20 (6H, t, $J=7.1\text{Hz}$).

STEP 3. Diethyl [6-chloro-2-(3-trifluoromethylbenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetoxy-[6-chloro-2-(3-trifluoromethylbenzoyl)-1H-indol-3-yl]malonate (step 2).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.91 (1H, br s), 8.08-8.03 (1H, m), 8.02-7.87 (2H, m), 7.76 (1H, d, $J=9.0\text{Hz}$), 7.73-7.64 (1H, m), 7.39-7.36 (1H, m), 7.15 (1H, dd, $J=1.8, 9.0\text{Hz}$), 5.15 (1H, s), 4.26-4.10 (4H, m), 1.22 (6H, t, $J=7.2\text{Hz}$).

STEP 4. [6-Chloro-2-(3-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [6-chloro-2-(3-trifluoromethylbenzoyl)-1H-indol-3-yl]malonate (step 3).

m.p.: 194-196 °C.

IR (KBr) ν : 3371, 1705, 1631, 1421, 1307, 1228, 1168, 1122, 1072 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.86 (1H, s), 8.10-7.98 (3H, m), 7.83 (1H, t, $J=7.7\text{Hz}$), 7.75 (1H, d, $J=8.6\text{Hz}$), 7.49 (1H, d, $J=2.0\text{Hz}$), 7.15 (1H, dd, $J=2.0, 8.6\text{Hz}$), 3.80 (2H, s).

EXAMPLE 15

[6-CHLORO-2-(4-TRIFLUOROMETHYLBENZOYL)-1H-INDOL-3-YL]ACETIC

ACID

STEP 1. 6-Chloro-1-(phenylsulfonyl)-2-(4-trifluoromethylbenzoyl)indole

The title compound was prepared according to the procedure described in step 2 of Example 2 (Method B) from 6-chloro-1-(phenylsulfonyl)indole (step 1 of Example 2, Method B) and 4-trifluoromethylbenzoyl chloride.

tlc: $R_f=0.2$ (ethyl acetate/hexane=1:4).

STEP 2. 6-Chloro-2-(4-trifluoromethylbenzoyl)indole

The title compound was prepared according to the procedure described in step 3 of Example 2 (Method B) from 6-chloro-1-phenylsulfonyl-2-(4-trifluoromethylbenzoyl) indole (step 1).

- 5 $^1\text{H-NMR}$ (CDCl_3) δ : 9.53 (1H, br s), 7.94 (4H, dd, $J=8.24$, 69.1Hz), 7.63 (1H, d, $J=8.56\text{Hz}$), 7.49-7.11 (3H, m).

STEP 3. Diethyl α -acetoxy-[6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]malonate

- 10 The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 6-chloro-2-(4-trifluoromethylbenzoyl)indole (step 2).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.66 (1H, br s), 7.98 (2H, d, $J=8.24\text{Hz}$), 7.87 (1H, d, $J=8.91\text{Hz}$), 7.75 (2H, d, $J=8.07\text{Hz}$), 7.39 (1H, d, $J=1.81\text{Hz}$), 7.19 (1H, dd, $J=1.81$, 7.10Hz), 4.36-4.16 (4H, m), 1.70 (3H, s), 1.34-1.22 (6H, m).

STEP 4. Diethyl [6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]malonate

- 15 The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetoxy-[6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]malonate (step 3).

$^1\text{H-NMR}$ (CDCl_3) δ : 9.24 (1H, br s), 7.84 (4H, dd, $J=7.91$, 25.7Hz), 7.69-7.09 (3H, m), 5.24 (1H, s), 4.21-4.06 (4H, m), 1.21 (6H, t, $J=7.07\text{Hz}$).

- 20 STEP 5. [6-Chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]malonate (step 4).

- 25 $^1\text{H-NMR}$ (DMSO-d_6) δ : 11.78 (1H, s), 7.94 (4H, s), 7.76 (1H, d, $J=8.72\text{Hz}$), 7.47 (1H, m), 7.14 (1H, d, $J=1.81$, 8.72), 3.81 (2H, s).

EXAMPLE 16[6-CHLORO-2-(3,4-DICHLOROBENZOYL)-1H-INDOL-3-YL]ACETIC ACIDSTEP 1. 6-Chloro-2-(3,4-dichlorobenzoyl)indole

The title compound was prepared according to the procedure described in step 2 of Example 7 from 6-chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (Example 7, step 1) and 1-bromo-3,4-dichlorobenzene.

tlc: R_f=0.7 (ethyl acetate/hexane=1:3).

5 STEP 2. Diethyl α-acetoxy-[6-chloro-2-(3,4-dichlorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 6-chloro-2-(3,4-dichlorobenzoyl)indole (step 1).

¹H-NMR (CDCl₃) δ: 8.80 (1H, br s), 7.90-7.79 (2H, m), 7.71 (1H, dd, J=2.0, 8.4Hz), 7.57 (1H, d, J=8.4Hz), 7.40-7.35 (1H, m), 7.18 (1H, dd, J=1.8, 8.8Hz), 4.30-4.14 (4H, m), 1.77 (3H, s), 1.20 (6H, t, J=7.1Hz).

10 STEP 3. Diethyl [6-chloro-2-(3,4-dichlorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α-acetoxy-[6-chloro-2-(3,4-dichlorobenzoyl)-1H-indol-3-yl]malonate (step 2).

15 ¹H-NMR (CDCl₃) δ: 8.95 (1H, br s), 7.88 (1H, d, J=1.6Hz), 7.72 (1H, d, J=8.7Hz), 7.68-7.58 (2H, m), 7.28 (1H, d, J=1.6Hz), 7.14 (1H, dd, J=1.6, 8.7Hz), 5.18 (1H, s), 4.28-4.10 (4H, m), 1.23 (6H, t, J=7.1Hz).

STEP 4. [6-Chloro-2-(3,4-dichlorobenzoyl)-1H-indol-3-yl]acetic acid

20 The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [6-chloro-2-(3,4-dichlorobenzoyl)-1H-indol-3-yl]malonate (step 3).

m.p.: 206-209 °C.

IR (KBr) ν: 3435, 1708, 1620, 1583, 1525, 1423, 1384, 1301, 1263, 1228 cm⁻¹.

25 ¹H-NMR (DMSO-d₆) δ: 11.62 (1H, br s), 7.95-7.75 (3H, m), 7.68 (1H, d, J=8.7Hz), 7.42 (1H, d, J=2.0Hz), 7.02 (1H, dd, J=2.0, 8.7Hz), 3.50 (2H, s).

EXAMPLE 17

(2-BENZOYL-4-CHLORO-1H-INDOL-3-YL)ACETIC ACID

STEP 1. 4-Chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole

30 The title compound was prepared according to the procedure described in step 1 of Example 7 employing 4-chloroindole-2-carboxylic acid (F. C. Uhle, *J. Amer. Chem. Soc.*, 1949, 71, 761).

¹H-NMR (CDCl₃) δ: 9.56 (1H, br s), 7.36-7.29 (2H, m), 7.24-7.12 (2H, m), 3.88 (3H, s), 3.45 (3H, s).

STEP 2. 2-Benzoyl-4-chloroindole

To a solution of 4-chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (step 1, 3.4 g, 0.014 mol) in THF (60 ml) cooled to -78 °C was added dropwise phenyllithium (1.8 M in cyclohexane/ether (7:3), 30 ml, 0.070 mol). After stirring for 1 h, the mixture was poured into water (80 ml) and extracted with ethyl acetate (80 ml x 2). After drying (MgSO₄) and removal of solvent, the crude product was purified by flash column chromatography eluting with ethyl acetate/hexane (1:10) to afford 3.6 g (100 %) of the title compound as white solids.

¹H-NMR (CDCl₃) δ: 9.46 (1H, br s), 8.05-8.00 (2H, m), 7.73-7.51 (3H, m), 7.40 (1H, dd, J=1.0, 8.3 Hz), 7.33-7.24 (2H, m), 7.18 (1H, d, J=7.6 Hz).

STEP 3. Diethyl α-acetoxy-(2-benzoyl-4-chloro-1H-indol-3-yl)malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 2-benzoyl-4-chloroindole (step 2).

¹H-NMR (CDCl₃) δ: 9.49 (1H, br s), 7.98-7.92 (2H, m), 7.66-7.57 (1H, m), 7.53-7.44 (2H, m), 7.34 (1H, dd, J=2.0, 7.1 Hz), 7.21-7.15 (2H, m), 4.30-3.90 (4H, m), 2.08 (3H, s), 1.15 (6H, t, J=7.3 Hz).

STEP 4. Diethyl (2-benzoyl-4-chloro-1H-indol-3-yl)malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 from diethyl α-acetoxy-(2-benzoyl-4-chloro-1H-indol-3-yl)malonate (step 3).

¹H-NMR (CDCl₃) δ: 8.88 (1H, br s), 7.83-7.77 (2H, m), 7.64-7.58 (1H, m), 7.55-7.46 (2H, m), 7.25-7.05 (3H, m), 5.86 (1H, s), 4.25-4.08 (4H, m), 1.23 (6H, t, J=7.3 Hz).

STEP 5. (2-Benzoyl-4-chloro-1H-indol-3-yl)acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl (2-benzoyl-4-chloro-1H-indol-3-yl)malonate. m.p.: 206-209 °C (recrystallized from ethyl acetate/hexane).

IR (KBr) ν: 1700, 1575, 1245 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.20 (1H, br s), 11.98 (1H, s), 7.85-7.67 (3H, m), 7.63-7.55 (2H, m), 7.45 (1H, d, J=8.2 Hz), 7.25 (1H, t, J=8.2 Hz), 7.12 (1H, d, J=7.6 Hz), 4.02 (2H, s).

EXAMPLE 18

5 [5-CHLORO-2-(3-METHYLBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

STEP1. 5-Chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole

The title compound was prepared according to the procedure described in step 1 of Example 7 from 5-chloroindole-2-carboxylic acid.

¹H-NMR (CDCl₃) δ: 9.68 (1H, br s), 7.68-7.65 (1H, m), 7.37 (1H, d, J=8.7 Hz), 7.23 (1H, d, J=1.6 Hz), 7.18-7.15 (1H, m), 3.85 (3H, s), 2.05 (3H, s).

STEP2. 5-Chloro-2-(3-methylbenzoyl)indole

The title compound was prepared according to the procedure described in step 2 of Example 17 from 5-chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (step 1) and 3-methylphenyllithium.

15 m.p.: 197.5-198 °C (recrystallized from ethyl acetate/hexane).

IR (KBr) v: 3310, 1626, 1603, 1583, 1516, 1406, 1377, 1337, 1269, 1178, 1134 cm⁻¹.

¹H-NMR (CDCl₃) δ: 9.39 (1H, br s), 7.28-7.76 (2H, m), 7.71-7.68 (1H, m), 7.46-7.38 (3H, m), 7.32 (1H, dd, J=8.7, 2.0 Hz), 7.08 (1H, dd, J=2.0, 0.8 Hz), 2.47 (3H, s).

STEP3. Diethyl α-acetoxy-[5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]malonate

20 The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 5-chloro-2-(3-methylbenzoyl)indole (step 2).

m.p.: 173-174 °C (recrystallized from ethyl acetate/hexane).

¹H-NMR (CDCl₃) δ: 8.64 (1H, br s), 7.91 (1H, br s), 7.69-7.61 (2H, m), 7.46-7.22 (4H, m), 4.30-4.16 (4H, m), 2.39 (3H, s), 1.72 (3H, s), 1.23 (3H, t, J=7.2 Hz).

STEP4. Diethyl [5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α-acetoxy-[5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl] malonate (step 3).

m.p.: 143-144 °C (recrystallized from ethyl acetate/hexane).

30 ¹H-NMR (CDCl₃) δ: 8.88 (1H, br s), 7.84-7.80 (1H, m), 7.62-7.55 (2H, m), 7.48-7.36

(2H, m), 7.32-7.28 (2H, m), 5.27 (1H, s), 4.30-4.10 (4H, m), 2.42 (3H, s), 1.24 (3H, t, J=7.1 Hz).

STEP 5. [5-Chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]malonate.

m.p.: 241-242 °C (recrystallized from ethyl acetate/hexane).

IR (KBr) ν : 3321, 1703, 1618, 1535, 1431, 1335, 1232, 1016, 808, 758 cm^{-1} .

$^1\text{H-NMR}$ (DMSO- d_6) δ : 11.7 (1H, br s), 7.62 (1H, d, J=1.8 Hz), 7.45-7.30 (6H, m), 7.19 (1H, dd, J=8.6, 1.8 Hz), 3.63 (2H, s), 2.27 (3H, s).

EXAMPLE 19

[5-CHLORO-2-(4-CHLOROBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

STEP 1. 5-Chloro-2-(4-chlorobenzoyl)indole

The title compound was prepared according to the procedure described in step 2 of Example 7 from 5-chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (Example 18, step 1) and 4-bromochlorobenzene.

$^1\text{H-NMR}$ (CDCl_3) δ : 9.32 (1H, br s), 7.94 (2H, d, J=8.4Hz), 7.70 (1H, s), 7.53 (2H, d, J=8.4Hz), 7.42 (1H, d, J=8.9Hz), 7.34 (1H, dd, J=2.0, 8.9Hz), 7.07 (1H, s).

STEP 2. Diethyl α -acetox-[5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 5-chloro-2-(4-chlorobenzoyl)indole (step 1).

$^1\text{H-NMR}$ (CDCl_3) δ : 9.11 (1H, br s), 7.87 (1H, s), 7.76 (2H, d, J=8.6Hz), 7.41 (2H, d, J=8.6Hz), 7.27 (1H, d, J=8.7Hz), 7.22 (1H, dd, J=1.8, 8.7Hz), 4.25-4.14 (4H, m), 1.72 (3H, s), 1.24-1.19 (6H, m).

STEP 3. Diethyl [5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetox-[5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]malonate (step 2).

$^1\text{H-NMR}$ (CDCl_3) δ : 9.13 (1H, br s), 7.73 (2H, d, J=8.4Hz), 7.68 (1H, d, J=2.0Hz), 7.48 (2H, d, J=8.4Hz), 7.11 (1H, dd, J=2.0, 8.9Hz), 7.02 (1H, d, J=8.9Hz), 5.28 (1H, s),

4.24-4.03 (4H, m), 1.27-1.21 (6H, m).

STEP 4. [5-Chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]malonate (step 3).

m.p.: 220-224.°C (recrystallized from ethyl acetate/hexane).

IR (KBr) ν : 3321, 1618, 1535, 1379, 1339, 1263, 1130, 1090, 1057, 1007 cm^{-1} .

$^1\text{H-NMR}$ (DMSO- d_6) δ : 11.83 (1H, s), 7.80-7.75 (3H, m), 7.67-7.62 (2H, m), 7.48 (1H, d, $J=8.7\text{Hz}$), 7.31 (1H, dd, $J=2.0, 8.7\text{Hz}$), 3.84 (2H, s).

10 EXAMPLE 20

[5-CHLORO-2-(3-CHLOROBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

STEP 1. 5-Chloro-2-(3-chlorobenzoyl)indole

The title compound was prepared according to the procedure described in step 2 of Example 7 from 5-chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (Example 15 18, step 1) and 3-bromochlorobenzene.

$^1\text{H-NMR}$ (CDCl_3) δ : 9.28 (1H, br s), 7.95 (1H, t, $J=1.7\text{Hz}$), 7.88-7.84 (1H, m), 7.71 (1H, d, $J=2.0\text{Hz}$), 7.63-7.59 (1H, m), 7.49 (1H, t, $J=7.7\text{Hz}$), 7.42 (1H, d, $J=8.7\text{Hz}$), 7.34 (1H, dd, $J=2.0, 8.7\text{Hz}$), 7.10-7.09 (1H, m).

STEP 2. Diethyl α -acetoxy-[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]malonate

20 The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 5-chloro-2-(3-chlorobenzoyl)indole (step 1).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.94 (1H, br s), 7.91 (1H, d, $J=1.8\text{Hz}$), 7.79-7.76 (1H, m), 7.74 (1H, d, $J=7.7\text{Hz}$), 7.60-7.56 (1H, m), 7.42 (1H, t, $J=7.7\text{Hz}$), 7.33 (1H, d, $J=8.7\text{Hz}$), 7.27 (1H, dd, $J=1.8, 8.7\text{Hz}$), 4.37-4.19 (4H, m), 1.75 (3H, s), 1.26-1.20 (6H, m).

25 STEP 3. Diethyl [5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetoxy-[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]malonate (step 2).

$^1\text{H-NMR}$ (CDCl_3) δ : 9.04 (1H, br.s), 7.77-7.75 (2H, m), 7.68-7.58 (2H, m), 7.46 (1H, t, $J=7.7\text{Hz}$), 7.24-7.20 (2H, m), 5.23 (1H, s), 4.27-4.14 (4H, m), 1.27-1.22 (6H, m).

30

STEP 4. [5-Chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]malonate (step 3).

5 m.p.: 243-247 °C (recrystallized from ethyl acetate/hexane).

IR (KBr) ν : 3329, 1707, 1618, 1535, 1431, 1406, 1375, 1333, 1279, 1232, 1053 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.87 (1H, s), 7.81-7.68 (4H, m), 7.64-7.58 (1H, m), 7.49 (1H, d, $J=8.9\text{Hz}$), 7.32 (1H, dd, $J=2.0, 8.9\text{Hz}$), 3.82 (2H, s).

EXAMPLE 21

10 [2-(4-CHLOROBENZOYL)-5-FLUORO-1H-INDOL-3-YL]ACETIC ACID

STEP 1. 5-fluoro-2-[(N-methoxy-N-methylamino)carbonyl]indole

The title compound was prepared according to the procedure described in step 1 of Example 7 from 5-fluoroindole-2-carboxylic acid.

$^1\text{H-NMR}$ (CDCl_3) δ : 10.15 (1H, br s), 7.41-7.36 (1H, m), 7.32 (1H, dd, $J=2.5, 9.1\text{Hz}$),
15 7.20-7.19 (1H, m), 7.09-7.01 (1H, m), 3.84 (3H, s), 3.47 (3H, s).

STEP 2. 2-(4-Chlorobenzoyl)-5-fluoro-indole

The title compound was prepared according to the procedure described in step 2 of Example 7 from 5-fluoro-2-[(N-methoxy-N-methylamino)carbonyl]indole (step 1) and 4-bromochlorobenzene.

20 $^1\text{H-NMR}$ (CDCl_3) δ : 9.27 (1H, br s), 7.94 (2H, d, $J=8.4\text{Hz}$), 7.52 (2H, d, $J=8.4\text{Hz}$), 7.45-7.40 (1H, m), 7.37-7.33 (1H, m), 7.21-7.12 (1H, m), 7.10-7.09 (1H, m).

STEP 3. Diethyl α -acetoxy-[2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 2-(4-chlorobenzoyl)-5-fluoroindole (step 2).

25 $^1\text{H-NMR}$ (CDCl_3) δ : 9.01 (1H, br s), 7.80-7.77 (2H, m), 7.58-7.54 (1H, m), 7.45-7.41 (2H, m), 7.36-7.27 (1H, m), 7.12-7.01 (1H, m), 4.29-4.15 (4H, m), 1.74 (3H, s), 1.28-1.17 (6H, m).

STEP 4. Diethyl [2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]malonate

30 The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetoxy-[2-(4-chlorobenzoyl)-5-fluoro-1H-

indol-3-yl]malonate (step 3).

¹H-NMR (CDCl₃) δ: 8.98 (1H, br s), 7.77-7.72 (2H, m), 7.51-7.46 (2H, m), 7.40 (1H, dd, J= 2.5, 9.7Hz), 7.18-7.13 (1H, m), 7.04-6.96 (1H, m), 5.28 (1H, s), 4.26-4.07 (4H, m), 1.30-1.18 (6H, m).

5 STEP 5. [2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]malonate (step 4).

m.p.: 233-238 °C (recrystallized from ethyl acetate/hexane).

10 IR (KBr) ν: 3317, 1707, 1624, 1609, 1587, 1526, 1458, 1408, 1344, 1263, 1242 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.73 (1H, s), 7.77 (2H, d, J=8.6Hz), 7.65 (2H, d, J=8.6Hz), 7.52-7.44 (2H, m), 7.22-7.15 (1H, m), 3.84 (2H, s).

EXAMPLE 22

[2-(3-CHLOROBENZOYL)-5-FLUORO-1H-INDOL-3-YL]ACETIC ACID

15 STEP 1. 2-(3-chlorobenzoyl)-5-fluoroindole

The title compound was prepared according to the procedure described in step 2 of Example 7 from 5-fluoro-2-[(N-methoxy-N-methylamino)carbonyl]indole (Example 21, step 1) and 3-bromochlorobenzene.

¹H-NMR (CDCl₃) δ: 9.29 (1H, m), 7.96-7.94 (1H, m), 7.88-7.84 (1H, m), 7.63-7.59 (1H, m), 7.48 (1H, t, J=8.0Hz), 7.45-7.40 (1H, m), 7.36 (1H, dd, J=2.6, 8.6Hz), 7.20-7.11 (2H, m).

STEP 2. Diethyl α-acetoxy-[2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 2-(3-chlorobenzoyl)-5-fluoroindole (step 1).

25 ¹H-NMR (CDCl₃) δ: 8.73 (1H, br s), 7.79-7.74 (2H, m), 7.61-7.56 (2H, m), 7.43 (1H, t, J=7.6Hz), 7.37-7.32 (1H, m), 7.13-7.06 (1H, m), 4.34-4.20 (4H, m), 1.76 (3H, s), 1.33-1.20 (6H, m).

STEP 3. Diethyl [2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α-acetoxy-[2-(3-chlorobenzoyl)-5-fluoro-1H-

30

indol-3-yl]malonate (step 2).

¹H-NMR (CDCl₃) δ: 9.05 (1H, br s), 7.76-7.75 (1H, m), 7.68-7.64 (1H, m), 7.61-7.56 (1H, m), 7.48-7.40 (2H, m), 7.24-7.19 (1H, m), 7.08-7.00 (1H, m), 5.25 (1H, s), 4.28-4.07 (4H, m), 1.33-1.21 (6H, m).

5 STEP 4. [2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]malonate (step 3).

m.p.: 208-212 °C (recrystallized from ethyl acetate/hexane).

10 IR (KBr) v: 3337, 1709, 1618, 1560, 1529, 1477, 1458, 1427, 1408, 1335, 1304 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.78 (1H, s), 7.77-7.58 (4H, m), 7.52-7.46 (2H, m), 7.24-7.16 (1H, m), 3.82 (2H, s).

EXAMPLE 23

[5-METHOXY-2-(3-METHYLBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

15 STEP 1. 5-methoxy-2-[(N-methoxy-N-methylamino)carbonyl]indole

The title compound was prepared according to the procedure described in step 1 of Example 7 from 5-methoxyindole-2-carboxylic acid.

¹H-NMR (CDCl₃) δ: 9.29 - 9.13 (1H, br), 7.33 (1H, d, J=8.9Hz), 7.19 - 7.14 (1H, m), 7.10 (1H, d, J=2.3Hz), 6.98 (1H, dd, J=8.9, 2.3Hz), 3.86 (3H, s), 3.84 (3H, s), 3.42 (3H, s).

20

STEP 2. 5-Methoxy-2-(3-methylbenzoyl)indole

The title compound was prepared according to the procedure described in step 2 of Example 17 from 5-methoxy-2-[(N-methoxy-N-methylamino)carbonyl]indole (step 1) and 3-methylphenyllithium.

25 ¹H-NMR (CDCl₃) δ: 9.65 - 9.45 (1H, br), 7.84 - 7.75 (2H, m), 7.46 - 7.35 (3H, m), 7.12 - 7.01 (3H, m), 3.85 (3H, s), 2.46 (3H, s).

STEP 3. Diethyl α-acetoxy-[5-methoxy-2-(3-methylbenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 5-methoxy-2-(3-methylbenzoyl)indole (step 2).

30 tlc: R_f=0.45 (ethyl acetate/hexane=1:3).

STEP 4. Diethyl [5-methoxy-2-(3-methylbenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetoxy-[5-methoxy-2-(3-methylbenzoyl)-1H-indol-3-yl]malonate (step 3).

- 5 $^1\text{H-NMR}(\text{CDCl}_3)\delta$: 9.52 - 9.13 (0.5H, m), 8.81 - 8.71 (0.5H, m), 7.80 - 7.72 (1H, m), 7.63 - 7.55 (1H, m), 7.47 - 7.00 (5H, m), 5.41(0.5H, s), 5.37 (0.5H, s), 4.27 - 4.15 (4H, m), 3.89 (1.5H, s), 3.84 (1.5H, s), 2.45 (1.5H, s), 2.43 (1.5H, s), 1.29 - 1.19 (6H, m).

STEP 5. [5-Methoxy-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid

- 10 The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [5-methoxy-2-(3-methylbenzoyl)-1H-indol-3-yl]malonate (step 4).

m.p.:230.4-232.0 °C (decomposed) (recrystallized from ethyl acetate).

IR (KBr) ν : 3310, 1705, 1614, 1583 cm^{-1} .

- 15 $^1\text{H-NMR}(\text{DMSO}-d_6)\delta$: 11.47 (1H, br s), 7.60-7.34 (5H, m), 7.16-7.09 (1H, m), 6.98 (1H, dd, $J=9.0, 2.4\text{Hz}$), 3.78 (5H, s), 2.40 (3H, s).

$^{13}\text{C-NMR}(\text{DMSO}-d_6)\delta$: 188.1, 172.1, 153.8, 138.9, 137.9, 132.7, 132.3, 132.0, 129.2, 128.5, 128.0, 125.8, 117.0, 115.6, 113.6, 100.7, 55.3, 30.6, 20.8.

EXAMPLE 24(2-BENZOYL-7-CHLORO-1H-INDOL-3-YL)ACETIC ACID

- 20 STEP 1. 7-Chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole

The title compound was prepared according to the procedure described in step 1 of Example 7 from 7-chloroindole-2-carboxylic acid (H. N. Rydon and J. C. Tweddle, *J. Chem. Soc.*, 1955, 3499).

- 25 $^1\text{H-NMR}(\text{CDCl}_3)\delta$: 9.40 (1H, br s), 7.59 (1H, d, $J=8.1\text{ Hz}$), 7.32-7.25 (2H, m), 7.08 (1H, t, $J=8.1\text{ Hz}$), 3.85 (3H, s), 3.43 (3H, s).

STEP 2. 2-Benzoyl-7-chloroindole

The title compound was prepared according to the procedure described in step 2 of Example 17 from 7-chloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (step 1) and phenyllithium.

- 30 $^1\text{H-NMR}(\text{CDCl}_3)\delta$: 9.40 (1H, br s), 8.01-7.96 (2H, m), 7.70-7.50 (4H, m), 7.38 (1H, d, $J=7.6\text{ Hz}$), 7.18 (1H, d, $J=2.0\text{ Hz}$), 7.11 (1H, t, $J=7.6\text{ Hz}$).

STEP 3. Diethyl α -acetoxy-(2-benzoyl-7-chloro-1H-indol-3-yl)malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 2-benzoyl-7-chloroindole (step 2).

¹H-NMR (CDCl₃) δ : 8.76 (1H, br s), 7.92-7.81 (3H, m), 7.68-7.60 (1H, m), 7.53-7.45 (2H, m), 7.33 (1H, d, J=7.6 Hz), 7.15 (1H, t, J=8.2 Hz), 4.60-4.20 (4H, m), 1.71 (3H, s), 1.98 (6H, t, J=7.1 Hz).

STEP 4. Diethyl (2-benzoyl-7-chloro-1H-indol-3-yl)malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetoxy-(2-benzoyl-7-chloro-1H-indol-3-yl)malonate (step 3).

¹H-NMR (CDCl₃) δ : 8.94 (1H, br s), 7.86-7.80 (2H, m), 7.75 (1H, d, J=8.1 Hz), 7.70-7.62 (1H, m), 7.55-7.50 (2H, m), 7.36 (1H, d, J=7.6 Hz), 7.13 (1H, t, J=7.6 Hz), 5.29 (1H, s), 4.25-4.11 (4H, m), 1.22 (6H, t, J=7.3 Hz).

STEP 5. (2-Benzoyl-7-chloro-1H-indol-3-yl)acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl (2-benzoyl-7-chloro-1H-indol-3-yl)malonate (step 4).

m.p.: 190-193 °C (recrystallized from ethyl acetate/hexane)

IR (KBr) ν : 1691, 1627, 1598, 1323, 1260 1199, 1010 cm⁻¹.

¹H-NMR (DMSO-d₆) δ : 11.90 (1H, br s), 7.82 (2H, m), 7.74-7.65 (2H, m), 7.62-7.38 (2H, m), 7.40 (1H, d, J=7.6 Hz), 7.13 (1H, t, J=7.6 Hz), 3.75 (2H, s).

EXAMPLE 25(2-BENZOYL-4,5-DICHLORO-1H-INDOL-3-YL)ACETIC ACIDSTEP 1. 4,5-dichloroindolyl-2-carboxylic acid

To a suspension of ethyl 4,5-dichloroindole-2-carboxylate (Ishii et al., *Chem. Pharm. Bull.*, 1974, 22, 1981, 1.8 g, 7.0 mmol) in ethanol (40 ml) was added 2N aqueous NaOH (10 ml) and the mixture was heated at reflux temperature for 2 h. The mixture was cooled to room temperature and concentrated. The residual solid was acidified with 2N aqueous HCl (30 ml) and extracted with diethyl ether (80 ml x 2). The organic extracts were dried (MgSO₄) and concentrated to give 1.5 g (94 %) of the

title compound as yellow solids.

¹H-NMR (DMSO-d₆) δ: 12.34 (1H, br s), 7.47-7.39 (2H, m), 7.08 (1H, d, J=1.8 Hz).

STEP 2. 4,5-Dichloro-2-[N-methoxy-N-methylamino)carbonyl]indole

The title compound was prepared according to the procedure described in step 1
5 of Example 7 from 4,5-dichloroindole-2-carboxylic acid (step 1).

¹H-NMR (CDCl₃) δ: 9.62 (1H, br s), 7.40-7.05 (3H, m), 3.88 (3H, s), 3.45 (3H, s).

STEP 3. 2-Benzoyl-4,5-dichloroindole

The titled compound was prepared according to the procedure described in step
2 of Example 17 from 4,5-dichloro-2-[(N-methoxy-N-methylamino)carbonyl]indole
10 (step 2) and phenyllithium.

m.p.: 206-210 °C (recrystallized from ethyl acetate/hexane).

¹H-NMR (CDCl₃) δ: 9.58 (1H, br s), 8.05-7.98 (2H, m), 7.71-7.53 (3H, m), 7.42 (1H, d,
J=8.9 Hz), 7.35 (1H, d, J=8.9 Hz), 7.22 (1H, s).

STEP 4. Diethyl α-acetoxy-(2-benzoyl-4,5-dichloro-1H-indol-3-yl)malonate

The title compound was prepared according to the procedure described in step 4
15 of Example 2 (Method B) from 2-benzoyl-4,5-dichloroindole (step 3).

¹H-NMR (CDCl₃) δ: 8.85 (1H, br s), 7.91-7.86 (2H, m), 7.66-7.58 (1H, m), 7.52-7.44
(2H, m), 7.36 (1H, d, J=8.7 Hz), 7.21 (1H, d, J=8.7 Hz), 4.20-3.98 (4H, m), 2.07 (3H,
s), 1.15 (6H, t, J=7.3 Hz).

20 STEP 5. Diethyl (2-benzoyl-4,5-dichloro-1H-indol-3-yl)malonate

The title compound was prepared according to the procedure described in step 5
of Example 2 (Method B) from diethyl α-acetoxy-(2-benzoyl-4,5-dichloro-1H-indol-3-
yl)malonate (step 4).

¹H-NMR (CDCl₃) δ: 9.24 (1H, br s), 7.82-7.76 (2H, m), 7.65-7.58 (1H, m), 7.55-7.44
25 (2H, m), 7.20 (1H, d, J=8.7 Hz), 7.03 (1H, d, J=8.7 Hz), 5.89 (1H, s), 4.16-4.02 (4H,
m), 1.20 (6H, t, J=7.3 Hz).

STEP 6. (2-Benzoyl-4,5-dichloro-1H-indol-3-yl)acetic acid

The title compound was prepared according to the procedure described in step 6
of Example 2 (Method B) from diethyl (2-benzoyl-4,5-dichloro-1H-indol-3-
30 yl)malonate (step 5).

m.p.: 249-252 °C (recrystallized from ethyl acetate/hexane).

IR (KBr) ν : 1701, 1625, 1523, 1450, 1330, 1257, 1012 cm^{-1} .

$^1\text{H-NMR}$ (CDCl_3) δ : 9.92 (1H, br s), 7.85-7.81 (2H, m), 7.66-7.42 (3H, m), 7.36 (1H, d, $J=8.8$ Hz), 7.29 (1H, d, $J=8.8$ Hz), 4.09 (2H, s).

5 **EXAMPLE 26**

(2-BENZOYL-4,6-DICHLORO-1H-INDOL-3-YL)ACETIC ACID

STEP 1. 4,6-Dichloro-2-(N-methoxy-N-methylamino)carbonyl]indole

The title compound was prepared according to the procedure described in step 1 of Example 7 from 4,6-dichloroindole-2-carboxylic acid (Salituro, Francesco G. et al.,
10 *J. Med. Chem.*, 1990, **33**, 2944).

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.09 (1H, br s), 7.48 (1H, dd, $J=1.0, 1.6$ Hz), 7.25 (1H, d, $J=1.6$ Hz), 7.11 (1H, s), 3.82 (3H, s), 3.63 (3H, s).

STEP 2. 2-Benzoyl-4,6-dichloroindole

The titled compound was prepared according to the procedure described in step
15 2 of Example 17 from 4,6-dichloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (step 1) and phenyllithium.

m.p.: 214-218 °C.

$^1\text{H-NMR}$ (CDCl_3) δ : 9.45 (1H, br s), 8.02-7.95 (2H, m), 7.70-7.52 (3H, m), 7.40 (1H, d, $J=1.6$ Hz), 7.25 (1H, s), 7.20 (1H, d, $J=1.6$ Hz).

20 **STEP 3. Diethyl α -acetoxy-(2-benzoyl-4,6-dichloro-1H-indol-3-yl)malonate**

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 2-benzoyl-4,6-dichloroindole (step 2).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.92 (1H, br s), 7.95-7.86 (2H, m), 7.68-7.43 (3H, m), 7.29 (1H, d, $J=1.8$ Hz), 7.23 (1H, d, $J=1.8$ Hz), 4.60-4.00 (4H, m), 2.04 (3H, s), 1.30-1.10 (6H, m)

25 **STEP 4. Diethyl (2-benzoyl-4,6-dichloro-1H-indol-3-yl)malonate**

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetoxy-(2-benzoyl-4,6-dichloro-1H-indol-3-yl)malonate (step 3).

m.p.: 170-172 °C.

¹H-NMR (CDCl₃) δ: 9.15 (1H, br s), 7.84-7.75 (2H, m), 7.66-7.45 (3H, m), 7.26 (1H, s), 7.12 (1H, s), 5.79 (1H, s), 4.20-4.00 (4H, m), 1.20 (6H, t, J=7.3 Hz).

STEP 5. (2-Benzoyl-4,6-dichloro-1H-indol-3-yl)acetic acid

The title compound was prepared according to the procedure described in step 6
5 of Example 2 (Method B) from diethyl [2-benzoyl-4,5-dichloro-1H-indol-3-yl]malonate (step 4).

m.p.: 239-243 °C (recrystallized from ethyl acetate/hexane).

IR (KBr) ν: 1725, 1555, 1525, 1330, 1287, 1250, 1005 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.40 (1H, br s), 12.12 (1H, br s), 7.80-7.67 (3H, m), 7.64-
10 7.54 (2H, m), 7.48 (1H, d, J=1.8 Hz), 7.23 (1H, d, J=1.8 Hz), 3.99 (2H, s).

EXAMPLE 27

(2-BENZOYL-5,6-DICHLORO-1H-INDOL-3-YL)ACETIC ACID

STEP 1. 5,6-dichloroindole-2-carboxylic acid

The title compound was prepared according to the procedure described in step 1
15 of Example 25 from ethyl 5,6-dichloroindole-2-carboxylate (Ishii et al., *Chem. Pharm. Bull.*, 1974, 22, 1981).

¹H-NMR (DMSO-d₆) δ: 12.06 (1H, br s), 7.95 (1H, s), 7.62 (1H, s), 7.09 (1H, s).

STEP 2. 5,6-Dichloro-2-[N-methoxy-N-methylamino)carbonyl]indole

The title compound was prepared according to the procedure described in step 1
20 of Example 7 from 5,6-dichloroindole-2-carboxylic acid (step 1).

¹H-NMR (CDCl₃) δ: 11.86 (1H, br s), 7.96 (1H, s), 7.66 (1H, s), 7.16 (1H, s), 3.80 (3H, s), 3.34 (3H, s).

STEP 3. 2-Benzoyl-5,6-dichloroindole

The titled compound was prepared according to the procedure described in step
25 2 of Example 17 from 5,6-dichloro-2-[(N-methoxy-N-methylamino)carbonyl]indole (step 2) and phenyllithium.

m.p.: 206-210 °C (recrystallized from ethyl acetate/hexane).

¹H-NMR (CDCl₃) δ: 9.35 (1H, br s), 8.00-7.93 (2H, m), 7.81 (1H, s), 7.69-7.51 (3H, m), 7.26 (1H, s), 7.08 (1H, s).

30 STEP 4. Diethyl α-acetoxy-(2-benzoyl-5,6-dichloro-1H-indol-3-yl)malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 2-benzoyl-5,6-dichloroindole (step 3).

¹H-NMR (CDCl₃) δ: 8.76 (1H, br s), 8.05 (1H, s), 7.90-7.82 (2H, m), 7.67-7.59 (1H, m), 7.54-7.44 (3H, m), 4.30-4.10 (4H, m), 1.69 (3H, s), 1.35-1.20 (6H, m).

5 STEP 5. Diethyl (2-benzoyl-5,6-dichloro-1H-indol-3-yl)malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α-acetoxy-(2-benzoyl-5,6-dichloro-1H-indol-3-yl)malonate (step 4).

¹H-NMR (CDCl₃) δ: 8.90 (1H, br s), 7.95 (1H, s), 7.82-7.76 (2H, m), 7.70-7.61 (1H, m), 7.58-7.46 (3H, m), 5.26 (1H, s), 4.30-4.05 (4H, m), 1.24 (6H, t, J=7.1 Hz).

10 STEP 6. (2-Benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl (2-benzoyl-4,5-dichloro-1H-indol-3-yl)malonate (step 5).

15 m.p.: 208-210 °C (recrystallized from ethyl acetate/hexane).

¹H-NMR (DMSO-d₆) δ: 11.91 (1H, br s), 8.04 (1H, s), 7.77-7.50 (6H, m), 3.81 (2H, s).

EXAMPLE 28

dl-2-(2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)PROPANOIC ACID

STEP 1. Diethyl α-methyl-(2-benzoyl-6-chloro-1H-indol-3-yl)malonate

20 The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 2-benzoyl-6-chloroindole (step 3 of Example 2; Method B) and diethyl methylmalonate.

m.p.: 193-196 °C.

¹H-NMR (CDCl₃) δ: 8.42 (1H, br s), 7.87 (2H, m), 7.66-7.48 (3H, m), 7.42 (1H, d, J=8.8 Hz), 7.35 (1H, d, J=1.8 Hz), 7.11 (1H, dd, J=1.8, 8.8 Hz), 4.30-4.02 (4H, m), 1.98 (3H, s), 1.15 (6H, t, J=7.1 Hz).

STEP 2. dl-2-(2-Benzoyl-6-chloro-1H-indol-3-yl)propanoic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl α-methyl-(2-benzoyl-6-chloro-1H-indol-3-yl)malonate (step 1).

30

m.p.: 204-208 °C (recrystallized from dichloromethane/hexane).

IR (KBr) ν : 1720, 1620, 1475, 1260, 1230 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.29 (1H, br s), 11.72 (1H, br s), 7.82-7.76 (2H, m), 7.75-7.68 (2H, m), 7.65-7.57 (2H, m), 7.48 (1H, d, $J=1.8$ Hz), 7.12 (1H, dd, $J=1.8, 8.4$ Hz),
5 4.15 (1H, q, $J=7.1$ Hz), 1.44 (3H, d, $J=7.1$ Hz).

EXAMPLE 29 and EXAMPLE 30

less polar antipode, 2-(2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)PROPANOIC ACID (EXAMPLE 29) and more polar antipode, 2-(2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)PROPANOIC ACID (EXAMPLE 30)

10 Chiral separation of *dl*-2-(2-Benzoyl-6-chloro-1H-indol-3-yl)propanoic acid (Example 28) was performed by DAICEL CHIRALCEL OJ (4.6 x 250 mm, eluent: hexane/isopropanol/trifluoroacetic acid = 85:15:0.1, flow rate: 1.0 ml/min) to afford; less polar compound (retention time: 17 min.) and more polar compound (retention time: 27 min.).

15 **EXAMPLE 31**

[6-CHLORO-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

STEP 1. Methyl *trans*-4-chloro-2-(ethoxycarbonylamino)cinnamate

To a stirred solution of methyl *trans*-2-amino-4-chlorocinnamate (R.W.Carling
20 et al., *J.Med.Chem.*, **1993**, 36, 3397., 32.6 g, 0.15 mol), pyridine (14.9 ml, 0.17 mol) and 4-dimethylaminopyridine (0.5 g) in dichloromethane (500 ml) was added dropwise ethyl chloroformate (16.2 ml, 0.17 mol) at room temperature. After stirring for 2h, the mixture was concentrated. The residue was diluted in ethyl acetate (1000ml) and washed with 10% aqueous citric acid (500ml). The organic layer was successively
25 washed with water (500 ml), saturated sodium bicarbonate (500 ml), water (500 ml), brine (500 ml). After drying (MgSO_4) and removal of solvent, the residual solids were recrystallized from ethyl acetate/hexane to give 39.13 g (92 %) of the title compound.

$^1\text{H-NMR}$ (CDCl_3) δ : 7.95 (1H, br s), 7.76 (1H, d, $J=15.8\text{Hz}$), 7.42 (1H, d, $J=8.6\text{Hz}$),
30 7.22 (1H, dd, $J=2.1, 8.5\text{Hz}$), 6.69 (1H, br), 6.37 (1H, d, $J=15.7\text{Hz}$), 4.26 (2H, q, $J=7.3\text{Hz}$), 3.82 (3H, s), 1.34 (3H, t, $J=7.25\text{Hz}$).

STEP 2. Methyl [6-chloro-1-ethoxycarbonyl-2-(4-methylpyridine-2-carbonyl)indolin-3-yl]acetate

A mixture of methyl *trans*-4-chloro-2-(ethoxycarbonylamino)cinnamate (step 1, 1.5 g, 5.3 mmol), 2-bromoacetyl-4-methylpyridine hydrobromide*, potassium carbonate (7.3 g, 53 mmol) and acetonitrile (50 ml) was heated at reflux temperature for 17h. The mixture was then cooled and concentrated. The residue was diluted in ethyl acetate (200 ml) and washed with water (200 ml) and brine (200 ml). After drying (MgSO₄) and removal of solvent, the crude product was purified by flash column chromatography eluting with ethyl acetate/hexane (1:5) to afford 433 mg (20 %) of the title compound.

MS (EI) m/z: 416 (M⁺).

* 2-bromoacetyl-4-methylpyridine hydrobromide was prepared as follows;

To a stirred solution of 2-acetyl-4-methylpyridine (F.H.Case et al, *J. Am. Chem. Soc.*, 1956, 78, 5842., 7.8 g, 57.7 mmol) in 25% HBr-AcOH (40 ml) was added dropwise a solution of bromine (10.1 g, 63.5 mmol) in AcOH (10 ml) with ice-cooling. After stirring for 1h, diethyl ether (100 ml) was added and the precipitates were collected by filtration to give 10.8 g (63 %) of the title compound.

¹H-NMR (DMSO-d₆) δ: 8.71 (1H, d, J=5.1Hz), 8.14 (1H, s), 7.75 (1H, d, J=5.1Hz), 5.07 (2H, s), 2.52 (3H, s).

STEP 3. [6-Chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid

A stirred solution of methyl [6-chloro-1-ethoxycarbonyl-2-(4-methylpyridine-2-carbonyl)indolin-3-yl]acetate (step 2, 930 mg, 2.2 mmol) in ethanol (20 ml) and 2N aqueous NaOH (10 ml) was heated at reflux temperature for 72h. After cooling to room temperature, the resulting mixture was neutralized with 2N aqueous HCl (10 ml) and concentrated. The residue was diluted in dichloromethane/methanol (10:1, 300 ml) and dried (MgSO₄). After removal of solvent, the crude product was purified by flash column chromatography eluting with dichloromethane/methanol (20:1) and then washed with *i*-PrOH (ca.20 ml) to give 120 mg (17 %) of the title compound as a yellow powder.

m.p.: 223 °C (decomposed).

IR (KBr) ν: 1707, 1647, 1595, 1533, 1487, 1429, 1276, 1289, 1196 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.30 (1H, s), 8.70 (1H, d, J=4.9Hz), 7.96 (1H, br s), 7.85-7.70 (2H, m), 7.65-7.55 (1H, m), 7.17-7.07 (1H, m), 4.08 (2H, s), 2.47 (3H, s).

EXAMPLE 32

[6-CHLORO-2-(5-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC

5 ACID

STEP 1. Methyl [6-chloro-1-ethoxycarbonyl-2-(5-methylpyridine-2-carbonyl)indolin-3-yl]acetate

Two diastereomers of the title compound were prepared according the procedure described in step 2 of Example 31 from methyl *trans*-4-chloro-2-
10 (ethoxycarbonylamino)cinnamate (Example 31, step 1) and 2-bromoacetyl-5-methylpyridine.*

Less polar product; tlc: R_f=0.30 (ethyl acetate/hexane=1/2), MS (EI) m/z: 416 (M⁺).

More polar product; tlc: R_f=0.25 (ethyl acetate/hexane=1/2), MS (EI) m/z: 416 (M⁺).

* 2-Bromoacetyl-5-methylpyridine was prepared as follows;

15 A mixture of 2-bromo-5-methylpyridine (5.00 g, 29.06 mmol), tributyl(1-ethoxyvinyl)tin (10.49 g, 29.07 mmol), and tetrakis(triphenylphosphine)palladium (3.36 g, 2.91 mmol) in toluene (40 ml) was heated at reflux temperature for 18 h. The mixture was cooled, filtered through a pad of Celite and then concentrated. The residue (~10 g) was dissolved in a mixture of THF (100 ml) and water (20 ml), cooled
20 to 0 °C and *N*-bromosuccinimide (5.43 g, 30.52 mmol) was added over 20 min. The resulting mixture was stirred for 0.5 h at the same temperature and then concentrated to ca. 20 ml. The mixture was diluted in ethyl acetate (300 ml), washed with water (100 ml x 3), and dried (MgSO₄). After removal of solvent, the crude product was purified by flash column chromatography eluting with ethyl acetate/hexane (1:15 to
25 1:10) to afford 2.30 g (37%) of the title compound as an oil.

¹H-NMR (CDCl₃) δ: 8.49 (1 H, br s), 8.01 (1 H, d, J=8.1 Hz), 7.68-7.64 (1 H, m), 4.84 (2 H, s), 2.44 (3 H, s).

STEP 2. [6-Chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid

Both of the diastereoisomers of methyl [6-chloro-1-ethoxycarbonyl-2-(5-
30 methylpyridine-2-carbonyl)indolin-3-yl]acetate (step 2) were converted to the title compound, respectively, according to the procedure described in step 3 of Example 31.

MS (EI) m/z: 328 (M^+)

m.p.: 235-238 °C (recrystallized from ethyl acetate).

IR (KBr) ν : 3281, 1699, 1638, 1529, 1310, 1150, 797, 702 cm^{-1} .

$^1\text{H-NMR}$ (DMSO- d_6) δ : 12.26 (1 H, br s), 8.68 (1 H, br s), 8.04 (1 H, d, $J=8.1$ Hz),
5 7.94 (1 H, br d, $J=9.1$ Hz), 7.79 (1 H, d, $J=8.7$ Hz), 7.74 (1 H, br s), 7.11 (1 H, br d, $J=8.6$ Hz), 4.10 (2 H, s), 2.47 (3 H, s). One signal due to NH or COOH was not observed.

EXAMPLE 33

10 METHYL [6-CHLORO-2-(4-CHLOROPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

A mixture of methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A, 675 mg, 1.92 mmol), 2-bromoacetyl-4-chloropyridine hydrobromide* (907 mg, 2.88 mmol), and potassium carbonate (2.65 g, 19.18 mmol) in acetone (20 ml) was heated at reflux temperature for 4 h. The mixture was cooled
15 and concentrated. The residue was diluted with ethyl acetate (150 ml) and washed with water (70 ml x 6). After drying (MgSO_4) and removal of solvent, the crude product was purified by flash column chromatography eluting with ethyl acetate/hexane (1:6-1:3) to afford 195 mg (28%) of the title compound (yellow solids) along with 264 mg (27%) of methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1-
20 (phenylsulfonyl)indolin-3-yl]acetate (brown crystals).

Methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate:

MS (EI) m/z: 362 (M^+).

$^1\text{H-NMR}$ (CDCl_3) δ : 12.09 (1 H, br s), 8.62 (1 H, d, $J=5.3$ Hz), 8.28 (1 H, d, $J=2.1$ Hz),
7.57 (1 H, d, $J=8.6$ Hz), 7.52 (1 H, dd, $J=2.0$ and 5.3 Hz), 7.45 (1 H, d, $J=1.8$ Hz), 7.08
25 (1 H, dd, $J=1.8$ and 8.7 Hz), 4.27 (2 H, s), 3.75 (3 H, s).

Methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1-(phenylsulfonyl)indolin-3-yl]acetate:

tlc: $R_f=0.35$ (ethyl acetate/hexane=1:2).

MS (EI) m/z (intensity): 504 (M^+ , 0.1), 363 (90), 335 (30), 304 (100), 275 (10), 223
30 (15).

* 2-Bromoacetyl-4-chloropyridine hydrobromide was prepared as follows;

4-Chloro-2-pyridinecarbonitrile: To a mixture of 4-chloropyridine-*N*-oxide (5.00 g, 38.6 mmol) and trimethylsilyl cyanide (4.84 g, 46.3 mmol) in dichloromethane (60 ml) cooled to 0 °C was added dropwise *N,N*-dimethylcarbamoyl chloride (3.8 ml, 40.5 mmol). The mixture was allowed to warm to ambient temperature and stirred for 16 h. The mixture was cooled to 0 °C and a 30% aqueous solution of potassium carbonate (100 ml) was added. The crude product was extracted with dichloromethane (100 ml x 2), the organic extracts dried (MgSO₄) and evaporated to give 4-chloro-2-pyridinecarbonitrile (5.35 g, 100%).

¹H-NMR (CDCl₃) δ: 8.63 (1 H, d, J=4.8 Hz), 7.72 (1 H, d, J=2.6 Hz), 7.55 (1 H, dd, J=1.8, 5.1 Hz).

2-Acetyl-4-chloropyridine: To a solution of 4-chloro-2-pyridinecarbonitrile (5.35 g, 38.6 mmol) in benzene (50 ml) and ether (50 ml) cooled to 0 °C was added dropwise over 20 min a 2M solution of MeMgI in ether (23 ml, 46.3 mmol). After 0.5 h, the mixture was allowed to warm to ambient temperature, and stirring continued for 2 h.

The mixture was cooled to 0 °C and 2M aqueous HCl (100 ml) added. The mixture was made basic with saturated aqueous sodium bicarbonate (~80 ml) and the organic layer separated and dried (MgSO₄). After removal of solvent, the residue was purified by flash chromatography eluting with ethyl acetate/hexane (1:5) to afford 3.60 g (60%) of 2-acetyl-4-chloropyridine.

¹H-NMR (DMSO-d₆) δ: 8.59 (1 H, d, J=5.1 Hz), 8.04 (1 H, d, J=1.8 Hz), 7.47 (1 H, dd, J=1.8, 5.1 Hz), 2.72 (3 H, s).

2-Bromoacetyl-4-chloropyridine hydrobromide: 2-(Bromoacetyl)-4-chloropyridine hydrobromide was prepared from 2-acetyl-4-chloropyridine according to the method of H. McKennis, Jr., L. B. Turnbull, E. R. Bowman, and E. Tamaki (in *J. Org. Chem.*, 1963, 28, 383-387).

¹H-NMR (DMSO-d₆) δ: 8.74 (1 H, d, J=5.5 Hz), 8.05 (1 H, d, J=1.8 Hz), 7.88 (1 H, dd, J=2.2 and 5.5 Hz), 5.02 (2 H, s).

EXAMPLE 34

[6-CHLORO-2-(4-CHLOROPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC

ACID

A suspension of methyl [6-chloro-2-[4-chloropyridine-2-carbonyl]-1H-indol-3-

yl]acetate (Example 33, 195 mg, 0.537 mmol) in ethanol (20 ml) and 2N sodium hydroxide (4 ml) was heated for 1 h at 50 °C. After cooling to room temperature, 2N hydrochloric acid (4 ml) was added and the mixture was concentrated. The residue was diluted in ethyl acetate (100 ml), washed with water (50 ml x 2), and dried
5 (MgSO₄). After removal of solvent, the crystalline residue was recrystallized from ethyl acetate to afford 175 mg (94%) of the title compound.

m.p.: 233-234 °C.

IR (KBr) ν : 3306, 1709, 1641, 1531, 1254, 1236, 741 cm⁻¹.

¹H-NMR (DMSO-d₆) δ : 12.20 (1 H, br s), 12.16 (1 H, br s), 8.80 (1 H, d, J=5.4 Hz),
10 8.12 (1 H, d, J=2.1 Hz), 7.90 (1 H, dd, J=2.1, 5.3 Hz), 7.81 (1 H, d, J=8.7 Hz), 7.70 (1 H, d, J=1.8 Hz), 7.13 (1 H, dd, J=2.0, 8.7 Hz), 4.07 (2 H, s).

EXAMPLE 35

[6-CHLORO-2-(PYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

STEP 1. Methyl [6-chloro-1-ethoxycarbonyl-2-(pyridine-2-carbonyl)indolin-3-
15 yl]acetate

The title compound were prepared according the procedure described in step 2 of Example 31 from methyl *trans*-4-chloro-2-(ethoxycarbonylamino)cinnamate (Example 31, step 1) and 2-bromoacetylpyridine hydrobromide (H.McKennis et al., *J.Org.Chem.*, 1963, 387.).

20 tlc: R_f=0.3 (ethyl acetate/hexane=1:3).

STEP 2. [6-Chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid

The title compound were prepared according the procedure described in step 3 of Example 31 from methyl [6-chloro-1-ethoxycarbonyl-2-(pyridine-2-carbonyl)indolin-3-yl]acetate (step 1).

25 m.p.: 210 °C (decomposed).

IR (KBr) ν : 3280, 1697, 1643, 1531, 1234, 1150 cm⁻¹.

¹H-NMR (DMSO-d₆) δ : 12.22 (1H, s), 8.84 (1H, d, J=4.9Hz), 8.15-8.05 (2H, m), 7.85-7.65 (3H, m), 7.11 (1H, dd, J=1.9, 8.7Hz), 4.04 (2H, s).

EXAMPLE 36

30 [5-CHLORO-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

STEP 1. Methyl *trans*-5-chloro-2-nitrocinnamate

A mixture of the 5-chloro-2-nitrobenzaldehyde (9.68 g, 52.16 mmol) and methyl (triphenylphosphoranylidene)acetate (18.31 g, 54.77 mmol) in toluene (200 ml) was heated at reflux temperature for 2 h. The mixture was concentrated and the crystalline residue was purified by flash column chromatography eluting with ethyl acetate/hexane (1:5) to afford crystals. Recrystallization from ethyl acetate/hexane gave 7.54 g (60%) of the title compound as pale yellow solids.

¹H-NMR (CDCl₃) δ: 8.09 (1 H, d, J=15.8 Hz), 8.04 (1 H, d, J=8.7 Hz), 7.60 (1 H, d, J=2.1 Hz), 7.51 (1 H, dd, J=2.1, 8.7 Hz), 6.36 (1 H, d, J=15.8 Hz), 3.84 (3 H, s).

10 STEP 2. Methyl *trans*-2-amino-5-chlorocinnamate

A mixture of methyl *trans*-5-chloro-2-nitrocinnamate (step 1, 3.00 g, 12.42 mmol), iron powder (3.65 g, 62.08 mmol), ammonium chloride (332 mg, 6.21 mmol), ethanol (60 ml) and water (10 ml) was heated at reflux temperature for 2 h. The mixture was cooled and filtered through a pad of Celite. The filtrate was concentrated. The residue was diluted with ethyl acetate (200 ml) and washed with water (100 ml x 2). After drying (MgSO₄), removal of solvent gave 2.57 g (98%) of the title compound as crystals.

¹H-NMR (CDCl₃) δ: 7.73 (1 H, d, J=15.8 Hz), 7.34 (1 H, d, J=2.5 Hz), 7.12 (1 H, dd, J=2.3, 8.6 Hz), 6.64 (1 H, d, J=8.6 Hz), 6.35 (1 H, d, J=15.8 Hz), 3.95 (2 H, br s), 3.81 (3 H, s).

20 STEP 3. Methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate

The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-2-amino-5-chlorocinnamate (step 2).

¹H-NMR (CDCl₃) δ: 7.72-7.67 (2 H, m), 7.58-7.51 (1 H, m), 7.47-7.40 (4 H, m), 7.36 (1 H, d, J=8.6 Hz), 7.31 (1 H, dd, J=2.1, 8.6 Hz), 6.14 (1 H, d, J=15.8 Hz), 3.78 (3 H, s). One signal due to NH was not observed.

STEP 4. Methyl [5-Chloro-2-(4-methylpyridine-2-carbonyl)-1-(phenylsulfonyl)indolin-3-yl]acetate

The title compound was prepared according to the procedure described in step 2 of Example 8 (Method A) from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (step 3) and 2-bromoacetyl-4-methylpyridine

hydrobromide (F.H.Case et al., *J.Am.Chem.Soc.*, **1956**, 78, 5842).

tlc: Rf=0.32 (ethyl acetate/hexane=1:2)

STEP 5. [5-Chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 3
5 of Example 31 from methyl [5-chloro-2-(4-methylpyridine-2-carbonyl)-1-(phenylsulfonyl)indolin-3-yl]acetate (step 4).

MS (EI) m/z: 328 (M⁺).

m.p.: 230-231 °C (recrystallized from ethyl acetate).

IR (KBr) ν : 3292, 1699, 1597, 1533, 1282, 1198, 1059, 802, 704 cm⁻¹.

10 ¹H-NMR (DMSO-d₆) δ : 12.26 (1 H, br s), 8.69 (1 H, d, J=5.1 Hz), 7.93 (1 H, br s), 7.82 (1 H, d, J=2.0 Hz), 7.66 (1 H, d, J=8.7 Hz), 7.56 (1 H, br d, J=4.9 Hz), 7.31 (1 H, dd, J=2.0, 8.7 Hz), 4.02 (2 H, s), 2.46 (3 H, s). One signal due to NH or COOH was not observed.

EXAMPLE 37

15 METHYL [5-CHLORO-2-(6-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

STEP 1. Methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1-(phenylsulfonyl)indolin-3-yl]acetate

The title compound was prepared according to the procedure described in step 2
20 of Example 8 (Method A) from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetyl-6-methylpyridine hydrobromide (H.Erlenmeyer, J.Jenni, and B.Prijs, *J.Med.Pharm.Chem.*, **1961**, 3, 561-566).

tlc: Rf=0.39 (ethyl acetate/hexane=2:3).

25 STEP 2. Methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate

The title compound was prepared according to the procedure described in step 3 of Example 8 (Method A) from methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1-(phenylsulfonyl)indolin-3-yl]acetate (step 1).

30 ¹H-NMR (CDCl₃) δ : 12.62 (1 H, br s), 8.15 (1 H, d, J=7.9 Hz), 7.84 (1 H, t, J=7.7 Hz), 7.67 (1 H, d, J=1.8 Hz), 7.43 (1 H, d, J=8.9 Hz), 7.40 (1 H, d, J=7.7 Hz), 7.32 (1 H, dd, J=2.0, 8.7 Hz), 4.28 (2 H, s), 3.73 (3 H, s), 2.76 (3 H, s).

Example 38**[5-CHLORO-2-(6-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID**

The title compound was prepared according to the procedure described in
5 Example 34 from methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate.

MS (EI) m/z: 328 (M^+).

m.p.: 225-226°C (recrystallized from ethyl acetate).

IR (KBr) ν : 1705, 1636, 1529, 1333, 1236, 1180, 1061, 669 cm^{-1} .

10 $^1\text{H-NMR}$ (DMSO- d_6) δ : 12.13 (1 H, br s), 12.09 (1 H, br s), 7.99 (1 H, t, $J=7.7$ Hz), 7.87 (1 H, d, $J=7.7$ Hz), 7.84 (1 H, d, $J=2.0$ Hz), 7.66 (1 H, d, $J=8.9$ Hz), 7.59 (1 H, d, $J=7.6$ Hz), 7.33 (1 h, dd, $J=2.0, 8.9$ Hz), 4.03 (2 H, s), 2.69 (3 H, s).

EXAMPLE 39**[6-CHLORO-2-(1-METHYLIMIDAZOLE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID**
15**STEP 1. Methyl [6-chloro-1-ethoxycarbonyl-2-(1-methylimidazole-2-carbonyl)indolin-3-yl]acetate**

The title compound was prepared according to the procedure described in step 2
of Example 31 from methyl *trans*-4-chloro-2-(ethoxycarbonylamino)cinnamate
20 (Example 31, step 1) and 2-bromoacetyl-1-methylimidazole hydrobromide.*

MS (EI) m/z: 405 (M^+).

* 2-Bromoacetyl-1-methylimidazole hydrobromide was prepared as follows;

To a stirred suspension of 2-acetyl-1-methylimidazole (D.H.Davis, J.Hall, and
E.H.Smith, *J.Chem.Soc.Perkin trans. 1*, 1991, 2691., 3.0 g, 26.8 mmol) in 25% HBr-
25 AcOH was added dropwise a solution of bromine (4.7 g, 29.5 mmol) with ice-cooling.
After stirring for 0.5h, the mixture was allowed to warm to room temperature and the
stirring was continued for an additional 1h. To the mixture was added diethyl ether
(150 ml) and the mixture was cooled with ice-bath. The precipitates were collected
by filtration to give 5.2 g (66 %) of the title compound as a pale yellow powder.

30 $^1\text{H-NMR}$ (DMSO- d_6) δ : 7.61 (1H, s), 7.27 (1H, s), 4.68 (2H, s), 3.81 (3H, s).

STEP 2. [6-Chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 3 of Example 31 from [6-chloro-1-ethoxycarbonyl-2-(1-methylimidazole-2-carbonyl)indolin-3-yl]acetate (step 1).

m.p.: 236 °C (decomposed).

5 MS (EI) m/z: 317 (M^+).

IR (KBr) ν : 3238, 1695, 1630, 1537, 1402, 1229, 1146 cm^{-1} .

$^1\text{H-NMR}$ ($\text{CDCl}_3 + \text{DMSO-d}_6$) δ : 12.30 (1H, s), 7.65 (1H, d, $J=8.7\text{Hz}$), 7.50 (1H, d, $J=1.8\text{Hz}$), 7.42 (1H, s), 7.28-7.23 (1H, m), 7.16 (1H, s), 7.09 (1H, dd, $J=1.8, 8.6\text{Hz}$), 4.25 (2H, s), 4.13 (3H, s).

10 **EXAMPLE 40**

METHYL [5-CHLORO-2-(THIAZOLE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

STEP 1. Methyl [5-chloro-1-phenylsulfonyl-2-(thiazole-2-carbonyl)indolin-3-yl]acetate

The title compound was prepared according to the procedure described in step 2 of Example 8 (Method A) from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetylthiazole hydrobromide (A.Dondoni, A.Marra, and P.Merino, *J.Am.Chem.Soc.*, 1994, **116**, 3324).
15 tlc: $R_f=0.07$ (ethyl acetate/hexane=1:2).

STEP 2. Methyl [5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetate

20 The title compound was prepared according to the procedure described in step 3 of Example 8 (Method A) from methyl [5-chloro-1-phenylsulfonyl-2-(thiazole-2-carbonyl)indolin-3-yl]acetate (step 1).

$^1\text{H-NMR}$ (CDCl_3) δ : 11.78 (1 H, br s), 8.12 (1 H, d, $J=3.1\text{ Hz}$), 7.75 (1 H, d, $J=3.1\text{ Hz}$), 7.68 (1 H, d, $J=1.8\text{ Hz}$), 7.44 (1 H, d, $J=8.7\text{ Hz}$), 7.34 (1 H, dd, $J=2.0, 8.9\text{ Hz}$), 4.29 (2
25 H, s), 3.74 (3 H, s).

EXAMPLE 41

[5-CHLORO-2-(THIAZOLE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [5-chloro-1-phenylsulfonyl-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetate (step 2).
30

MS (EI) m/z: 320 (M^+).

m.p.; 230-231°C (recrystallized from ethyl acetate).

IR (KBr) ν : 3302, 1703, 1636, 1541, 1387, 1335, 1267, 1232, 1186, 1003, 766 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.23 (1 H, br s), 12.10 (1 H, br s), 8.33 (1 H, d, $J=3.1$ Hz), 8.31 (1 H, d, $J=3.1$ Hz), 7.89 (1 H, d, $J=2.0$ Hz), 7.77 (1 H, d, $J=8.9$ Hz), 7.36 (1 H, dd, $J=2.0, 8.7$ Hz), 4.17 (2 H, s).

EXAMPLE 42

METHYL (2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)ACETATE

A mixture of (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid (Example 2, 50 mg, 0.16 mmol) and 10% HCl in methanol (3 ml) was stirred for 3 h at room temperature.

The mixture was concentrated and the residue was purified by flash column chromatography eluting with ethyl acetate/hexane (1:5) to afford 23 mg (44 %) of the title compound as white solids.

m.p.: 134-137 °C.

IR (KBr) ν : 1735, 1620, 1529, 1434, 1325, 1147, 1013, 945 cm^{-1} .

$^1\text{H-NMR}$ (CDCl_3) δ : 8.96 (1H, br s), 7.85-7.74 (2H, m), 7.66-7.47 (4H, m), 7.38 (1H, d, $J=1.8$ Hz), 7.15 (1H, dd, $J=1.8, 8.6$ Hz), 3.18 (2H, s), 3.65 (3H, s).

EXAMPLE 43

(2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)-N,N-DIMETHYLACETAMIDE

To a solution of (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid (Example 2, 140 mg, 0.45 mmol), dimethylamine hydrochloride (45 mg, 0.54 mmol) and triethylamine (0.1 ml, 0.54 mmol) in DMF (2 ml) at 0 °C was added diethyl phosphorocyanidate (DEPC, 0.1 ml, 0.54 mmol). The mixture was then stirred at room temperature for 1 h. The mixture was poured into water (20 ml) and extracted with diethyl ether (50 ml x 2). The organic extracts were washed with water (30 ml x 2), dried (MgSO_4) and concentrated. The residual solids were recrystallized from ethyl acetate/hexane to afford 50 mg (33%) of the title compound.

m.p.: 190-193 °C (recrystallized from ethyl acetate/hexane).

IR (KBr) ν : 1631, 1232, 1007, 908 cm^{-1} .

$^1\text{H-NMR}$ (CDCl_3) δ : 11.72 (1H, br s), 7.72-7.64 (4H, m), 7.60-7.52 (2H, m), 7.45 (1H, d, $J=1.2$ Hz), 7.08 (1H, d, $J=8.6$ Hz), 3.32 (2H, s), 2.80 (3H, s), 2.76 (3H, s).

EXAMPLE 44**(2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)-N-METHYLACETAMIDE**

The title compound was prepared according to the procedure described in Example 43 from (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid (Example 2) and methylamine hydrochloride.

m.p.: 242-246 °C (recrystallized from ethyl acetate/hexane).

IR (KBr) ν : 1618, 1527, 1409, 1325 cm^{-1} .

$^1\text{H-NMR}$ (CDCl_3) δ : 11.71 (1H, br s), 7.82-7.75 (2H, m), 7.62-7.52 (5H, m), 7.49-7.46 (1H, m), 7.15-7.08 (1H, m), 3.64 (2H, s), 3.31 (3H, s).

10 EXAMPLE 45**(2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)ACETAMIDE**

The title compound was prepared according to the procedure described in Example 43 from (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid (Example 2) and a solution of ammonia in THF.

15 m.p.: 234-237 °C.

IR (KBr) ν : 1665, 1618, 1566, 1523, 1325, 1259, 943 cm^{-1} .

$^1\text{H-NMR}$ (CDCl_3) δ : 11.66 (1H, br s), 7.82-7.75 (2H, m), 7.74-7.65 (2H, m), 7.62-7.53 (2H, m), 7.46 (1H, d, $J=1.8$ Hz), 7.27 (1H, br s), 7.12 (1H, dd, $J=1.8, 8.6$ Hz), 6.85 (1H, br s), 3.63 (2H, s).

20 EXAMPLE 46**(2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)-N-METHOXY-N-METHYLACETAMIDE**

The title compound was prepared according to the procedure described in Example 43 from (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid (Example 2) and *N,O*-dimethylhydroxylamine hydrochloride.

25 dimethylhydroxylamine hydrochloride.

m.p.: 109.9 - 112.2 °C (decomposed).

IR (KBr) ν : 3179, 2970, 2937, 1634, 1599, 1570 cm^{-1} .

$^1\text{H-NMR}$ (CDCl_3) δ : 9.20 - 8.90 (1H, m), 7.84 - 7.75 (2H, m), 7.66 - 7.45 (4H, m), 7.33 (1H, d, $J=1.3$ Hz), 7.12 (1H, dd, $J=8.6, 1.8$ Hz), 3.94 (2H, s), 3.51 (3H, s), 3.13 (3H, s).

30 EXAMPLE 47

2-(2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)-1-PIPERIDINO-1-ETHANONESTEP 1. 7-Chloro-1-phenyl-9H-pyrano[3,4-b]indole-3-one

A solution of (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid (Example 2, 200 mg, 0.64 mmol), diethyl phosphorocyanidate (DEPC, 0.12 ml, 0.76 mmol) and triethylamine (0.11 ml, 0.76 mmol) in DMF (3.0 ml) was stirred at room temperature for 5 min. The mixture was then poured into water (20 ml) and the orange precipitates were collected by filtration to give 20 mg (11%) of the title compound as orange solids.

¹H-NMR (CDCl₃) δ: 10.79 (1H, br s), 8.10 (1H, d, J=8.4 Hz), 7.93-7.84 (2H, m), 7.67-7.50 (3H, m), 7.29 (1H, s), 7.13 (1H, d, J=8.4 Hz), 6.87 (1H, s).

STEP 2. 2-(2-Benzoyl-6-chloro-1H-indol-3-yl)-1-piperidino-1-ethanone

A mixture of 7-chloro-1-phenyl-9H-pyrano[3,4-b]indole-3-one (step 1, 0.30 g, 1.0 mmol), and piperidine (1.0 ml, 10 mmol) in methanol (20 ml) was heated under reflux temperature for 2 h. After cooling down to rt, the yellow mixture was concentrated and the residual solids were recrystallized from methanol/hexane to give 0.12 g (32 %) of the title compound.

m.p.: 223-224 °C.

IR (KBr) ν: 3310, 2928, 1655, 1570, 1533, 1447, 1323, 1256, 1225, 1059, 945, 858, 737, 700 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.7 (1H, br s), 7.74-7.63 (4H, m), 7.55 (2H, t, J=8.7 Hz), 7.46 (1H, d, J=1.8 Hz), 7.09 (1H, dd, J=8.7, 1.8 Hz), 3.80 (2H, s), 3.45-3.20 (4H, m), 1.55-1.20 (6H, m)

EXAMPLE 482-(2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)-N-(4-METHYL-1-PIPERAZINYL)-1-ETHANONE

The title compound was prepared according to the procedure described in Example 43 from (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid (Example 2) and 1-methylpiperazine.

m.p.: 184-1°5 °C (recrystallized from methanol/hexane).

IR (KBr) ν: 2939, 2795, 1634, 1531, 1435, 1325, 1229, 1144, 1001, 737, 700 cm⁻¹.

¹H-NMR (CDCl₃) δ: 9.37 (1H, br s), 7.78-7.75 (1H, m), 7.73 (1H, d, J=1.5 Hz), 7.62-

7.52 (2H, m), 7.52-7.43 (2H, m), 7.18-7.10 (1H, m), 7.10-7.02 (1H, m), 3.81 (2H, s), 3.54 (2H, br s), 3.36 (2H, t, J=4.8 Hz), 2.36-2.22 (4H, m), 2.28 (3H, s).

EXAMPLE 49

(2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)-N-(2-CYANOETHYL)ACETAMIDE

5 The title compound was prepared according to the procedure described in Example 43 from (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid (Example 2) and aminopropionitrile.

m.p.: 233-233.5 °C (recrystallized from methanol/hexane).

¹H-NMR (DMSO-d₆) δ: 11.7 (1H, br s), 8.16 (1H, t, J=6.3Hz), 7.81-7.75 (2H, m),
10 7.72-7.64 (2H, m), 7.62-7.53 (2H, m), 7.47 (1H, d, J=1.9 Hz), 7.10 (1H, dd, J=8.7, 1.9 Hz), 3.67 (2H, s), 3.23 (2H, q, J=6.3 Hz), 2.58 (2H, t, J=6.3 Hz)

EXAMPLE 50

(2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)-N-(2-HYDROXYETHYL)ACETAMIDE

15 The title compound was prepared according to the procedure described in Example 43 from (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid (Example 2) and 2-aminoethanol.

m.p.: 178-179.5 °C (recrystallized from methanol/hexane).

¹H-NMR (DMSO-d₆) δ: 11.7 (1H, br s), 7.84-7.75 (3H, m), 7.72-7.64 (2H, m), 7.61-
20 7.52 (2H, m), 7.46 (1H, d, J=2.0 Hz), 7.11 (1H, dd, J=8.6, 2.0 Hz), 3.64 (2H, s), 3.38-3.30 (3H, m), 3.06 (2H, q, J=5.9 Hz).

EXAMPLE 51

2-(2-BENZOYL-6-CHLORO-1H-INDOL-3-YL)-1-MORPHOLINO-1-ETHANONE

25 The title compound was prepared according to the procedure described in Example 43 from (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid (Example 2) and morpholine.

m.p.: 187.7-189.5 °C.

IR (KBr) ν: 3339, 2964, 2849, 1653, 1612, 1568 cm⁻¹.

¹H-NMR (CDCl₃) δ: 9.08 - 8.92 (1H, m), 7.81 - 7.72 (2H, m), 7.69 - 7.58 (2H, m), 7.56
30 - 7.47 (2H, m), 7.29 (1H, d, J=1.8Hz), 7.12 (1H, dd, J=8.6, 1.8Hz), 3.87 (2H, s), 3.67 - 3.46 (6H, m), 3.41 - 3.31 (2H, m).

EXAMPLE 52**[2-(4-CHLOROBENZOYL)-1H-INDOL-3-YL]ACETIC ACID****STEP 1. 2-(4-Chlorobenzoyl)-1-(phenylsulfonyl)indole**

To a solution of 1-(phenylsulfonyl)indole (500 mg, 1.94 mmol) in THF (5 ml)
5 was added dropwise *tert*-butyllithium (1.4 ml, 2.33 mmol) under nitrogen atmosphere
at -78 °C. The yellow solution was cannulated directly into a solution of *p*-
chlorobenzoyl chloride (0.3 ml, 2.33 mmol) in THF (3 ml) cooled to -78 °C. The
reaction mixture was stirred at -78 °C for 2 h. The mixture was quenched with
saturated ammonium chloride and extracted with ethyl acetate (100 ml). The organic
10 layer was washed with water (50 ml), brine (50 ml) and dried (MgSO₄). After
removal of solvent, the crude product was purified by flash column chromatography
eluting with hexane/ethyl acetate (10:1) to afford 339 mg (44.1 %) of the title
compound as yellow amorphous solids.

¹H-NMR (CDCl₃) δ: 8.13 (1H, d, J=8.4Hz), 8.04-8.00 (2H, m), 7.93-7.90 (2H, m),
15 7.58-7.44 (7H, m), 7.30 (1H, t, J=7.4Hz), 6.95 (1H, s).

STEP 2. 2-(4-Chlorobenzoyl)indole

A mixture of 2-(4-chlorobenzoyl)-1-(phenylsulfonyl)indole (step 1, 334 mg,
0.84 mmol) and 2N sodium hydroxide (1.5 ml, 2.78 mmol) in ethanol (5 ml) was
heated at reflux temperature for 15 min. The mixture was concentrated and the
20 residue was diluted with ethyl acetate (100 ml). The organic layer was washed with
water and dried (MgSO₄), and concentrated to afford 211 mg (98.2 %) of the title
compound as yellow solids.

¹H-NMR (CDCl₃) δ: 9.45 (1H, br.s), 7.97-7.92 (2H, m), 7.74-7.70 (1H, m), 7.54-7.47
(3H, m), 7.42-7.36 (1H, m), 7.21-7.13 (2H, m).

STEP 3. Diethyl α-acetoxy-[2-(4-chlorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4
of Example 2 (Method B) employing 2-(4-chlorobenzoyl)indole (step 2).

¹H-NMR (CDCl₃) δ: 8.94 (1H, br.s), 7.90 (1H, d, J=8.4Hz), 7.81-7.77 (2H, m), 7.43-
7.36 (3H, m), 7.32-7.26 (1H, m), 7.22-7.16 (1H, m), 4.27-4.14 (4H, m), 1.75 (3H, s),
30 1.29-1.16 (6H, m).

STEP 4. Diethyl [2-(4-chlorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) employing diethyl α -acetoxy-[2-(4-chlorobenzoyl)-1H-indol-3-yl]malonate (step 3).

- 5 $^1\text{H-NMR}$ (CDCl_3) δ : 8.73 (1H, br.s), 7.85-7.82 (1H, m), 7.79-7.76 (2H, m), 7.52-7.46 (2H, m), 7.39-7.37 (1H, m), 7.26-7.19 (2H, m), 5.27 (1H, s), 4.26-4.16 (4H, m), 1.26-1.21 (6H, m)

STEP 5. [2-(4-Chlorobenzoyl)-1H-indol-3-yl]acetic acid

- 10 The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) employing diethyl [2-(4-chlorobenzoyl)-1H-indol-3-yl]malonate (step 4).

m.p.: 221-224 °C (recrystallized from ethyl acetate/hexane).

IR (KBr) ν : 3321, 1697, 1607, 1576, 1529, 1433, 1408, 1339, 1263, 1223, 1202 cm^{-1} .

- 15 $^1\text{H-NMR}$ (DMSO-d_6) δ : 11.62 (1H, s), 7.79-7.75 (2H, m), 7.71-7.63 (3H, m), 7.46 (1H, d, $J=8.2\text{Hz}$), 7.35-7.29 (1H, m), 7.14-7.08 (1H, m), 3.84 (2H, s)

EXAMPLE 53**[6-CHLORO-2-(2-FURYL CARBONYL)-1H-INDOL-3-YL]ACETIC ACID****STEP 1. 6-chloro-2-(2-furylcarbonyl)-1-(phenylsulfonyl)indole**

- 20 The title compound was prepared according the procedure deacribed in step 2 of Example 2 (Method B) from 6-chloro-1-(phenylsulfonyl)indole (step 1 of Example 2, Method B) and 2-furoyl chloride.

$^1\text{H-NMR}$ (CDCl_3) δ : 8.11-8.19 (3 H, m), 7.73-7.74 (1 H, m), 7.51-7.65 (4 H, m), 7.27-7.31 (2 H, m), 7.10 (1 H, s), 6.62-6.64 (1 H, m).

STEP 2. 6-Chloro-2-(2-furylcarbonyl)indole

- 25 The title compound was prepared according to the procedure deacribed in step 3 of Example 2 (Method B) from 6-chloro-2-(2-furylcarbonyl)-1-(phenylsulfonyl)indole (step 1).

$^1\text{H-NMR}$ (CDCl_3) δ : 9.33 (1 H, br s), 7.65-7.73 (3 H, m), 7.46-7.48 (2 H, m), 7.12-7.16 (1 H, m), 6.64-6.66 (1 H, m).

- 30 **STEP 3. Diethyl α -Acetoxy[6-chloro-2-(2-furylcarbonyl)indol-3-yl]malonate**

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 6-chloro-2-(2-furylcarbonyl)indole (step 2).

¹H-NMR (CDCl₃) δ: 9.60 (1 H, br s), 7.76 (1 H, d, J=8.9 Hz), 7.62-7.63 (1 H, m), 7.43 (1 H, d, J=1.3 Hz), 7.28-7.29 (1 H, m), 7.13 (1 H, dd, J=1.8 Hz, 8.7 Hz), 6.59 (1 H, dd, J=1.6 Hz, 3.5 Hz), 4.18-4.32 (4 H, m), 1.88 (3 H, s), 1.18-1.28 (6 H, m).

STEP 4. Diethyl [6-chloro-2-(2-furylcarbonyl)indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α-acetoxy-[6-chloro-2-(2-furylcarbonyl)indol-3-yl]malonate (step 3).

¹H-NMR (CDCl₃) δ: 9.83 (1 H, br s), 7.67 (1 H, t, J=0.8 Hz), 7.63 (1 H, d, J=8.9 Hz), 7.40 (1 H, d, J=3.6 Hz), 7.30 (1 H, d, J=1.8 Hz), 7.01 (1 H, dd, J=1.8 Hz, 8.9 Hz), 6.62 (1 H, dd, J=1.6 Hz, 2.1 Hz), 6.19 (1 H, s), 4.20-4.32 (4 H, m), 1.27 (6 H, t, J=7.3 Hz).

STEP 5. [6-Chloro-2-(2-furylcarbonyl)indol-3-yl]acetic acid.

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [6-chloro-2-(2-furylcarbonyl)indol-3-yl]malonate (step 4).

¹H-NMR (DMSO-d₆) δ: 12.22 (1 H, br s), 11.76 (1 H, br s), 8.13 (1 H, d, J=1.0 Hz), 7.75 (1 H, d, J=8.6 Hz), 7.56 (1 H, d, J=1.8 Hz), 7.48 (1 H, d, J=3.6 Hz), 7.14 (1 H, dd, J=1.8 Hz, 8.6 Hz), 6.85 (1 H, dd, J=1.8 Hz, 3.6 Hz), 4.02 (2 H, s).

EXAMPLE 54

[6-CHLORO-2-(CYCLOHEXANECARBONYL)-1H-INDOL-3-YL]ACETIC ACID

STEP 1. 6-chloro-2-cyclohexanecarbonyl-1-(phenylsulfonyl)indole

The title compound was prepared according to the procedure described in step 2 of Example 2 (Method B) from 6-chloro-1-(phenylsulfonyl)indole (step 1 of Example 2, Method B) and cyclohexanecarbonyl chloride.

tlc: R_f=0.4 (ethyl acetate/hexane=1:4).

STEP 2. 6-Chloro-2-(cyclohexanecarbonyl)indole

The title compound was prepared according to the procedure described in step 3 of Example 2 (Method B) from 6-chloro-2-cyclohexanecarbonyl-1-(phenylsulfonyl)indole (step 1).

¹H-NMR (CDCl₃) δ: 10.08 (1H, br s), 8.08-7.04 (4H, m), 2.28-1.20 (11H, m).

STEP 3. Diethyl α-acetoxy-[6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 6-chloro-2-(cyclohexanecarbonyl)indole (step 2).

5 ¹H-NMR (CDCl₃) δ: 8.94 (1H, br s), 8.12-7.09 (3H, m), 4.34-4.21 (4H, m), 2.20 (3H, s), 1.81-1.20 (17H, m).

STEP 4. Diethyl [6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α-acetoxy-[6-chloro-2-(cyclohexanecarbonyl)-
10 1H-indol-3-yl]malonate (step 3).

¹H-NMR (CDCl₃) δ: 8.90 (1H, br s), 7.72 (1H, d, J=8.72Hz), 7.36-7.09 (2H, m), 5.70 (1H, s), 4.28-4.19 (4H, m), 1.91-1.22 (17H, m).

STEP 5. [6-Chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]
15 malonate (step 4).

m.p. : 206-209 °C

IR(KBr) ν : 3314, 2924, 2856, 1734, 1650, 1537, 1396, 1248, cm⁻¹

¹H-NMR (DMSO-d₆) δ: 11.78 (1H, s), 7.71 (1H, d, J=8.64Hz), 7.46-7.07 (2H, m), 4.01
20 (2H, s), 1.78-1.16 (11H).

EXAMPLE 55

METHYL [6-CHLORO-2-(4-METHOXYBENZOYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 8 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of
25 Example 8) and 4-methoxyphenacyl bromide.

¹H-NMR (CDCl₃) δ: 8.85 (1H, s), 7.82 (2H, d, J=8.9Hz), 7.56 (1H, d, J=8.6Hz), 7.40 (1H, d, J=1.8Hz), 7.15 (1H, dd, J=1.8, 8.6Hz), 6.99 (2H, d, J=8.6Hz), 3.90 (3H, s), 3.86 (2H, s), 3.67 (3H, s).

EXAMPLE 56

30 **[6-CHLORO-2-(4-METHOXYBENZOYL)-1H-INDOL-3-YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate (step 1).

m.p.: 187-190 °C

- 5 ¹H-NMR (DMSO-d₆) δ: 11.72 (1H, s), 7.78 (2H, d, J=8.7Hz), 7.70 (1H, d, J=8.6Hz), 7.49-7.45 (1H, m), 7.15-7.07 (3H, m), 3.87 (3H, s), 3.81 (2H, s).

EXAMPLE 57

METHYL [6-CHLORO-2-(4-ETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

- 10 A mixture of methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A, 700 mg, 1.99 mmol), 2-bromoacetyl-4-ethylpyridine* (545 mg, 2.39 mmol), potassium carbonate (1.37 g, 13.9 mmol) and acetone (20 ml) was stirred at room temperature. After stirring for 3 h, 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 0.6 ml, 3.98 mmol) was added. The resulting mixture was stirred for an
- 15 additional 19 h and then concentrated. The residue was diluted with dichloromethane (200 ml) and washed with water (100 ml x 2). The organic layer was dried (MgSO₄) and concentrated. The residue was purified by flash column chromatography eluting with ethyl acetate/hexane/dichloromethane (1:4:1) to give the title compound including impurity. The crude product was washed with ethyl acetate to give 297 mg (42%) of
- 20 the title compound as yellow solids.

¹H-NMR (CDCl₃) δ: 12.52 (1H, br s), 8.64 (1H, d, J=4.9Hz), 8.21 (1H, br s), 7.62 (1H, d, J=8.7Hz), 7.52 (1H, d, J=1.8Hz), 7.39-7.35 (1H, m), 7.13 (1H, dd, J=1.8, 8.6Hz), 4.31 (2H, s), 3.73 (3H, s), 2.78 (2H, q, J=7.6Hz), 1.32 (3H, t, J=7.6Hz).

* 2-Bromoacetyl-4-ethylpyridine was prepared as follows;

- 25 To a solution of 2-acetyl-4-ethylpyridine (E. C. Constable et al., *J. Am. Chem. Soc.*, 1997, 119, 5606., 8.37 g, 56.1 mmol) in 25% hydrobromic acid-acetic acid (150 ml) was added dropwise a solution of bromine (9.86 g, 61.7 mmol) in acetic acid (30 ml) with ice-cooling. The mixture was allowed to warm to room temperature and stirred for 2h. Diethyl ether (500 ml) was added to the mixture and the resulting
- 30 mixture was cooled with an ice-bath. A brown oil, separated out from the solution, was collected by decantation. The oil was treated with saturated aqueous sodium

bicarbonate (50 ml) and extracted with diethyl ether (300 ml). The organic layer was dried (MgSO₄) and concentrated to give 15.3 g (88%) of the title compound.

¹H-NMR (CDCl₃) δ: 8.56 (1H, d, J=5.1Hz), 7.95 (1H, br s), 7.36-7.34 (1H, m), 4.86 (2H, s), 2.74 (2H, q, J=7.7Hz), 1.29 (3H, t, J=7.6Hz).

5 **EXAMPLE 58**

[6-CHLORO-2-(4-ETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

A yellow suspension of methyl [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 57, 297 mg, 0.83 mmol) in 2N aqueous sodium hydroxide (2.5 ml) and ethanol (20 ml) was heated at reflux temperature for 3 h. After cooling to room temperature, the mixture was neutralized with 2N aqueous hydrochloric acid (2.5 ml) and concentrated. The residue was diluted with THF (150 ml), dried (MgSO₄) and concentrated. The residual solids were recrystallization from ethyl acetate to give 251 mg (88%) of the title compound as yellow solids.

15 MS (EI) m/z: 342 (M⁺).

m.p.: 215-216 °C (decomposition).

IR (KBr) ν: 3206, 1707, 1643, 1595, 1535, 1421, 1227, 1192, 1140, 912, 777 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.30 (1H, br s), 12.18 (1H, br s), 8.73 (1H, d, J=4.9Hz), 7.98 (1H, br s), 7.79 (1H, d, J=8.7Hz), 7.74 (1H, d, J=1.8Hz), 7.62 (1H, br d, J=5.1Hz), 7.12 (1H, dd, J=1.8, 8.6Hz), 4.08 (2H, s), 2.78 (2H, q, J=7.7Hz), 1.26 (3H, t, J=7.7Hz).

EXAMPLE 59

METHYL [5-CHLORO-2-(4-ETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetyl-4-ethylpyridine (Preparation is described in Example 57).

¹H-NMR (CDCl₃) δ: 12.57 (1H, br s), 8.65 (1H, d, J=5.1Hz), 8.21 (1H, br s), 7.68 (1H, br s), 7.45 (1H, d, J=8.9Hz), 7.40-7.36 (1H, m), 7.31 (1H, dd, J=2.0, 8.7Hz), 4.28 (2H, s), 3.74 (3H, s), 2.78 (2H, q, J=7.7Hz), 1.32 (3H, t, J=7.7Hz).

30 **EXAMPLE 60**

[5-CHLORO-2-(4-ETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 59).

MS (EI) m/z: 342 (M⁺).

m.p.: 217-218 °C.

IR (KBr) ν : 3269, 1705, 1643, 1595, 1533, 1418, 1335, 1225, 1200, 1059, 779 cm⁻¹.

¹H-NMR (DMSO-d₆) δ : 12.33 (1H, br s), 8.73 (1H, d, J=4.9Hz), 7.98 (1H, br s), 7.86 (1H, d, J=2.0Hz), 7.69 (1H, d, J=8.9Hz), 7.61 (1H, dd, J=4.8, 1.6Hz), 7.33 (1H, dd, J=2.0, 8.7Hz), 4.07 (2H, s), 2.78 (2H, q, J=7.6Hz), 1.26 (3H, t, J=7.6Hz).

EXAMPLE 61

METHYL [6-CHLORO-2-(4-ISOPROPYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 2-bromoacetyl-4-isopropylpyridine*.

¹H-NMR (CDCl₃) δ : 12.53 (1H, br s), 8.65 (1H, d, J=4.9Hz), 8.24 (1H, d, J=1.6Hz), 7.62 (1H, d, J=8.7Hz), 7.52 (1H, d, J=1.6Hz), 7.40 (1H, dd, J=1.8, 4.9Hz), 7.13 (1H, dd, J=1.8, 8.7Hz), 4.30 (2H, s), 3.73 (3H, s), 2.97-3.07 (1H, m), 1.32 (6H, d, J=6.9Hz).

*2-Bromoacetyl-4-isopropylpyridine was prepared from 2-acetyl-4-isopropylpyridine (K.Ishihama et al., *J. Agric. Food Chem.*, 1992, **40**, 1647) according to the procedure for preparing 2-bromoacetyl-4-ethylpyridine described in Example 57.

¹H-NMR (CDCl₃) δ : 8.57 (1H, d, J=5.8Hz), 7.97-7.98 (1H, m), 7.36-7.38 (1H, m), 4.86 (2H, s), 2.94-3.04 (1H, m), 1.27-1.32 (6H, m).

EXAMPLE 62

[6-CHLORO-2-(4-ISOPROPYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 61).

m.p.: 194-196 °C.

IR (KBr) ν : 3244, 2965, 1692, 1647, 1597, 1537, 1254, 1200, 1178, 1150, 764 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.28 (1H, br s), 8.74 (1H, d, $J=5.3\text{Hz}$), 8.01 (1H, s), 7.80 (1H, d, $J=8.9\text{Hz}$), 7.74 (1H, d, $J=1.6\text{Hz}$), 7.64-7.66 (1H, m), 7.12 (1H, dd, $J=1.8, 8.6\text{Hz}$),

5 4.08 (2H, s), 3.02-3.13 (1H, m), 1.28 (6H, d, $J=6.9\text{Hz}$).

EXAMPLE 63

METHYL [5-CHLORO-2-(4-ISOPROPYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in
10 Example 57 from methyl *trans*-5-chloro-2-(penylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetyl-4-isopropylpyridine (Preparation is described in Example 61).

$^1\text{H-NMR}$ (CDCl_3) δ : 12.57 (1H, br s), 8.65 (1H, d, $J=5.1\text{Hz}$), 8.24 (1H, d, $J=1.6\text{Hz}$),
7.67 (1H, d, $J=2.0\text{Hz}$), 7.44 (1H, d, $J=8.2\text{Hz}$), 7.40 (1H, dd, $J=1.8, 4.9\text{Hz}$), 7.31 (1H,
15 dd, $J=2.0, 8.7\text{Hz}$), 4.28 (2H, s), 3.74 (3H, s), 2.97-3.07 (1H, m), 1.32 (6H, d, $J=6.9\text{Hz}$).

EXAMPLE 64

[5-CHLORO-2-(4-ISOPROPYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in
20 Example 58 from methyl [5-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 63).

MS (EI) m/z : 356 (M^+).

m.p.: 227-228 °C.

IR (KBr) ν : 2964, 1703, 1643, 1595, 1537, 1202, 1059, 768 cm^{-1} .

25 $^1\text{H-NMR}$ (DMSO-d_6) δ : 12.32 (1H, br s), 12.15 (1H, br s), 8.74 (1H, d, $J=4.9\text{Hz}$), 8.00 (1H, s), 7.86 (1H, s), 7.64-7.71 (2H, m), 7.31-7.35 (1H, m), 4.08 (2H, s), 3.03-3.13 (1H, m), 1.28 (6H, d, $J=6.9\text{Hz}$).

EXAMPLE 65

30 METHYL [6-CHLORO-2-(4-PROPYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(penylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 2-bromoacetyl-4-propylpyridine hydrobromide*.

¹H-NMR (CDCl₃) δ: 12.53 (1H, br s), 8.64 (1H, d, J=4.9Hz), 8.19 (1H, s), 7.62 (1H, d, J=8.7Hz), 7.53 (1H, d, J=1.6Hz), 7.35-7.37 (1H, m), 7.13 (1H, dd, J=1.8, 8.7Hz), 4.31 (2H, s), 3.73 (3H, s), 2.71 (2H, t, J=7.3Hz), 1.69-1.77 (2H, m), 0.98 (3H, t, J=7.3Hz).

*2-Bromoacetyl-4-propylpyridine hydrobromide was prepared as follows;

4-Propyl-2-pyridinecarbonitrile:

The title compound was prepared from 4-propylpyridine-*N*-oxide (S.Ghersetti et al., *J. Heterocycl. Chem.*, **1969**, 6, 859) according to the procedure for preparing 4-chloro-2-pyridinecarbonitrile described in Example 33.

¹H-NMR (CDCl₃) δ: 8.59 (1H, d, J=5.1Hz), 7.53 (1H, s), 7.32-7.34 (1H, m), 2.66 (2H, t, J=7.3Hz), 1.62-1.76 (2H, m), 0.97 (3H, t, J=7.3Hz).

2-Acetyl-4-propylpyridine:

The title compound was prepared from 4-propyl-2-pyridinecarbonitrile according to the procedure for preparing 2-acetyl-4-chloropyridine described in Example 33.

¹H-NMR (CDCl₃) δ: 8.56 (1H, d, J=4.9Hz), 7.88 (1H, s), 7.27-7.30 (1H, m), 2.72 (3H, s), 2.66 (2H, t, J=7.4Hz), 1.62-1.76 (2H, m), 0.95 (3H, t, J=7.4Hz).

2-Bromoacetyl-4-propylpyridine hydrobromide:

The title compound was prepared from 2-acetyl-4-propylpyridine according to the procedure for preparing 2-bromoacetyl-4-methylpyridine hydrobromide described in step 2 of Example 31.

¹H-NMR (DMSO-d₆) δ: 8.64 (1H, d, J=4.9Hz), 7.90 (1H, d, J=1.0Hz), 7.59 (1H, dd, J=1.6, 4.9Hz), 5.02 (2H, s), 2.70 (2H, t, J=7.4Hz), 1.57-1.71 (2H, m), 0.89 (3H, t, J=7.3Hz).

EXAMPLE 66

[6-CHLORO-2-(4-PROPYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl 2-[6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 65).

m.p.: 189-191 °C.

5 IR (KBr) ν : 2964, 2928, 1711, 1645, 1595, 1533, 1281, 1225, 1192, 799 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.28 (1H, br s), 8.73 (1H, d, $J=5.1\text{Hz}$), 7.96 (1H, s), 7.73-7.81 (2H, m), 7.59 (1H, d, $J=4.9\text{Hz}$), 7.10-7.13 (1H, m), 4.08 (2H, s), 2.73 (2H, t, $J=7.1\text{Hz}$), 1.63-1.72 (2H, m), 0.92 (3H, t, $J=7.3\text{Hz}$).

EXAMPLE 67

10 METHYL [5-CHLORO-2-(4-PROPYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(penylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetyl-4-propylpyridine hydrobromide (Preparation is described in Example 65).

$^1\text{H-NMR}$ (CDCl_3) δ : 12.56 (1H, br s), 8.64 (1H, d, $J=4.9\text{Hz}$), 8.18 (1H, s), 7.67 (1H, d, $J=2.0\text{Hz}$), 7.45 (1H, d, $J=8.7\text{Hz}$), 7.29-7.37 (2H, m), 4.28 (2H, s), 3.74 (3H, s), 2.71 (2H, t, $J=7.4\text{Hz}$), 1.80-1.66 (2H, m), 0.97 (3H, t, $J=7.3\text{Hz}$).

EXAMPLE 68

20 [5-CHLORO-2-(4-PROPYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 67).

25 m.p.: 208-209 °C.

IR (KBr) ν : 3296, 2957, 1705, 1645, 1595, 1535, 1329, 1273, 1204, 1057, 795 cm^{-1}

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.33 (1H, br s), 12.15 (1H, br s), 8.73 (1H, d, $J=4.9\text{Hz}$), 7.95 (1H, s), 7.86 (1H, s), 7.69 (1H, d, $J=8.7\text{Hz}$), 7.58-7.61 (1H, m), 7.31-7.35 (1H, m), 4.08 (2H, s), 2.73 (2H, t, $J=7.4\text{Hz}$), 1.64-1.72 (2H, m), 0.92 (3H, t, $J=7.4\text{Hz}$).

30 EXAMPLE 69

METHYL [2-(4-*tert*-BUTYLPYRIDINE-2-CARBONYL)-6-CHLORO-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(penylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 2-bromoacetyl-4-*tert*-butylpyridine*.

¹H-NMR (CDCl₃) δ: 12.55 (1H, br s), 8.67 (1H, d, J=5.3Hz), 8.39 (1H, d, J=2.1Hz), 7.63 (1H, d, J=8.7Hz), 7.53-7.55 (2H, m), 7.13 (1H, dd, J=1.8, 8.7Hz), 4.31 (2H, s), 3.73 (3H, s), 1.38 (9H, s).

*2-Bromoacetyl-4-*tert*-butylpyridine was prepared from 2-acetyl-4-*tert*-butylpyridine (E.C.Constable et al., *J. Am. Chem. Soc.*, **1997**, 119, 5606) according to the procedure for preparing 2-bromoacetyl-4-ethylpyridine described in Example 57.

¹H-NMR (CDCl₃) δ: 8.58 (1H, d, J=4.8Hz), 8.11 (1H, d, J=1.6Hz), 7.51 (1H, dd, J=1.8, 5.1Hz), 4.86 (2H, s), 1.35 (9H, s).

EXAMPLE 70

[2-(4-*tert*-BUTYLPYRIDINE-2-CARBONYL)-6-CHLORO-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [2-(4-*tert*-butylpyridine-2-carbonyl)-6-chloro-1H-indol-3-yl]acetate (Example 69).

MS (EI) m/z: 370 (M⁺).

m.p.: 203-205 °C.

IR (KBr) ν: 2966, 1699, 1647, 1591, 1535, 1229 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.29 (1H, br s), 8.76 (1H, d, J=5.3Hz), 8.10 (1H, d, J=2.0Hz), 7.74-7.81 (3H, m), 7.12 (1H, dd, J=1.8, 8.6Hz), 4.08 (2H, s), 1.36 (9H, s).

EXAMPLE 71

METHYL [2-(4-*tert*-BUTYLPYRIDINE-2-CARBONYL)-5-CHLORO-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(penylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetyl-4-*tert*-butylpyridine (Preparation is described in Example 69).

¹H-NMR (CDCl₃) δ: 12.59 (1H, br s), 8.67 (1H, d, J=5.3Hz), 8.38 (1H, d, J=2.0Hz), 7.68 (1H, d, J=2.0Hz), 7.54 (2H, dd, 2.0, 5.3Hz), 7.45 (1H, d, J=8.9Hz), 7.32 (1H, dd, J=2.0, 8.9Hz), 4.29 (2H, s), 3.74 (3H, s), 1.38 (9H, s).

EXAMPLE 72

5 [2-(4-*tert*-BUTYLPYRIDINE-2-CARBONYL)-5-CHLORO-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [2-(4-*tert*-butylpyridine-2-carbonyl)-5-chloro-1H-indol-3-yl]acetate (Example 71).

10 MS (EI) m/z: 370 (M⁺).

m.p.: 209-211 °C.

IR (KBr) ν: 3269, 2968, 1746, 1705, 1589, 1531, 1236, 1207, 1177, 1150, 1059, 737 cm⁻¹.

15 ¹H-NMR (DMSO-d₆) δ: 12.33 (1H, br s), 8.76 (1H, d, J=5.3Hz), 8.10 (1H, d, J=2.0Hz), 7.86 (1H, d, J=2.0Hz), 7.78 (1H, dd, J=2.0, 5.1Hz), 7.69 (1H, d, J=8.7Hz), 7.33 (1H, dd, J=2.0, 8.7Hz), 4.08 (2H, s), 1.36 (9H, s).

EXAMPLE 73

METHYL [6-CHLORO-2-(3-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

20 The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(penylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 2-bromoacetyl-3-methylpyridine hydrobromide*.

¹H-NMR (CDCl₃) δ: 11.19 (1H, br s), 8.54 (1H, d, J=4.6Hz), 7.65 (1H, d, J=7.7Hz), 7.55 (1H, d, J=8.6Hz), 7.33-7.38 (2H, m), 7.06 (1H, dd, J=1.8, 8.7Hz), 4.15 (2H, s),
25 3.69 (3H, s), 2.59 (3H, s).

*2-Bromoacetyl-3-methylpyridine hydrobromide was prepared from 2-acetyl-3-methylpyridine (T.A.Crabb et al., *Org. Magn. Reson.*, 1982, 20, 242) according to the procedure for preparing 2-bromoacetyl-4-methylpyridine hydrobromide described in step 2 of Example 31.

¹H-NMR (DMSO-d₆) δ: 8.56 (1H, d, J=3.6Hz), 7.84 (1H, d, J=7.7Hz), 7.56-7.60 (1H, m), 5.01 (2H, s), 4.01 (3H, s).

EXAMPLE 74

[6-CHLORO-2-(3-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC

5 ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 73).

MS (EI) m/z: 328 (M⁺).

10 m.p.: 195-196 °C.

IR (KBr) ν: 3314, 1703, 1636, 1526, 1418, 1396, 1231, 1196, 1150, 1109 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.14 (1H, br s), 11.76 (1H, br s), 8.53 (1H, d, J=3.8Hz), 7.85 (1H, d, J=7.9Hz), 7.74 (1H, d, J=8.6Hz), 7.52-7.56 (2H, m), 7.10 (1H, dd, J=1.8, 8.6Hz), 3.66 (2H, s), 2.33 (3H, s).

15 EXAMPLE 75

METHYL [5-CHLORO-2-(3-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(penylsulfonylamino)cinnamate (Example 20 36, step 3) and 2-bromoacetyl-3-methylpyridine hydrobromide (Preparation is described in Example 73).

¹H-NMR (CDCl₃) δ: 11.28 (1H, br s), 8.56 (1H, d, J=3.1Hz), 7.61-7.69 (2H, m), 7.23-7.41 (3H, m), 4.15 (2H, s), 3.70 (3H, s), 2.61 (3H, s).

EXAMPLE 76

25 [5-CHLORO-2-(3-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 75).

30 m.p.: 209-211°C (decomposed).

IR (KBr) ν : 3379, 3271, 1728, 1649, 1638, 1528, 1231, 1195, 1182, 1165, 1015 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.81 (1H, br s), 8.53 (1H, d, $J=4.4\text{Hz}$), 7.85 (1H, d, $J=7.7\text{Hz}$), 7.80 (1H, d, $J=2.0\text{Hz}$), 7.49-7.56 (2H, m), 7.31 (1H, dd, $J=2.1, 8.7\text{Hz}$), 3.66 (2H, s), 2.33 (3H, s).

5 **EXAMPLE 77**

METHYL [6-CHLORO-2-(6-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of
10 Example 8, Method A) and 2-bromoacetyl-6-methylpyridine hydrobromide (H.Erlenmeyer, J.Jenni, and B.Prijs, *J.Med.Pharm.Chem.*, 1961, 3, 561-566).

$^1\text{H-NMR}$ (CDCl_3) δ : 12.58 (1H, br s), 8.15 (1H, d, $J=7.9\text{Hz}$), 7.83 (1H, t, $J=7.7\text{Hz}$), 7.62 (1H, d, $J=8.7\text{Hz}$), 7.51 (1H, d, $J=1.8\text{Hz}$), 7.40 (1H, d, $J=7.7\text{Hz}$), 7.13 (1H, dd, $J=1.8, 8.7\text{Hz}$), 4.31 (2H, s), 3.72 (3H, s), 2.76 (3H, s).

15 **EXAMPLE 78**

[6-CHLORO-2-(6-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-
20 yl]acetate (Example 77).

MS (EI) m/z : 328 (M^+).

m.p.: 230-231°C.

IR (KBr) ν : 3273, 1697, 1643, 1535, 1308, 1227, 1183, 1150, 797, 760, 671 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.06 (1H, br s), 7.99 (1H, t, $J=7.7\text{Hz}$), 7.87 (1H, d, $J=7.6\text{Hz}$),
25 7.78 (1H, d, $J=8.9\text{Hz}$), 7.74 (1H, d, $J=2.0\text{Hz}$), 7.59 (1H, d, $J=7.6\text{Hz}$), 7.13 (1H, dd, $J=2.0, 8.7\text{Hz}$), 4.04 (2H, s), 2.69 (3H, s).

EXAMPLE 79

METHYL [5-CHLORO-2-(5-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

30 The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example

36, step 3) and 2-bromoacetyl-5-methylpyridine (Preparation is described in step 1 of Example 32).

¹H-NMR (CDCl₃) δ: 12.48 (1H, br s), 8.59 (1H, d, J=2.1Hz), 8.24 (1H, d, J=8.1Hz), 7.75 (1H, dd, J=2.1, 8.1Hz), 7.67 (1H, d, J=2.0Hz), 7.44 (1H, d, J=8.7Hz), 7.31 (1H, dd, J=2.0, 8.9Hz), 4.29 (2H, s), 3.74 (3H, s), 2.48 (3H, s).

EXAMPLE 80

[5-CHLORO-2-(5-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 79).

MS (EI) m/z: 328 (M⁺).

m.p.: 247-248 °C.

IR (KBr) ν: 3288, 1699, 1638, 1531, 1427, 1329, 1285, 1246, 1209, 1177, 1059, 1016, 800, 700 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.30 (1H, br s), 12.15 (1H, br s), 8.68 (1H, br s), 8.04 (1H, d, J=7.9Hz), 7.93 (1H, br d, J=8.9Hz), 7.84 (1H, br s), 7.69 (1H, d, J=8.9Hz), 7.33 (1H, dd, J=2.0, 8.9Hz), 4.09 (2H, s), 2.46 (3H, s).

EXAMPLE 81

METHYL [6-CHLORO-2-[5-(TRIFLUOROMETHYL)PYRIDINE-2-CARBONYL]-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 2-bromoacetyl-5-(trifluoromethyl)pyridine*.

¹H-NMR (CDCl₃) δ: 11.97 (1H, br s), 9.07 (1H, br s), 8.48 (1H, d, J=8.4Hz), 8.23 (1H, dd, J=2.1, 8.4Hz), 7.63 (1H, d, J=8.6Hz), 7.52 (1H, d, J=1.8Hz), 7.15 (1H, dd, J=1.8, 8.6Hz), 4.31 (2H, s), 3.74 (3H, s).

* 2-Bromoacetyl-5-(trifluoromethyl)pyridine was prepared from 2-chloro-5-(trifluoromethyl)pyridine according to the procedure for preparing 2-bromoacetyl-5-methylpyridine described in step 1 of Example 32.

¹H-NMR (CDCl₃) δ: 8.96 (1H, br s), 8.23 (1H, br d, J=8.2Hz), 8.13 (1H, dd, J=2.1, 8.1Hz), 4.83 (2H, s).

EXAMPLE 82

[6-CHLORO-2-[5-(TRIFLUOROMETHYL)PYRIDINE-2-CARBONYL]-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(5-trifluoromethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 81).

MS (EI) m/z: 382 (M⁺).

10 m.p.: 228-229°C.

IR (KBr) ν: 3325, 1707, 1636, 1529, 1333, 1310, 1138, 1078, 1020 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.05 (1H, br s), 9.17 (1H, br s), 8.54 (1H, dd, J=2.0, 8.4Hz), 8.24 (1H, d, J=8.2Hz), 7.82 (1H, d, J=8.6Hz), 7.64 (1H, d, J=1.8Hz), 7.14 (1H, dd, J=2.0, 8.7Hz), 4.06 (2H, s).

15 **EXAMPLE 83**

METHYL [5-CHLORO-2-[5-(TRIFLUOROMETHYL)PYRIDINE-2-CARBONYL]-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 20 36, step 3) and 2-bromoacetyl-5-(trifluoromethyl)pyridine (Preparation is described in Example 81).

¹H-NMR (CDCl₃) δ: 12.01 (1H, br s), 9.05 (1H, br s), 8.45 (1H, d, J=8.2Hz), 8.21 (1H, dd, J=2.3, 8.4Hz), 7.65 (1H, br s), 7.43 (1H, d, J=8.7Hz), 7.32 (1H, dd, J=2.0, 8.7Hz), 4.27 (2H, s), 3.76 (3H, s).

25 **EXAMPLE 84**

[5-CHLORO-2-[5-(TRIFLUOROMETHYL)PYRIDINE-2-CARBONYL]-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate (Example 83).

30

MS (EI) m/z: 382 (M^+).

m.p.: 230-231°C.

IR (KBr) ν : 3300, 1720, 1701, 1641, 1531, 1327, 1235, 1163, 1130, 1078, 1020, 864 cm^{-1} .

- 5 $^1\text{H-NMR}$ (DMSO-d_6) δ : 12.09 (1H, br s), 9.17 (1H, br s), 8.54 (1H, dd, $J=2.0, 8.2\text{Hz}$), 8.23 (1H, d, $J=8.2\text{Hz}$), 7.89 (1H, d, $J=1.8\text{Hz}$), 7.62 (1H, d, $J=8.9\text{Hz}$), 7.36 (1H, dd, $J=1.8, 8.9\text{Hz}$), 4.06 (2H, s).

EXAMPLE 85

METHYL [5-CHLORO-2-(5-CHLOROPYRIDINE-2-CARBONYL)-1H-INDOL-3- 10 YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetyl-5-chloropyridine *.

- 15 $^1\text{H-NMR}$ (CDCl_3) δ : 12.01 (1H, br s), 8.72 (1H, d, $J=2.5\text{Hz}$), 8.29 (1H, d, $J=8.6\text{Hz}$), 7.93 (1H, dd, $J=2.5, 8.6\text{Hz}$), 7.65 (1H, d, $J=2.0\text{Hz}$), 7.43 (1H, d, $J=8.7\text{Hz}$), 7.31 (1H, dd, $J=2.0, 8.7\text{Hz}$), 4.27 (2H, s), 3.75 (3H, s).

* 2-Bromoacetyl-5-chloropyridine was prepared from 2-bromo-5-chloropyridine (Case, *J. Am. Chem. Soc.*, **1946**, 68, 2574) according to the procedure for preparing 2-bromoacetyl-5-methylpyridine described in step 1 of Example 32.

- 20 $^1\text{H-NMR}$ (CDCl_3) δ : 8.63 (1H, dd, $J=0.6, 2.5\text{Hz}$), 8.06 (1H, dd, $J=0.6, 8.4\text{Hz}$), 7.85 (1H, dd, $J=2.3, 8.4\text{Hz}$), 4.79 (2H, s).

EXAMPLE 86

[5-CHLORO-2-(5-CHLOROPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

- 25 The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 85).

MS (EI) m/z: 348 (M^+).

m.p.: 259-260 °C.

- 30 IR (KBr) ν : 3314, 1703, 1632, 1528, 1331, 1236, 1178, 1111, 1059, 1015, 806, 698 cm^{-1} .

¹H-NMR (DMSO-d₆) δ: 12.18 (1H, br s), 12.08 (1H, br s), 8.84 (1H, d, J=2.5Hz), 8.25 (1H, dd, J=2.5, 8.4Hz), 8.10 (1H, d, J=8.6Hz), 7.87 (1H, d, J=2.0Hz), 7.63 (1H, d, J=8.7Hz), 7.34 (1H, dd, J=2.1, 8.9Hz), 4.06 (2H, s).

EXAMPLE 87

5 METHYL [6-CHLORO-2-(5-CHLOROPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 2-bromoacetyl-5-chloropyridine (Preparation is described in Example 85).

¹H-NMR (CDCl₃) δ: 11.98 (1H, br s), 8.74 (1H, dd, J=0.7 and 2.3Hz), 8.31 (1H, dd, J=0.7, 8.6Hz), 7.94 (1H, dd, J=2.3, 8.4Hz), 7.63 (1H, d, J=8.7Hz), 7.52 (1H, d, J=1.6Hz), 7.14 (1H, dd, J=1.8, 8.7Hz), 4.30 (2H, s), 3.73 (3H, s).

EXAMPLE 88

15 [6-CHLORO-2-(5-CHLOROPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 87).

20 MS (EI) m/z: 348 (M⁺).

m.p.: 242-244 °C.

IR (KBr) ν: 3306, 1703, 1636, 1529, 1308, 1234, 1151, 1109, 698 cm⁻¹.

25 ¹H-NMR (DMSO-d₆) δ: 12.05 (1H, br s), 8.84 (1H, d, J=2.3Hz), 8.26 (1H, dd, J=2.3, 8.4Hz), 8.10 (1H, d, J=8.4Hz), 7.80 (1H, d, J=8.7Hz), 7.67 (1H, d, J=1.8Hz), 7.13 (1H, dd, J=1.8, 8.7Hz), 4.06 (2H, s).

EXAMPLE 89

METHYL [5-CHLORO-2-(4-CHLOROPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example

36, step 3) and 2-bromoacetyl-4-chloropyridine hydrobromide (Preparation is described in Example 33).

¹H-NMR (CDCl₃) δ: 12.20 (1H, br s), 8.67 (1H, d, J=5.3Hz), 8.33 (1H, d, J=2.1Hz), 7.66 (1H, d, J=2.0Hz), 7.56 (1H, dd, J=2.1, 5.3Hz), 7.43 (1H, d, J=8.7Hz), 7.32 (1H, dd, J=2.0, 8.9Hz), 4.27 (2H, s), 3.75 (3H, s).

EXAMPLE 90

[5-CHLORO-2-(4-CHLOROPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 89).

MS (EI) m/z: 348 (M⁺).

m.p.: 243-244 °C.

IR (KBr) ν: 3000, 1719, 1643, 1528, 1242, 1202, 741 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.20 (1H, br s), 8.80 (1H, d, J=5.3Hz), 8.12 (1H, d, J=2.1Hz), 7.90 (1H, dd, J=2.1, 5.4Hz), 7.66 (1H, d, J=8.9Hz), 7.34 (1H, dd, J=2.0, 8.7Hz), 4.06 (2H, s).

EXAMPLE 91

METHYL [6-CHLORO-2-(PYRIDINE-3-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 3-bromoacetylpyridine hydrobromide (G. B. Barlin, L. P. Davies, S. J. Ireland, M. M. L. Ngu, *Aust. J. Chem.*, **1989**, 42, 1735).

¹H-NMR (CDCl₃) δ: 9.26 (1H, br s), 9.00 (1H, dd, J=0.8, 2.1Hz), 8.80 (1H, dd, J=1.6, 4.8Hz), 8.10 (1H, dt, J=2.0, 2.0, 7.9Hz), 7.58 (1H, d, J=8.7Hz), 7.47 (1H, ddd, J=0.8, 4.9, 7.9Hz), 7.37 (1H, d, J=1.8Hz), 7.16 (1H, dd, J=1.8, 8.6Hz), 3.84 (2H, s), 3.65 (3H, s).

EXAMPLE 92

[6-CHLORO-2-(PYRIDINE-3-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(pyridine-3-carbonyl)-1H-indol-3-yl]acetate (Example 91).

MS (EI) m/z: 314 (M^+).

5 m.p.: 267-268 °C.

IR (KBr) ν : 3346, 1705, 1609, 1566, 1528, 1433, 1418, 1327, 1267, 1215, 943, 761 cm^{-1} .

$^1\text{H-NMR}$ (DMSO- d_6) δ : 12.28 (1H, br s), 11.86 (1H, br s), 8.90 (1H, br s), 8.85 (1H, br d, $J=4.9\text{Hz}$), 8.12 (1H, dt, $J=2.0, 2.0, 7.9\text{Hz}$), 7.77 (1H, t, $J=8.6\text{Hz}$), 7.62 (1H, dd, $J=4.9, 8.0\text{Hz}$), 7.49 (1H, d, $J=2.0\text{Hz}$), 7.15 (1H, dd, $J=1.6, 8.4\text{Hz}$), 3.85 (2H, s).

EXAMPLE 93

METHYL [6-CHLORO-2-(PYRIDINE-4-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 4-bromoacetylpyridine hydrobromide (L. W. Deady, M. S. Stanborough, *Aust. J. Chem.*, **1981**, 34, 1295).

$^1\text{H-NMR}$ (CDCl_3) δ : 9.21 (1H, br s), 8.82-8.79 (2H, m), 7.59-7.57 (2H, m), 7.56 (1H, d, $J=8.7\text{Hz}$), 7.33 (1H, d, $J=1.8\text{Hz}$), 7.16 (1H, dd, $J=1.8, 8.7\text{Hz}$), 3.80 (2H, s), 3.65 (3H, s).

EXAMPLE 94

[6-CHLORO-2-(PYRIDINE-4-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(pyridine-4-carbonyl)-1H-indol-3-yl]acetate (Example 93).

MS (EI) m/z: 314 (M^+).

m.p.: 256-257 °C.

IR (KBr) ν : 3352, 1709, 1607, 1528, 1431, 1329, 1259, 1202, 772, 687 cm^{-1} .

$^1\text{H-NMR}$ (DMSO- d_6) δ : 11.82 (1H, br s), 8.82 (2H, d, $J=5.8\text{Hz}$), 7.77 (1H, d, $J=8.6\text{Hz}$), 7.64-7.62 (2H, m), 7.48 (1H, d, $J=1.8\text{Hz}$), 7.15 (1H, dd, $J=2.0, 8.7\text{Hz}$), 3.83 (2H, s).

EXAMPLE 95**METHYL [6-CHLORO-2-[4-(HYDROXYMETHYL)PYRIDINE-2-CARBONYL]-1H-INDOL-3-YL]ACETATE****STEP 1. Methyl [6-chloro-2-[4-(tert-butyldimethylsilyloxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate**

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(penylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 2-bromoacetyl-4-(*tert*-butyldimethylsilyloxymethyl)pyridine*.

¹H-NMR (CDCl₃) δ: 12.47 (1H, br s), 8.70 (1H, d, J=4.9Hz), 8.21-8.22 (1H, m), 7.55-7.58 (2H, m), 7.50-7.51 (1H, m), 7.10 (1H, dd, J=1.8, 8.7Hz), 4.83 (2H, s), 4.31 (2H, s), 3.72 (3H, s), 0.98 (9H, s), 0.15 (6H, s).

2-Bromoacetyl-4-(*tert*-butyldimethylsilyloxymethyl)pyridine was prepared as follows;

2-Acetyl-4-(*tert*-butyldimethylsilyloxymethyl)pyridine:

The title compound was prepared from 4-(*tert*-butyldimethylsilyloxymethyl)pyridinecarbonitrile (A.Hadri et al., *J. Heterocycl. Chem.*, **1993**, 30, 631) according to the procedure for preparing 2-acetyl-4-chloropyridine described in Example 33.

¹H-NMR (CDCl₃) δ: 8.64 (1H, d, J=4.8Hz), 7.96 (1H, s), 7.50 (1H, d, J=4.8Hz), 4.80 (2H, s), 2.73 (3H, s), 0.96 (9H, s), 0.12 (6H, s).

2-Bromoacetyl-4-(*tert*-butyldimethylsilyloxymethyl)pyridine:

To a solution of 2-acetyl-4-(*tert*-butyldimethylsilyloxymethyl)pyridine (1.84 g, 6.932 mmol) in THF (50 ml) was added dropwise a solution of lithium bis(trimethylsilyl)amide (1M in THF, 8.3 ml, 8.3 mmol) at -78 °C. After stirring for 1h, chlorotriethylsilane (1.7 ml, 10.4 mmol) was added to the mixture at -78 °C. The mixture was stirred at the same temperature for 1h, then allowed to warm to 0 °C. After stirring for 1h, saturated aqueous ammonium chloride (50 ml) was added. The mixture was extracted with diethyl ether (100 ml). The organic layer was washed with water (50 ml), dried (MgSO₄) and concentrated. The residue was dissolved in THF (20 ml), and then water (4 ml) and NBS were added at 0 °C. After stirring for 1h, the mixture was diluted with diethyl ether (200 ml), washed with water (50 ml) and

dried (MgSO₄). Removal of solvent gave the crude product. Purification by flash column chromatography eluting with ethyl acetate/hexane (1:20) to afford 0.74 g (31%) of the title compound as crystals.

¹H-NMR (CDCl₃) δ: 8.63 (1H, d, J=4.8Hz), 8.02 (1H, q, J=0.8Hz), 7.53 (1H, dt, J=0.8, 4.9Hz), 4.87 (2H, s), 4.81 (2H, s), 0.96 (9H, s), 0.13 (3H, s).

STEP 2. Methyl [6-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate

To a solution of methyl [6-chloro-2-[4-(*tert*-butyldimethylsilyloxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate (step 1, 171.5 mg, 0.3625 mmol) in THF (5 ml) was added a solution of tetrabutylammonium fluoride (1M in THF, 0.54 ml, 0.54 mmol) at room temperature. After stirring for 1h, the mixture was concentrated. The residue was diluted with ethyl acetate (100 ml), washed with water (20 ml x 2) and dried (MgSO₄). Removal of solvent gave the crude product. Purification by flash column chromatography eluting with ethyl acetate/hexane/dichloromethane (1:1:1) to afford 69.6 mg (54%) of the title compound as crystals.

¹H-NMR (CDCl₃) δ: 12.43 (1H, br s), 8.74 (1H, d, J=5.1Hz), 8.29 (1H, s), 7.59-7.64 (2H, m), 7.52 (1H, d, J=1.5Hz), 7.13 (1H, dd, J=1.6, 8.6Hz), 4.86 (2H, d, J=5.1Hz), 4.31 (2H, s), 3.73 (3H, s).

20 EXAMPLE 96

[6-CHLORO-2-[4-(HYDROXYMETHYL)PYRIDINE-2-CARBONYL]-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate (Example 95).

MS (EI) m/z: 344 (M⁺).

m.p.: 210-212 °C.

IR (KBr) ν: 3304, 1728, 1713, 1622, 1583, 1526, 1194 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.31 (1H, br s), 8.78 (1H, d, J=4.8Hz), 8.08 (1H, s), 7.80 (1H, d, J=8.6Hz), 7.74 (1H, d, J=1.8Hz), 7.67 (1H, d, J=4.9Hz), 7.12 (1H, dd, J=1.8, 8.7Hz), 4.69 (2H, s), 4.09 (2H, s).

EXAMPLE 97**METHYL [5-CHLORO-2-[4-(HYDROXYMETHYL)PYRIDINE-2-CARBONYL]-1H-INDOL-3-YL]ACETATE**

5 **STEP 1. Methyl [5-chloro-2-[4-(*tert*-buthyldimethylsilyloxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate**

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(penylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetyl-4-(*tert*-buthyldimethylsilyloxymethyl)pyridine (Preparation is described in step 1 of Example 95).

10 ¹H-NMR (CDCl₃) δ: 12.47 (1H, br s), 8.66 (1H, d, J=4.9Hz), 8.18 (1H, s), 7.60-7.61 (1H, m), 7.54-7.57 (1H, m), 7.39 (1H, d, J=8.9Hz), 7.25 (1H, dd, J=2.0, 8.9Hz), 4.82 (2H, s), 4.27 (2H, s), 3.74 (3H, s), 0.98 (9H, s), 0.14 (6H, s).

STEP 2. Methyl [5-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate

15 The title compound was prepared according to the procedure described in step 2 of Example 95 from methyl [5-chloro-2-[4-(*tert*-buthyldimethylsilyloxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate (step 1).

¹H-NMR (CDCl₃) δ: 12.48 (1H, br s), 8.75 (1H, d, J=4.9Hz), 8.31 (1H, s), 7.68 (1H, s), 7.60-7.62 (1H, m), 7.45 (1H, d, J=8.91Hz), 7.32 (1H, dd, J=2.0, 8.9Hz), 4.87 (2H, d, J=5.4Hz), 4.28 (2H, s), 3.74 (3H, s).

20

EXAMPLE 98**[5-CHLORO-2-[4-(HYDROXYMETHYL)PYRIDINE-2-CARBONYL]-1H-INDOL-3-YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate (Example 97).

25

MS (EI) m/z: 344 (M⁺).

m.p.: 218-219 °C.

IR (KBr) ν: 3263, 1705, 1641, 1595, 1528, 1327, 1198, 1061 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.34 (1H, br s), 8.77 (1H, d, J=4.9Hz), 8.08 (1H, s), 7.85 (1H, s), 7.66-7.71 (2H, m), 7.33 (1H, dd, J=1.8, 8.9Hz), 5.63 (1H, br s), 4.68 (2H, s), 4.09 (2H, s).

EXAMPLE 99

5 METHYL [5-CHLORO-2-(3,4-DIMETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetyl-3,4-dimethylpyridine hydrobromide*.

10 ¹H-NMR (CDCl₃) δ: 12.20 (1H, br s), 8.67 (1H, d, J=5.3Hz), 8.33 (1H, d, J=2.1Hz), 7.66 (1H, d, J=2.0Hz), 7.56 (1H, dd, J=2.1, 5.3Hz), 7.43 (1H, d, J=8.7Hz), 7.32 (1H, dd, J=2.0, 8.9Hz), 4.27 (2H, s), 3.75 (3H, s).

* 2-Bromoacetyl-3,4-dimethylpyridine hydrobromide was prepared as follows;

3,4-Dimethylpyridine-2-carbonitrile:

15 The title compound including 4,5-dimethylpyridine-2-carbonitrile in the ratio of 5.5 to 1 was prepared from 3,4-dimethylpyridine-*N*-oxide (Abramovitch et al., *J. Org. Chem.*, 1972, 37, 1690) according to the procedure for preparing 4-chloro-2-pyridinecarbonitrile described in Example 33.

2-Acetyl-3,4-dimethylpyridine:

20 The title compound was prepared along with 2-acetyl-4,5-dimethylpyridine from 3,4-dimethylpyridine-2-carbonitrile including 4,5-dimethylpyridine-2-carbonitrile in the ratio of 5.5 to 1 according to the procedure for preparing 2-acetyl-4-chloropyridine described in Example 33.

2-acetyl-3,4-dimethylpyridine: ¹H-NMR (CDCl₃) δ: 8.34 (1H, d, J=4.6Hz), 7.19 (1H, d, J=4.8Hz), 2.69 (3H, s), 2.43 (3H, s), 2.34 (3H, s).

2-acetyl-4,5-dimethylpyridine: ¹H-NMR (CDCl₃) δ: 8.39 (1H, s), 7.83 (1H, s), 2.70 (3H, s), 2.33 (3H, s), 2.32 (3H, s).

2-Bromoacetyl-3,4-dimethylpyridine hydrobromide:

30 The title compound was prepared from 2-acetyl-3,4-dimethylpyridine according to the method of H. McKennis, Jr., L.B. Turnbull, E.R. Bowman, and E. Tamaki (in *J. Org. Chem.*, 1963, 23, 383-387).

¹H-NMR (DMSO-d₆) δ: 8.39 (1H, d, J=4.8Hz), 7.46 (1H, d, J=4.8Hz), 4.96 (2H, s), 2.38 (3H, s), 2.35 (3H, s).

EXAMPLE 100

[5-CHLORO-2-(3,4-DIMETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-

5 **YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(3,4-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 99).

MS (EI) m/z: 342 (M⁺).

10 m.p.: 236-237 °C.

IR (KBr) ν: 3395, 1710, 1641, 1526, 1339, 1281, 1196, 1053, 1007 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.76 (1H, br s), 8.37 (1H, d, J=4.9Hz), 7.79 (1H, d, J=2.0Hz), 7.48 (1H, d, J=8.7Hz), 7.41 (1H, d, J=4.9Hz), 7.31 (1H, dd, J=2.0, 8.7Hz), 3.56 (2H, s), 2.35 (3H, s), 2.16 (3H, s).

15 **EXAMPLE 101**

METHYL [5-CHLORO-2-(4,5-DIMETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 20 36, step 3) and 2-bromoacetyl-4,5-dimethylpyridine*.

¹H-NMR (CDCl₃) δ: 12.59 (1H, br s), 8.47 (1H, s), 8.12 (1H, s), 7.67 (1H, d, J=1.8Hz), 7.44 (1H, d, J=9.4Hz), 7.30 (1H, dd, J=2.0, 8.7Hz), 4.28 (2H, s), 3.73 (3H, s), 2.38 (6H, s).

* 2-Bromoacetyl-3,4-dimethylpyridine was prepared from 2-acetyl-4,5-dimethylpyridine (Preparation is described in Example 99) according to the procedure for preparing 2-bromoacetyl-4-ethylpyridine described in Example 57.

¹H-NMR (CDCl₃) δ: 8.39 (1H, s), 7.87 (1H, s), 4.83 (2H, s), 2.35 (3H, s), 2.34 (3H, s).

EXAMPLE 102

[5-CHLORO-2-(4,5-DIMETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-

30 **YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 101).

MS (EI) m/z: 342 (M^+).

5 m.p.: 245-246 °C.

IR (KBr) ν : 3281, 1697, 1638, 1589, 1535, 1254, 1232, 1188, 1060, 802 cm^{-1} .

$^1\text{H-NMR}$ (DMSO- d_6) δ : 12.34 (1H, br s), 12.15 (1H, br s), 8.57 (1H, s), 7.93 (1H, s), 7.84 (1H, d, $J=1.8\text{Hz}$), 7.69 (1H, d, $J=8.7\text{Hz}$), 7.32 (1H, dd, $J=2.0, 8.7\text{Hz}$), 4.09 (2H, s), 2.39 (3H, s), 2.38 (3H, s).

10 **EXAMPLE 103**

METHYL [6-CHLORO-2-(4,5-DIMETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of
15 Example 8, Method A) and 2-bromoacetyl-4,5-dimethylpyridine (Preparation is described in Example 99).

$^1\text{H-NMR}$ (CDCl_3) δ : 12.55 (1H, br s), 8.47 (1H, s), 8.12 (1H, s), 7.62 (1H, d, $J=8.6\text{Hz}$), 7.52 (1H, d, $J=1.8\text{Hz}$), 7.13 (1H, dd, $J=1.8, 8.6\text{Hz}$), 4.31 (2H, s), 3.72 (3H, s), 2.39 (6H, s).

20 **EXAMPLE 104**

[6-CHLORO-2-(4,5-DIMETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 103).
25

MS (EI) m/z: 342 (M^+).

m.p.: 226-228 °C.

IR (KBr) ν : 3275, 1697, 1638, 1537, 1258, 1188, 799 cm^{-1} .

$^1\text{H-NMR}$ (DMSO- d_6) δ : 12.30 (1H, br s), 8.57 (1H, s), 7.94 (1H, s), 7.78 (1H, d, $J=8.7\text{Hz}$), 7.74 (1H, d, $J=1.8\text{Hz}$), 7.11 (1H, dd, $J=2.0, 8.6\text{Hz}$), 4.09 (2H, s), 2.39 (3H, s), 2.38 (3H, s).
30

EXAMPLE 105**METHYL [6-CHLORO-2-(4-METHOXYPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE**

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(penylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 2-bromoacetyl-4-methoxypyridine hydrobromide*.

¹H-NMR (CDCl₃) δ: 12.61 (1H, br s), 8.57 (1H, d, J=5.9Hz), 7.89 (1H, d, J=2.6Hz), 7.63 (1H, d, J=8.6Hz), 7.52 (1H, d, J=1.8Hz), 7.13 (1H, dd, J=1.8, 8.7Hz), 7.04 (1H, dd, J=2.6, 5.8Hz), 4.30 (2H, s), 3.96 (3H, s), 3.73 (3H, s).

*2-Bromoacetyl-4-methoxypyridine hydrobromide was prepared from 2-acetyl-4-methoxypyridine (B. Case et al., *J. Org. Chem.*, 1961, 26, 4415) according to the procedure for preparing 2-bromoacetyl-4-methylpyridine hydrobromide described in step 2 of Example 31.

¹H-NMR (DMSO-d₆) δ: 8.59-8.62 (1H, m), 7.63-7.65 (1H, m), 7.33-7.37 (1H, m), 5.03 (2H, s), 3.96 (3H, s).

EXAMPLE 106**[6-CHLORO-2-(4-METHOXYPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 105).

MS (EI) m/z: 344 (M⁺).

m.p.: 213 °C (decomposed).

IR (KBr) ν: 3200, 1709, 1645, 1589, 1533, 1225, 1207 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.35 (1H, br s), 12.18 (1H, br s), 8.66 (1H, d, J=5.6Hz), 7.80 (1H, d, J=8.9Hz), 7.74 (1H, d, J=1.8Hz), 7.63 (1H, d, J=2.6Hz), 7.31 (1H, dd, J=2.9, 5.1Hz), 7.12 (1H, dd, J=1.3, 8.7Hz), 4.09 (2H, s), 3.96 (3H, s).

EXAMPLE 107**METHYL [5-CHLORO-2-(4-METHOXYPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE**

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(penylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetyl-4-methoxypyridine hydrobromide (Preparation is described in Example 105).

- 5 ¹H-NMR (CDCl₃) δ: 12.65 (1H, br s), 8.57 (1H, d, J=5.8Hz), 7.88 (1H, d, J=2.5Hz), 7.68 (1H, d, J=2.0Hz), 7.45 (1H, dd, J=0.5, 8.7Hz), 7.32 (1H, dd, J=1.8, 8.7Hz), 7.05 (1H, dd, J=2.8, 5.8Hz), 4.28 (2H, s), 3.96 (3H, s), 3.74 (3H, s).

EXAMPLE 108

[5-CHLORO-2-(4-METHOXYPYRIDINE-2-CARBONYL)-1H-INDOL-3-

YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 107).

m.p.: 228°C (decomposed).

- 15 IR (KBr) ν: 3230, 1707, 1647, 1595, 1566, 1533, 1477, 1331, 1300, 1219, 1180, 1038, 1013 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.39 (1H, br s), 12.16 (1H, br s), 8.66 (1H, d, J=5.8Hz), 7.86 (1H, d, J=1.5Hz), 7.70 (1H, d, J=8.7Hz), 7.63 (1H, d, J=2.5Hz), 7.29-7.35 (2H, m), 4.09 (2H, s), 3.96 (3H, s).

20 EXAMPLE 109

METHYL [6-CHLORO-2-(3,5-DIMETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(penylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 2-bromoacetyl-3,5-dimethylpyridine hydrobromide*.

- 25 ¹H-NMR (CDCl₃) δ: 11.43 (1H, br s), 8.43 (1H, s), 7.60 (1H, d, J=8.7Hz), 7.51 (1H, s), 7.45 (1H, d, J=1.8Hz), 7.12 (1H, dd, J=1.8, 8.7Hz), 4.22 (2H, s), 3.69 (3H, s), 2.62 (3H, s), 2.43 (3H, s).

*2-Bromoacetyl-3,5-dimethylpyridine hydrobromide was prepared as follows;

- 30 2-Acetyl-3,5-dimethylpyridine:

The title compound was prepared from 3,5-dimethylpyridinecarbonitrile (K.Takahashi et al., *J. Heterocycl. Chem.*, 1978, 15, 893) according to the procedure for preparing 2-acetyl-4-chloropyridine described in Example 33.

¹H-NMR (CDCl₃) δ: 8.32 (1H, s), 7.37 (1H, s), 2.69 (3H, s), 2.56 (3H, s), 2.36 (3H, s).

5 2-Bromoacetyl-3,5-dimethylpyridine hydrobromide:

The title compound was prepared from 2-acetyl-3,5-dimethylpyridine according to the method of H. McKennis, Jr., L.B.Turnbull, E.R.Bowman, and E.Tamaki (in *J.Org.Chem.*, 1963, 23, 383-387).

¹H-NMR (DMSO-d₆) δ: 8.43 (1H, s), 7.69 (1H, s), 5.00 (2H, s), 2.52 (3H, s), 2.39 (3H, s).

EXAMPLE 110

[6-CHLORO-2-(3,5-DIMETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(3,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 109).

MS (EI) m/z: 342 (M⁺).

m.p.: 199-200 °C.

IR (KBr) ν: 3279, 1705, 1639, 1526, 1238, 1177, 827, 797 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.12 (1H, br s), 11.72 (1H, br s), 8.38 (1H, s), 7.74 (1H, d, J=8.6Hz), 7.68 (1H, s), 7.52 (1H, d, J=2.0Hz), 7.10 (1H, dd, J=1.8, 8.6Hz), 3.72 (2H, s), 2.38 (3H, s), 2.33 (3H, s).

EXAMPLE 111

METHYL [5-CHLORO-2-(4-ETHYL-3-FLUOROPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(penylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetyl-4-ethyl-3-fluoropyridine*.

¹H-NMR (CDCl₃) δ: 11.29 (1H, br s), 8.45 (1H, d, J=4.8Hz), 7.64 (1H, d, J=1.8Hz), 7.44 (1H, t, J=4.8Hz), 7.38 (1H, d, J=8.9Hz), 7.29 (1H, dd, J=2.0, 8.7Hz), 4.21 (2H, s), 3.72 (3H, s), 2.81 (2H, q, J=7.6Hz), 1.31 (3H, t, J=7.6Hz).

*2-Bromoacetyl-4-ethyl-3-fluoropyridine was prepared as follows;

4-Ethyl-3-fluoropyridine-*N*-oxide:

To a mixture of 4-ethyl-3-fluoropyridine (R.P.Dikinson et al., *Bioorg. Med. Chem. Lett.*, 1996, 6, 2031, 28.51 g, 205.9 mmol) and 30% hydrogen peroxide (30 ml) in acetic acid (300 ml) was heated at reflux temperature for 3h. After cooling to room temperature, the resulting mixture was concentrated. The residue was diluted in dichloromethane (300 ml) and dried (MgSO₄). Removal of solvent gave 32.30 g (100%) of the title compound as an oil.

¹H-NMR (DMSO-d₆) δ: 8.43-8.46 (1H, m), 8.08 (1H, d, J=6.1Hz), 7.38 (1H, dd, J=6.6, 9.7Hz), 2.61 (2H, q, J=7.6Hz), 1.17 (3H, t, J=7.6Hz).

4-Ethyl-3-fluoro-2-pyridinecarbonitrile:

The title compound was prepared from 4-ethyl-3-fluoropyridine-*N*-oxide according to the procedure for preparing 4-chloro-2-pyridinecarbonitrile described in Example 33.

¹H-NMR (CDCl₃) δ: 8.42 (1H, d, J=4.8Hz), 7.43 (1H, t, J=5.1Hz), 2.78 (2H, q, J=7.6Hz), 1.30 (3H, t, J=7.6Hz).

2-Acetyl-4-ethyl-3-fluoropyridine:

The title compound was prepared from 4-ethyl-3-fluoro-2-pyridinecarbonitrile according to the procedure for preparing 2-acetyl-4-chloropyridine described in Example 33.

¹H-NMR (CDCl₃) δ: 8.376 (1H, d, J=4.6Hz), 7.36 (1H, t, J=4.8Hz), 2.76 (2H, q, J=7.6Hz), 2.71 (3H, s), 1.28 (3H, t, J=7.6Hz).

2-Bromoacetyl-4-ethyl-3-fluoropyridine:

The title compound was prepared from 2-acetyl-4-ethyl-3-fluoropyridine according to the procedure for preparing 2-bromoacetyl-4-ethylpyridine described in Example 57.

¹H-NMR (CDCl₃) δ: 8.38 (1H, d, J=4.6Hz), 7.43 (1H, d, J=4.6Hz), 4.75 (2H, s), 2.78 (2H, q, J=7.6Hz), 1.29 (3H, t, J=7.6Hz).

EXAMPLE 112

[5-CHLORO-2-(4-ETHYL-3-FLUOROPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

A stirred solution of methyl [5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 111, 391.2 mg, 1.044 mmol) in acetic acid (12 ml) and 2N aqueous HCl (4 ml) was heated at reflux temperature for 24h. After cooling to room temperature, the resulting mixture was concentrated. The residue
5 was diluted in THF (100 ml), dried (MgSO₄) and concentrated. The crude product was purified by recrystallization to afford 349.2 mg (93 %) of the title compound.

MS (EI) m/z: 361 (M⁺).

m.p.: 208 °C.

IR (KBr) ν : 3217, 1720, 1632, 1516, 1429, 1234, 1180, 1057 cm⁻¹.

10 ¹H-NMR (DMSO-d₆) δ : 11.78 (1H, br s), 8.35 (1H, d, J=4.8Hz), 7.73 (1H, d, J=2.0Hz), 7.56 (1H, t, J=5.1Hz), 7.39 (1H, d, J=8.9Hz), 7.22 (1H, dd, J=2.0, 8.9Hz), 3.65 (2H, s), 2.65 (2H, q, J=7.6Hz), 1.14 (3H, t, J=7.6Hz).

EXAMPLE 113

METHYL [6-CHLORO-2-(4-ETHYL-3-FLUOROPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE
15

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(penylsulfonylamino)cinnamate (step 1 of Example 8, Method A).

¹H-NMR (CDCl₃) δ : 11.23 (1H, br s), 8.46 (1H, d, J=4.8Hz), 7.60 (1H, d, J=9.2Hz),
20 7.42-7.46 (2H, m), 7.10-7.14 (1H, m), 4.23 (2H, s), 3.71 (3H, s), 2.81 (2H, q, J=7.7Hz), 1.31 (3H, t, J=7.7Hz).

EXAMPLE 114

[6-CHLORO-2-(3-ETHOXY-4-ETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

25 The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 113).

m.p.: 165 °C.

IR (KBr) ν : 3300, 2974, 1705, 1645, 1529, 1339, 1178 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.65 (1H, br s), 8.32 (1H, d, J=4.6Hz), 7.73 (1H, d, J=8.9Hz), 7.47-7.52 (2H, m), 7.10 (1H, dd, J=1.8, 8.6Hz), 3.89 (2H, q, J=6.9Hz), 3.70 (2H, s), 2.72 (2H, q, J=7.6Hz), 1.13-1.26 (6H, m).

EXAMPLE 115

5 METHYL [6-CHLORO-2-(3-CHLORO-4-ETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(penylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 2-bromoacetyl-3-chloro-4-ethylpyridine hydrobromide*.

10 ¹H-NMR (CDCl₃) δ: 10.05 (1H, br s), 8.47 (1H, d, J=4.9Hz), 7.53 (1H, d, J=8.7Hz), 7.32-7.34 (2H, m), 7.05 (1H, dd, J=1.8, 8.7Hz), 3.79 (2H, s), 3.61 (3H, s), 2.83 (2H, q, J=7.6Hz), 1.28 (3H, t, J=7.6Hz).

*2-Bromoacetyl-3-chloro-4-ethylpyridine hydrobromide was prepared as follows;

3-Chloro-4-ethylpyridine-*N*-oxide:

15 The title compound was prepared from 3-chloro-4-ethylpyridine (F.Marsais et al., *J.Organomet.Chem.*, **1981**, 216, 139) according to the procedure for preparing 4-ethyl-3-fluoropyridine-*N*-oxide described in Example 111.

¹H-NMR (DMSO-d₆) δ: 8.46 (1H, d, J=1.8Hz), 8.16 (1H, d, J=6.6Hz), 7.40 (1H, dd, J=1.8, 6.6Hz), 2.67 (2H, q, J=7.6Hz), 1.14-1.19 (3H, m).

20 3-Chloro-4-ethyl-2-pyridinecarbonitrile:

The title compound was prepared from 3-chloro-4-ethylpyridine-*N*-oxide according to the procedure for preparing 4-chloro-2-pyridinecarbonitrile described in Example 33.

25 ¹H-NMR (CDCl₃) δ: 8.50 (1H, d, J=4.9Hz), 7.41 (1H, t, J=4.9Hz), 2.84 (2H, q, J=7.4Hz), 1.30 (3H, t, J=7.6Hz).

2-Acetyl-3-chloro-4-ethylpyridine:

The title compound was prepared from 3-chloro-4-ethyl-2-pyridinecarbonitrile according to the procedure for preparing 2-acetyl-4-chloropyridine described in Example 33.

¹H-NMR (CDCl₃) δ: 8.41 (1H, d, J=4.8Hz), 7.29 (1H, d, J=4.8Hz), 2.83 (2H, q, J=7.6Hz), 2.67 (3H, s), 1.27 (3H, t, J=7.6Hz).

2-Bromoacetyl-3-chloro-4-ethylpyridine hydrobromide:

The title compound was prepared from 2-acetyl-3-chloro-4-ethylpyridine according to the procedure for preparing 2-bromo-4-methylpyridine hydrobromide described in step 2 of Example 31.

¹H-NMR (DMSO-d₆) δ: 8.55 (1H, d, J=4.9Hz), 7.66 (1H, d, J=4.8Hz), 4.93 (2H, s), 2.82 (2H, q, J=7.3Hz), 1.21 (3H, t, J=7.4Hz).

EXAMPLE 116

[6-CHLORO-2-(3-CHLORO-4-ETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 115).

m.p.: 158 °C.

IR (KBr) ν: 3335, 1730, 1630, 1529, 1325, 1200 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.83 (1H, br s), 8.54 (1H, d, J=4.8Hz), 7.75 (1H, d, J=8.7Hz), 7.63 (1H, d, J=4.9Hz), 7.46 (1H, s), 7.12 (1H, dd, J=1.8, 8.7Hz), 3.57 (2H, s), 2.82 (2H, q, J=7.4Hz), 1.25 (3H, t, J=7.4Hz).

EXAMPLE 117

METHYL [5-CHLORO-2-(3-CHLORO-4-ETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(penylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetyl-3-chloro-4-ethylpyridine hydrobromide (Preparation is described in Example 115).

¹H-NMR (CDCl₃) δ: 9.87 (1H, br s), 8.50 (1H, d, J=4.9Hz), 7.64 (1H, s), 7.29-7.40 (3H, m), 3.86 (2H, s), 3.63 (3H, s), 2.88 (2H, q, J=7.6Hz), 1.32 (3H, t, J=7.6Hz).

EXAMPLE 118

[5-CHLORO-2-(3-CHLORO-4-ETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 117).

m.p.: 220 °C.

- 5 IR (KBr) ν : 3341, 1695, 1626, 1533, 1458, 1429, 1229 cm^{-1} .

^1H -NMR (DMSO-d_6) δ : 11.88 (1H, br s), 8.54 (1H, d, $J=4.9\text{Hz}$), 7.82 (1H, s), 7.63 (1H, d, $J=4.9\text{Hz}$), 7.46 (1H, d, $J=8.9\text{Hz}$), 7.34 (1H, dd, $J=2.0, 8.9\text{Hz}$), 3.60 (2H, s), 2.82 (2H, q, $J=7.4\text{Hz}$), 1.25 (3H, t, $J=7.4\text{Hz}$).

EXAMPLE 119

- 10 **METHYL [5-CHLORO-2-(4,6-DIMETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE**

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetyl-4,6-dimethylpyridine*.

- 15 ^1H -NMR (CDCl_3) δ : 12.73 (1H, br s), 7.98 (1H, s), 7.67 (1H, br s), 7.42 (1H, d, $J=8.9\text{Hz}$), 7.30 (1H, dd, $J=1.8$ and 8.7Hz), 7.22 (1H, s), 4.27 (2H, s), 3.73 (3H, s), 2.70 (3H, s), 2.43 (3H, s).

- * 2-Bromoacetyl-3,4-dimethylpyridine was prepared from 2-acetyl-4,6-dimethylpyridine (Sundberg et al., *J. Am. Chem. Soc.*, **1969**, 91, 658) according to the
20 procedure for preparing 2-bromoacetyl-4-ethylpyridine described in Example 57.

^1H -NMR (CDCl_3) δ : 7.72 (1H, s), 7.19 (1H, s), 5.13 (2H, s), 2.55 (3H, s), 2.38 (3H, s).

EXAMPLE 120

[5-CHLORO-2-(4,6-DIMETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

- 25 The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 119).

MS (EI) m/z : 342 (M^+).

m.p.: 233-235 °C.

- 30 IR (KBr) ν : 3288, 2919, 1742, 1630, 1599, 1529, 1333, 1232, 1180, 1067, 772 cm^{-1} .

^1H -NMR (DMSO-d_6) δ : 11.99 (1H, br s), 7.69 (1H, s), 7.57 (1H, s), 7.52 (1H, d,

J=8.9Hz), 7.28 (1H, s), 7.18 (1H, dd, J=2.1, 8.9Hz), 3.87 (2H, s), 2.49 (3H, s), 2.27 (3H, s).

EXAMPLE 121

METHYL [6-CHLORO-2-(4,6-DIMETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 2-bromoacetyl-4,6-dimethylpyridine (Preparation is described in Example 119).

- ¹H-NMR (CDCl₃) δ: 12.69 (1H, br s), 7.98 (1H, s), 7.62 (1H, d, J=8.7Hz), 7.50 (1H, d, J=0.7Hz), 7.22 (1H, s), 7.12 (1H, dd, J=1.6, 8.6Hz), 4.30 (2H, s), 3.72 (3H, s), 2.70 (3H, s), 2.43 (3H, s).

EXAMPLE 122

[6-CHLORO-2-(4,6-DIMETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-

- YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 121).

MS (EI) m/z: 342 (M⁺).

- m.p.: 219-220 °C.

IR (KBr) ν: 3300, 1701, 1639, 1603, 1535, 1225, 1180 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.95 (1H, br s), 7.62 (1H, d, J=8.7Hz), 7.58 (1H, d, J=1.3Hz), 7.56 (1H, s), 7.27 (1H, s), 6.96 (1H, dd, J=2.0, 8.7Hz), 3.87 (2H, s), 2.49 (3H, s), 2.26 (3H, s).

- EXAMPLE 123**

METHYL [5,6-DICHLORO-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

STEP 1. Methyl *trans*-4,5-dichloro-2-nitrocinnamate

- The title compound was prepared according to the procedure described in step 1 of Example 36 from 4,5-dichloro-2-nitrobenzaldehyde (J. Kenneth et al., *J. Med. Chem.*, **1968**, 11, 946).

¹H-NMR (CDCl₃) δ: 8.20 (1H, s), 8.04 (1H, d, J=15.8Hz), 7.72 (1H, s), 6.36 (1H, d, J=15.8Hz).

STEP 2. Methyl *trans*-2-amino-4,5-dichlorocinnamate

The title compound was prepared according to the procedure described in step 2 of Example 36 from methyl *trans*-4,5-dichloro-2-nitrocinnamate (step 1).

¹H-NMR (CDCl₃) δ: 7.65 (1H, d, J=15.8Hz), 7.42 (1H, s), 7.26 (1H, s), 6.81 (1H, s), 6.28 (1H, d, J=15.8Hz).

STEP 3. Methyl *trans*-4,5-dichloro-2-(phenylsulfonylamino)cinnamate

The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-2-amino-4,5-dichlorocinnamate (step 2).

¹H-NMR (CDCl₃) δ: 7.80-7.70 (2H, m), 7.60-7.40 (6H, m), 7.02 (1H, br s), 6.13 (1H, d, J=16.1Hz), 3.79 (3H, s).

STEP 4. Methyl [(5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4,5-dichloro-2-(phenylsulfonylamino)cinnamate (step 3) and 2-bromoacetyl-4-methylpyridine hydrobromide (Preparation is described in step 2 of Example 31).

¹H-NMR (DMSO-d₆) δ: 12.45 (1H, br s), 8.69 (1H, d, J=5.1Hz), 8.14 (1H, s), 8.00-7.93 (2H, m), 7.59 (1H, d, J=4.3Hz), 4.16 (2H, s), 3.59 (3H, s), 3.32 (3H, s).

EXAMPLE 124

[5,6-DICHLORO-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The titled compound was prepared according to the procedure described in Example 58 from methyl [5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 123).

m.p.: 220-224°C.

¹H-NMR (DMSO-d₆) δ: 12.41 (1H, s), 12.20 (1H, br s), 8.69 (1H, d, J=5.1Hz), 8.11 (1H, s), 7.96-7.58 (2H, m), 7.58 (1H, d, J=4.8Hz), 4.08 (2H, s), 2.47 (3H, s).

EXAMPLE 125

METHYL [5-METHYL-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-

YL]ACETATESTEP 1. Methyl *trans*-2-amino-5-methylcinnamate

The title compound was prepared according to the procedure described in step 2 of Example 36 from methyl *trans*-5-methyl-2-nitrocinnamate (C.Venkatasubban, J.Annamalai Univ., 1933, 2, 227).

¹H-NMR (CDCl₃) δ: 7.82 (1H, d, J=15.8Hz), 7.20-7.19 (1H, m), 7.01-6.98 (1H, m), 6.63 (1H, d, J=8.2Hz), 6.35 (1H, d, J=15.8Hz), 3.83 (2H, m), 3.80 (3H, s), 2.24 (3H, s)

STEP 2. Methyl *trans*-5-methyl-2-(phenylsulfonylamino)cinnamate

The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-2-amino-5-methylcinnamate (step 1).

¹H-NMR (CDCl₃) δ: 7.70-7.67 (2H, m), 7.55-7.38 (4H, m), 7.26-7.14 (3H, m), 6.55 (1H, br s), 6.14 (1H, d, J=16.0Hz), 3.77 (3H, s), 2.33 (3H, s)

STEP 3. Methyl [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-methyl-2-(phenylsulfonylamino)cinnamate (step 2) and 2-bromoacetyl-4-methylpyridine hydrobromide (Preparation is described in step 2 of Example 31).

¹H-NMR (CDCl₃) δ: 12.32 (1H, br s), 8.61 (1H, d, J=5.0Hz), 8.17 (1H, s), 7.46 (1H, s), 7.40 (1H, d, J=8.6Hz), 7.33-7.32 (1H, m), 7.22-7.19 (1H, m), 4.31 (2H, s), 3.72 (3H, s), 2.46 (6H, m)

EXAMPLE 126[5-METHYL-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 125).

m.p.: 220-226 °C.

IR (KBr) ν: 3736, 3649, 1697, 1533, 1508, 1398, 1194, 802 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.08 (1H, br s), 8.69 (1H, d, J=4.8Hz), 7.94 (1H, s), 7.56-7.49 (3H, m), 7.17 (1H, dd, J=8.4Hz), 4.05 (2H, s), 2.46 (3H, s), 2.40 (3H, s).

EXAMPLE 127**METHYL [5-FLUORO-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE****STEP 1. Methyl *trans*-5-fluoro-2-nitrocinnamate**

- 5 The title compound was prepared according to the procedure described in step 1 of Example 36 from 5-fluoro-2-nitrobenzaldehyde.

¹H-NMR (CDCl₃) δ: 8.17-8.10 (2H, m), 7.32-7.19 (2H, m), 6.36 (1H, d, J=15.8Hz), 3.84 (3H, s).

STEP 2. Methyl *trans*-5-fluoro-2-aminocinnamate

- 10 The title compound was prepared according to the procedure described in step 2 of Example 36 from methyl *trans*-5-fluoro-2-nitrocinnamate (step 1).

¹H-NMR (CDCl₃) δ: 7.77 (1H, dd, J=15.8, 1.5Hz), 7.10-7.06 (1H, m), 6.94-6.87 (1H, m), 6.68-6.63 (1H, m), 6.33 (1H, d, J=15.8Hz), 3.85 (2H, m), 3.81 (3H, s).

STEP 3. Methyl *trans*-5-fluoro-2-(phenylsulfonylamino)cinnamate

- 15 The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-5-fluoro-2-aminocinnamate (step 2).

¹H-NMR (CDCl₃) δ: 7.68-7.65 (2H, m), 7.55-7.49 (2H, m), 7.44-7.33 (3H, m), 7.18-7.13 (1H, m), 7.10-7.04 (1H, m), 6.10 (1H, d, J=15.8Hz), 3.78 (3H, s).

STEP 4. Methyl [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate

- 20 The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-fluoro-2-(phenylsulfonylamino)cinnamate (step 3).

¹H-NMR (CDCl₃) δ: 12.46 (1H, br s), 8.58-8.56 (1H, m), 8.12 (1H, s), 7.43-7.39 (1H, m), 7.33-7.26 (2H, m), 7.14-7.06 (1H, m), 4.26 (2H, s), 3.74 (3H, s), 2.45 (3H, s).

EXAMPLE 128

- 25 **[5-Fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid**

 The title compound was prepared according to the procedure described in Example 58 from methyl [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 127).

m.p.: 223 °C (decomposed).

- 30 IR (KBr) ν: 1705, 1643, 1570, 1277, 1227, 1204, 1186 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.28 (1H, br s), 8.70 (1H, d, J=4.9Hz), 7.95 (1H, s), 7.71-7.66 (1H, m), 7.58-7.52 (2H, m), 7.25-7.17 (1H, m), 4.06 (2H, s), 2.47 (3H, s).

EXAMPLE 129

METHYL [5-METHOXY-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

STEP 1. Methyl *trans*-5-methoxy-2-nitrocinnamate

The title compound was prepared according to the procedure described in step 1 of Example 36 from 5-methoxy-2-nitrobenzaldehyde.

¹H-NMR (CDCl₃) δ: 8.21 (1H, d, J=15.8Hz), 8.16-8.12 (1H, m), 7.00-6.96 (2H, m), 6.30 (1H, d, J=15.8Hz), 3.93 (3H, s), 3.83 (3H, s).

STEP 2. Methyl *trans*-5-methoxy-2-aminocinnamate

The title compound was prepared according to the procedure described in step 2 of Example 36 from methyl *trans*-5-methoxy-2-nitrocinnamate (step 1).

¹H-NMR (CDCl₃) δ: 7.83 (1H, d, J=15.8Hz), 6.92-6.91 (1H, m), 6.82 (1H, dd, J=8.7, 2.8Hz), 6.66 (1H, d, J=8.7Hz), 6.35 (1H, d, J=15.8Hz), 3.80 (3H, s), 3.76 (3H, s).

STEP 3. Methyl *trans*-5-methoxy-2-(phenylsulfonylamino)cinnamate

The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-5-methoxy-2-aminocinnamate (step 2).

¹H-NMR (CDCl₃) δ: 7.67-7.64 (2H, m), 7.54-7.37 (4H, m), 7.24 (1H, d, J=8.7Hz), 6.95 (1H, d, J=2.8Hz), 6.89 (1H, dd, J=8.7, 2.8Hz), 6.82 (1H, br s), 6.10 (1H, d, J=15.8Hz), 3.81 (3H, s), 3.77 (3H, s).

STEP 4. Methyl [5-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-methoxy-2-(phenylsulfonylamino)cinnamate (step 3).

¹H-NMR (CDCl₃) δ: 12.38 (1H, br s), 8.61 (1H, d, J=4.9Hz), 8.17 (1H, s), 7.40 (1H, d, J=8.7Hz), 7.34-7.32 (1H, m), 7.08-7.03 (2H, m), 4.31 (2H, s), 3.88 (3H, s), 3.73 (3H, s), 2.47 (3H, s).

EXAMPLE 130

[5-METHOXY-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-

YL]ACETIC ACID

The titled compound was prepared according to the procedure described in Example 58 from methyl [5-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 129).

m.p.: 235 °C (decomposed).

5 IR (KBr) ν : 1701, 1638, 1595, 1528, 1448, 1423, 1340, 1279, 1263, 1234, 1223, 1192, 1111 cm^{-1} .

$^1\text{H-NMR}$ (DMSO- d_6) δ : 12.11 (1H, br s), 8.69 (1H, d, $J=4.9\text{Hz}$), 7.95 (1H, s), 7.58-7.55 (2H, m), 7.18 (1H, m), 7.02-6.98 (1H, m), 4.07 (2H, s), 3.79 (3H, s), 2.46 (3H, s).

EXAMPLE 131

10 METHYL [6-METHOXY-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

STEP 1. Methyl *trans*-(4-methoxy-2-nitro)cinnamate

A mixture of 4-bromo-3-nitroanisole (4.177 g, 18.0 mmol), methyl acrylate (3.24 ml, 36.0 ml), palladium(II) acetate (606.1 mg, 2.7 mmol), triphenylphosphine (1.416 g, 5.4 mmol), triethylamine (3.76 ml, 27.0 mmol) in DMF (45.0 ml) was stirred
15 at 130 °C for 4h. After cooling to ambient temperature, the mixture was concentrated. To the residue was added water (30 ml), and then the mixture was extracted with ethyl acetate-toluene (3:1, 40 ml). The aqueous layer was further extracted with ethyl acetate (2 x 30 ml). The combined organic layers were dried (MgSO_4) and
20 concentrated to afford 6.42g (quant.) of the title compound as a brown oil.

$^1\text{H-NMR}$ (CDCl_3) δ : 8.07-6.97 (4H, m), 6.31 (1H, d, 15.8Hz), 3.90 (3H, s), 3.82 (3H, s).

STEP 2. Methyl *trans*-2-amino-4-methoxycinnamate

The title compound was prepared according to the procedure described in step 2 of Example 36 from methyl *trans*-(4-methoxy-2-nitro)cinnamate (step 1).

25 $^1\text{H-NMR}$ (CDCl_3) δ : 7.80-6.21 (5H, m), 4.03 (2H, brs), 3.79 (6H, s).

STEP 3. Methyl *trans*-4-methoxy-2-(*p*-toluenesulfonylamino)cinnamate

The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-4-methoxy-2-aminocinnamate (step 2).

30 $^1\text{H-NMR}$ (CDCl_3) δ : 7.62-6.73 (9H, m), 6.07 (1H, d, 15.8Hz), 3.80 (3H, s), 3.77 (3H, s), 2.37 (3H, s).

STEP 4. Methyl [6-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-methoxy-2-(*p*-toluenesulfonylamino)cinnamate (step 3) and 2-bromoacetyl-4-methylpyridine hydrobromide (Preparation is described in step 2 of Example 31).

¹H-NMR (CDCl₃) δ: 12.32 (1H, br s), 8.61 (1H, d, 4.94Hz), 8.19-7.55 (2H, m), 7.34-6.82 (3H, m), 4.31 (2H, s), 3.89 (3H, s), 3.72 (3H, s), 2.47 (3H, s).

EXAMPLE 132[6-METHOXY-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 131).

m.p.: 204 °C.

IR (KBr) ν: 3198, 1709, 1630, 1593, 1531, 1460, 1423, 1334, 1275, 1213, 1161, 1136, 1039, 1001 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.11 (1H, br s), 8.68 (1H, d, 4.94Hz), 7.94 (1H, m), 7.62 (1H, d, 8.88Hz), 7.56-7.52 (1H, m), 7.14 (1H, d, 2.30Hz), 6.76 (1H, dd, 8.88Hz, 2.30Hz), 4.05 (2H, s), 3.82 (3H, s), 2.47 (3H, s).

EXAMPLE 133METHYL [5-ETHYL-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATESTEP 1. Methyl *trans*-5-ethyl-2-aminocinnamate

A mixture of 2-bromo-4-ethylaniline (1.0 g, 5.10 mmol), methyl acrylate (1.15 ml, 12.74 mmol), palladium(II) acetate (137 mg, 0.61 mmol), tri-*o*-tolylphosphine (745 mg, 2.45 mmol), and triethylamine (2.5 ml) in acetonitrile (10 ml) was heated at 110 °C. After stirring for 2 h, methyl acrylate (0.6 ml, 6.37 mmol), palladium(II) acetate (69 mg, 0.30 mmol), tri-*o*-tolylphosphine (372 mg, 1.22 mmol), triethylamine (1.3 ml) were added and the mixture was stirred at 110 °C for 7 h. The solvent was removed and the residue was diluted with ethyl acetate (100 ml), washed with water (100 ml), dried (MgSO₄) and concentrated. The residue was purified by flash column

chromatography eluting with ethyl acetate/hexane (1:5/1:4) to afford 793 mg (75.8 %) of the title compound as yellow solids.

¹H-NMR (CDCl₃) δ: 7.83 (1H, d, J=15.8Hz), 7.21 (1H, m), 7.04-7.01 (1H, m), 6.64 (1H, d, J=8.2Hz), 6.36 (1H, d, J=15.8Hz), 3.87 (2H, m), 3.80 (3H, s), 2.51 (2H, q, J=7.6Hz), 1.19 (3H, t, J=7.6Hz).

STEP 2. Methyl *trans*-5-ethyl-2-(phenylsulfonylamino)cinnamate

The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-5-ethyl-2-aminocinnamate (step 1).

¹H-NMR (CDCl₃) δ: 7.70-7.48 (4H, m), 7.42-7.36 (2H, m), 7.29-7.26 (2H, m), 7.20-7.17 (2H, m), 6.14 (1H, d, J=15.8Hz), 3.76 (3H, s), 2.62 (2H, q, J=7.6Hz), 1.21 (3H, t, J=7.6Hz).

STEP 3. Methyl [5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-ethyl-2-(phenylsulfonylamino)cinnamate (step 2).

¹H-NMR (CDCl₃) δ: 12.32 (1H, br s), 8.59-8.57 (1H, m), 8.14 (1H, m), 7.47-7.39 (2H, m), 7.30-7.21 (2H, m), 4.32 (2H, s), 3.73 (3H, s), 2.75 (2H, q, J=7.6Hz), 2.43 (3H, s), 1.27 (3H, t, J=7.6Hz).

EXAMPLE 134

[5-ETHYL-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC

ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 133).

m.p.: 215 °C (decomposed).

IR (KBr) ν: 1701, 1638, 1597, 1535, 1443, 1420, 1398, 1331, 1279, 1236, 1205 cm⁻¹.

¹H-NMR (DMSO) δ: 12.09 (1H, br s), 8.69 (1H, d, J=4.9Hz), 7.94 (1H, m), 7.58-7.51 (3H, m), 7.23-7.20 (1H, m), 4.06 (2H, s), 2.69 (2H, q, J=7.6Hz), 2.46 (3H, s), 1.23 (3H, t, J=7.6Hz).

EXAMPLE 135

METHYL [5-ETHYL-2-(4-ETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-

YL]ACETATE)

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-ethyl-2-(phenylsulfonylamino)cinnamate (Example 133, step 2) and 2-bromoacetyl-4-ethylpyridine (Preparation is described in Example 57).

¹H-NMR (CDCl₃) δ: 12.35 (1H, br s), 8.63 (1H, d, J=5.1Hz), 8.20 (1H, m), 7.48 (1H, s), 7.42 (1H, d, J=8.6Hz), 7.34 (1H, dd, J=5.1, 1.8Hz), 7.26-7.22 (1H, m), 4.32 (2H, s), 3.73 (3H, s), 2.76 (4H, q, J=7.6Hz), 1.30 (6H, t, J=7.6Hz)

EXAMPLE 136[5-ETHYL-2-(4-ETHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [5-ethyl-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 135).

m.p.: 206.2 °C.

IR (KBr) ν: 1703, 1636, 1595, 1533, 1431, 1333, 1283, 1236, 1188, 1117 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.09 (1H, br s), 8.73 (1H, d, J=4.9Hz), 7.97 (1H, s), 7.60-7.52 (3H, m), 7.22 (1H, d, J=8.6Hz), 4.06 (2H, s), 2.82-2.65 (4H, m), 1.28-1.21 (6H, m).

EXAMPLE 137METHYL [6-ETHYL-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATESTEP 1. Methyl *trans*-(4-acetyl-2-nitro)cinnamate

The title compound was prepared according to the procedure described in Example 131 from 4-bromo-3-nitroacetophenone and methyl acrylate .

¹H-NMR (CDCl₃) δ: 8.58-8.10 (3H, m), 7.77-7.74 (1H, m), 6.48-6.42 (1H, m), 3.85 (3H, s), 2.68 (3H, s).

STEP 2. Methyl *trans*-4-(1-hydroxyethyl)-2-nitrocinnamate

To a solution of methyl *trans*-4-acetyl-2-nitrocinnamate (462.2 mg, 1.854 mmol) in methanol (55 ml) was added sodium borohydride (176.3 mg, 4.66 mmol) at room temperature. After stirring for 10 min, the mixture was concentrated. The

residue was diluted in dichloromethane (20 ml) and washed with brine (20 ml). The aqueous layer was extracted with dichloromethane (20 ml x 2). The combined organic layers were dried (MgSO₄) and concentrated to afford 442 mg (quant.) of the title compound.

- 5 ¹H-NMR (CDCl₃) δ: 8.12-6.33 (5H, m), 5.0 (1H, m), 3.83 (3H, s), 1.54 (3H, d, 9.56Hz).

STEP 3. Methyl *trans*-(4-ethyl-2-nitro)cinnamate

- To a mixture of sodium iodide (1.668 g, 11.1 mmol) and acetonitrile (581 μl, 11.1 mmol) was added chlorotrimethylsilane (1.41 ml, 11.1 mmol) at room temperature. After stirring for 10 min, a solution of methyl *trans*-4-(1-hydroxyethyl-2-nitro)cinnamate (563.6 mg, step 1) in hexane-toluene-acetonitrile (1:1:1, 6.0 ml) was added and the mixture was stirred for an additional 48h. The reaction mixture was diluted with toluene-ethyl acetate (2:3, 25 ml) and poured into water (25 ml). The organic layer was separated and washed with 5% aqueous sodium thiosulfate (30 ml), brine (30 ml) and dried (MgSO₄). Removal of solvent gave 460 mg (quant.) of the title product as a brown oil.
- 15

¹H-NMR (CDCl₃) δ: 8.12-6.32 (5H, m), 3.83 (3H, s), 2.76 (2H, q, 7.59Hz), 1.29 (3H, t, 7.59Hz).

STEP 4. Methyl *trans*-2-amino-4-ethylcinnamate

- The title compound was prepared according to the procedure described in step 2 of Example 36 from methyl *trans*-4-ethyl-2-nitrocinnamate (step 3).
- 20

¹H-NMR (CDCl₃) δ: 7.84-6.29 (5H, m), 3.80 (3H, s), 2.57 (2H, m), 1.21 (3H, t, 7.59Hz).

STEP 5. Methyl *trans*-4-ethyl-2-(*p*-toluenesulfonylamino)cinnamate

- The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-2-amino-4-ethylcinnamate (step 4).
- 25

¹H-NMR (CDCl₃) δ: 7.57 (2H, d, 8.40Hz), 7.46 (1H, d, 15.8Hz), 7.39-7.08 (5H, m), 6.43 (1H, s), 6.43 (1H, d, 15.8Hz), 3.78 (3H, s), 2.61 (2H, m), 2.37 (3H, s), 1.18 (3H, t, 7.75Hz).

STEP 6. Methyl [6-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-

- 30 yl]acetate

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-ethyl-2-(*p*-toluenesulfonylamino)cinnamate (step 5) and 2-bromoacetyl-4-methylpyridine hydrobromide (Preparation is described in step 2 of Example 31).

- 5 ¹H-NMR (CDCl₃) δ: 12.30 (1H, br s), 8.62 (1H, d, 4.94Hz), 8.18-7.32 (4H, m), 7.05-7.02 (2H, m), 4.33 (2H, s), 3.72 (3H, s), 2.79 (2H, q, 7.59Hz), 2.48 (3H, s), 1.31 (3H, t, 7.59Hz).

EXAMPLE 138

[6-ETHYL-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC

10 ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 137).

m.p.: 204 -207 °C.

- 15 IR (KBr) ν: 3240, 1709, 1636, 1593, 1531, 1394, 1204, 1144, 1001 cm⁻¹.

¹H-NMR (CDCl₃) δ: 12.07 (1H, br s), 8.69 (1H, d, 4.94Hz), 7.93 (1H, m), 7.63 (1H, d, 8.40Hz), 7.55 (1H, m), 7.45 (1H, s), 7.00-6.96 (1H, m), 4.05 (2H, s), 2.73 (2H, q, 7.56Hz), 2.46 (3H, s), 1.24 (3H, t, 7.56Hz).

EXAMPLE 139

20 METHYL [5-ISOPROPYL-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

STEP 1. Methyl *trans*-5-isopropyl-2-aminocinnamate

The title compound was prepared according to the procedure described in step 1 of Example 133 from 2-bromo-4-isopropylaniline.

- 25 ¹H-NMR (CDCl₃) δ: 7.83 (1H, d, J=15.8Hz), 7.23 (1H, m), 7.08-7.05 (1H, m), 6.66 (1H, d, J=8.4Hz), 6.37 (1H, d, J=15.8Hz), 3.85 (2H, m), 3.81 (3H, s), 2.84-2.76 (1H, m), 1.21 (6H, m).

STEP 2. Methyl *trans*-5-isopropyl-2-(phenylsulfonylamino)cinnamate

- 30 The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-5-isopropyl-2-aminocinnamate (step 1).

¹H-NMR (CDCl₃) δ: 7.71-7.68 (2H, m), 7.58 (1H, d, J=15.8Hz), 7.51-7.20 (6H, m),

7.09 (1H, br.s), 6.16 (1H, d, J=15.8Hz), 4.11 (3H, s), 2.94-2.83 (1H, m), 1.24-1.21 (6H, m).

STEP 3. Methyl [5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate

The title compound was prepared according to the procedure described in
5 Example 57 from methyl *trans*-5-isopropyl-2-(phenylsulfonylamino)cinnamate (step 2).
¹H-NMR (CDCl₃) δ: 12.33 (1H, br s), 8.61 (1H, d, J=5.1Hz), 8.17-8.16 (1H, m), 7.50-
7.42 (2H, m), 7.33-7.26 (2H, m), 4.32 (2H, s), 3.73 (3H, s), 3.08-2.97 (1H, m), 2.46
(3H, s), 1.33-1.30 (6H, m).

EXAMPLE 140

10 [5-ISOPROPYL-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-
YL]ACETIC ACID

The title compound was prepared according to the procedure described in
Example 58 from methyl [5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-
yl]acetate (Example 139).

15 m.p.: 213 °C (decomposed).

IR (KBr) v: 1717, 1641, 1597, 1537, 1327, 1273, 1196 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.10 (1H, br s), 8.70 (1H, d, J=4.9Hz), 7.95-7.94 (1H, m),
7.59-7.54 (2H, m), 7.27 (1H, dd, J=8.6, 1.7Hz), 4.07 (2H, s), 3.03 (1H, m), 2.46 (3H, s),
1.27-1.25 (6H, m).

20 EXAMPLE 141

METHYL [2-(4-METHYLPYRIDINE-2-CARBONYL)-6-TRIFLUOROMETHYL-
1H-INDOL-3-YL]ACETATE

STEP 1. Methyl *trans*-2-amino-4-(trifluoromethyl)cinnamate

The title compound was prepared according to the procedure described in step
25 1 of Example 133 from 2-bromo-5-(trifluoromethyl)aniline.

¹H-NMR (CDCl₃) δ: 7.79 (1H, d, J=15.8Hz), 7.46-7.43 (1H, m), 7.00-6.94 (2H, m),
6.41 (1H, d, J=15.8Hz), 4.18 (2H, m), 3.82 (3H, m).

STEP 2. Methyl *trans*-2-phenylsulfonylamino-4-(trifluoromethyl)cinnamate

The title compound was prepared according to the procedure described in step
30 1 of Example 8 (Method A) from methyl *trans*-2-amino-4-(trifluoromethyl)cinnamate
(step 1).

¹H-NMR (CDCl₃) δ: 7.77-7.68 (2H, m), 7.65-7.53 (4H, m), 7.47-7.41 (3H, m), 6.24 (1H, d, J=15.8Hz), 3.80 (3H, s).

STEP 3. Methyl [2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetate

5 The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-2-phenylsulfonylamino-4-(trifluoromethyl)cinnamate (step 2).

¹H-NMR (CDCl₃) δ: 12.74 (1H, br s), 8.58 (1H, d, J=4.9Hz), 8.12 (1H, s), 7.81-7.74 (2H, m), 7.35-7.32 (2H, m), 4.32 (2H, s), 3.75 (3H, s), 2.46 (3H, s).

10 EXAMPLE 142

[2-(4-METHYLPYRIDINE-2-CARBONYL)-6-TRIFLUOROMETHYL-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetate (Example 141).

m.p.: 210-220 °C.

IR (KBr) ν: 1697, 1651, 1599, 1537, 1506, 1339, 1283, 1196, 1103, 1055 cm⁻¹.

¹H-NMR (DMSO) δ: 12.62 (1H, br s), 8.71 (1H, d, J=4.9Hz), 8.10 (1H, s), 7.99-7.97 (2H, m), 7.61-7.59 (1H, m), 7.39-7.36 (1H, m), 4.14 (2H, s), 2.48 (3H, s).

20 EXAMPLE 143

METHYL [5-*tert*-BUTYL-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

STEP 1. Methyl *trans*-5-*tert*-butyl-2-aminocinnamate

The title compound was prepared according to the procedure described in step 1 of Example 133 from 2-bromo-4-*tert*-butylaniline.

¹H-NMR (CDCl₃) δ: 7.85 (1H, d, 15.8Hz), 7.38 (1H, d, 2.30Hz), 7.23 (1H, dd, 8.56Hz, 2.30Hz), 6.67 (1H, d, 8.56Hz), 6.37 (1H, d, 15.8Hz), 3.87 (2H, br s), 3.81 (3H, s), 1.28 (9H, s).

STEP 2. Methyl *trans*-5-*tert*-butyl-2-(*p*-toluenesulfonylamino)cinnamate

30 The title compound was prepared according to the procedure described in step

1 of Example 8 (Method A) from methyl *trans*-5-*tert*-butyl-2-aminocinnamate and *p*-toluenesulfonyl chloride.

¹H-NMR (CDCl₃) δ: 7.60-7.20 (8H, m), 6.53 (1H, s), 6.17 (1H, d, 15.8Hz), 3.79 (3H, s), 2.46 (3H, s), 1.29 (s, 9H).

5 STEP 3. Methyl [5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate

The compound was prepared according to the procedure described in Example of 57 from methyl *trans*-5-*tert*-butyl-2-(*p*-toluenesulfonylamino)cinnamate (step 2) and 2-bromoacetyl-4-methylpyridine hydrobromide (Preparation is described in step 2 of Example 31).

10 ¹H-NMR (CDCl₃) δ: 12.33 (1H, br s), 8.63 (1H, d, 4.94Hz), 8.18 (1H, m), 7.62-7.32 (4H, m), 4.43 (2H, s), 3.74 (3H, s), 2.47 (3H, s), 1.40 (s, 9H).

EXAMPLE 144

[5-*tert*-BUTYL-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

15 The compound was prepared according to the procedure described in Example 9 (Method B) from methyl [5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 143).

m.p.: 191 °C.

IR (KBr) ν: 2963, 1717, 1645, 1597, 1533, 1437, 1281, 1211, 1080, 1032, 1003 cm⁻¹.

20 ¹H-NMR (DMSO-d₆) δ: 12.10 (1H, br s), 7.70 (1H, d, 5.1Hz), 7.95-7.44 (6H, m), 4.09 (2H, s), 2.46 (3H, s), 1.35 (9H, s).

EXAMPLE 145

METHYL [2-(4-METHYLPYRIDINE-2-CARBONYL)-5-TRIFLUOROMETHOXY-1H-INDOL-3-YL]ACETATE

25 STEP 1. Methyl *trans*-2-amino-5-(trifluoromethoxy)cinnamate

The title compound was prepared according to the procedure described in step 1 of Example 133 from 2-bromo-4-(trifluoromethoxy)aniline.

¹H-NMR (CDCl₃) δ: 7.74 (1H, d, 15.8Hz), 7.23-7.02 (2H, m), 6.68 (1H, d, 8.72Hz), 6.36 (1H, d, 15.8Hz), 4.00 (2H, br s), 3.81 (3H, s).

30 STEP 2. Methyl *trans*-2-*p*-toluenesulfonylamino-5-(trifluoromethoxy)cinnamate

The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-2-amino-4-(trifluoromethoxy)cinnamate (step 1) and *p*-toluenesulfonyl chloride.

¹H-NMR (CDCl₃) δ: 7.59-7.19 (8H, m), 6.76 (1H, s), 6.15 (1H, d, 15.8Hz), 3.80 (3H, s), 2.39 (3H, s).

STEP 3. Methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetate

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-2-*p*-toluenesulfonylamino-5-(trifluoromethoxy)cinnamate (step 2) and 2-bromoacetyl-4-methylpyridine hydrobromide (Preparation is described in step 2 of Example 31).

¹H-NMR (CDCl₃) δ: 12.7 (1H, br s), 8.64 (1H, d, 5.10Hz), 8.19 (1H, m), 7.55-7.23 (4H, m), 4.30 (2H, s), 3.74 (3H, s), 2.49 (3H, s).

EXAMPLE 146

15 [2-(4-METHYL-2-PYRIDINE-2-CARBONYL)-5-TRIFLUOROMETHOXY-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [2-(4-methylpyridine-2-carbonyl)]-5-trifluoromethoxy-1H-indol-3-yl]acetate (Example 145).

20 m.p.: 235-238 °C.

IR (KBr) ν: 3248, 1701, 1645, 1597, 1537, 1447, 1420, 1333, 1259, 1203, 1115, 1034, 1003 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.40 (1H, br s), 8.71 (1H, d, 4.94Hz), 7.97-7.96 (1H, m), 7.80-7.30 (4H, m), 4.09 (2H, s), 2.47 (3H, s).

25 EXAMPLE 147

METHYL [2-(4-ETHYLPYRIDINE-2-CARBONYL)-5-TRIFLUOROMETHOXY-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-2-*p*-toluenesulfonylamino-5-(trifluoromethoxy)cinnamate (step 2 of Example 145) and 2-bromoacetyl-4-ethylpyridine (Preparation is described in Example 57).

¹H-NMR (CDCl₃) δ: 8.66 (1H, d, 4.97Hz), 8.22 (1H, br s), 7.55-7.26 (4H, m), 4.30 (2H, s), 3.74 (3H, s), 2.79 (2H, q, 7.59Hz), 1.32 (3H, t, 7.59Hz).

EXAMPLE 148

[2-(4-ETHYLPYRIDINE-2-CARBONYL)-5-TRIFLUOROMETHOXY-1H-INDOL-3-

5 **YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetate (Example 147).

m.p.: 223 °C.

10 IR (KBr) ν: 3271, 1697, 1645, 1597, 1539, 1423, 1400, 1337, 1258, 1198, 1117, 1028, 1003 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.41 (1H, br s), 8.74 (1H, d, 4.94Hz), 7.99 (1H, m), 7.80-7.75 (2H, m), 7.63-7.30 (2H, m), 4.10 (2H, s), 2.78 (2H, q, 7.59Hz), 1.26 (3H, t, 7.59Hz).

EXAMPLE 149

15 **METHYL [6-METHYL-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE**

STEP 1. Methyl *trans*-2-amino-4-methylcinnamate

The title compound was prepared according to the procedure described in step 2 of Example 36 from methyl *trans*-4-methyl-2-nitrocinnamate.

20 ¹H-NMR (CDCl₃) δ: 7.81 (1H, d, J=15.8Hz), 7.30-7.26 (1H, m), 6.60-6.57 (1H, m), 6.52 (1H, m), 6.31 (1H, d, J=15.8Hz), 3.92 (2H, br s), 3.79 (3H, s), 2.26 (2H, s).

STEP 2. Methyl *trans*-4-methyl-2-(phenylsulfonylamino)cinnamate

The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-2-amino-4-methylcinnamate (step 1).

25 ¹H-NMR (CDCl₃) δ: 7.72-7.68 (2H, m), 7.55-7.49 (1H, m), 7.44-7.34 (4H, m), 7.26-7.22 (1H, m), 7.07-7.04 (1H, m), 6.62 (1H, br s), 6.12 (1H, d, J=15.8Hz), 3.76 (3H, s), 2.34 (3H, s).

STEP 3. Methyl [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate

30 The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-methyl-2-(phenylsulfonylamino)cinnamate (step 2).

¹H-NMR (CDCl₃) δ: 12.26 (1H, br s), 8.61 (1H, d, J=4.9Hz), 8.17 (1H, m), 7.58 (1H, d, J=8.4Hz), 7.34-7.29 (2H, m), 7.01-6.98 (1H, m), 4.32 (2H, s), 3.72 (3H, s), 2.49 (3H, s), 2.47 (3H, s).

EXAMPLE 150

5 [6-METHYL-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 149).

10 m.p.: 208 °C (decomposed).

IR (KBr) ν: 1707, 1638, 1593, 1531, 1277, 1205, 1142 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.02 (1H, br s), 8.68 (1H, d, J=4.9Hz), 7.93 (1H, s), 7.61 (1H, d, J=8.2Hz), 7.56-7.54 (1H, m), 7.42 (1H, s), 6.96-6.93 (1H, m), 4.04 (2H, s), 2.46 (3H, s), 2.43 (3H, s).

15 EXAMPLE 151

METHYL [2-(4-METHYLPYRIDINE-2-CARBONYL)-5-TRIFLUOROMETHYL-1H-INDOL-3-YL]ACETATE

STEP 1. Methyl *trans*-2-amino-5-(trifluoromethyl)cinnamate

The title compound was prepared according to the procedure described in step 20 1 of Example 133 from 2-bromo-4-(trifluoromethyl)aniline.

¹H-NMR (CDCl₃) δ: 7.77 (1H, d, J=15.8Hz), 7.61 (1H, s), 7.39 (1H, d, J=8.4Hz), 6.74 (1H, d, J=8.4Hz), 6.41 (1H, dd, J=15.8, 1.5Hz), 4.29 (2H, m), 3.82 (3H, m).

STEP 2. Methyl *trans*-2-phenylsulfonylamino-5-(trifluoromethyl)cinnamate

The title compound was prepared according to the procedure described in step 25 1 of Example 8 (Method A) from methyl *trans*-2-amino-5-(trifluoromethyl)cinnamate (step 1).

¹H-NMR (CDCl₃) δ: 7.79-7.76 (2H, m), 7.66 (1H, m), 7.60-7.44 (6H, m), 7.06 (1H, br s), 6.26 (1H, d, J=15.8Hz), 3.81 (3H, s).

30 STEP 3. Methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-2-phenylsulfonylamino-5-(trifluoromethyl)cinnamate (step 2).

¹H-NMR (CDCl₃) δ: 12.70 (1H, br s), 8.62 (1H, d, J=4.9Hz), 8.17 (1H, s), 8.00 (1H, s),
5 7.61-7.54 (2H, m), 7.38-7.36 (1H, m), 4.34 (2H, s), 3.75 (3H, s), 2.48 (3H, s).

EXAMPLE 152

[2-(4-METHYLPYRIDINE-2-CARBONYL)-5-TRIFLUOROMETHYL-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in
10 Example 58 from methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate (Example 151).

m.p.: 218-225 °C.

IR (KBr) ν: 1697, 1645, 1599, 1541, 1337, 1277, 1202, 1161, 1111, 1053 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.52 (1H, br s), 8.71 (1H, d, J=4.9Hz), 8.23 (1H, s), 7.97 (1H,
15 m), 7.85 (1H, d, J=8.9Hz), 7.62-7.58 (2H, m), 4.16 (2H, s), 2.48 (3H, s).

EXAMPLE 153

METHYL [2-(4-ETHYLPYRIDINE-2-CARBONYL)-5-TRIFLUOROMETHYL-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in
20 Example 57 from methyl *trans*-2-phenylsulfonylamino-5-(trifluoromethyl)cinnamate (step 2 of Example 151) and 2-bromoacetyl-4-ethylpyridine (Preparation is described in Example 57).

¹H-NMR (CDCl₃) δ: 12.65 (1H, br s), 8.64 (1H, d, J=5.1Hz), 8.21 (1H, m), 8.01 (1H, s),
7.61-7.52 (2H, m), 7.40-7.37 (1H, m), 4.34 (2H, s), 3.76 (3H, s), 2.78 (2H, q,
25 J=7.6Hz), 1.31 (3H, t, J=7.6Hz).

EXAMPLE 154

[2-(4-ETHYLPYRIDINE-2-CARBONYL)-5-TRIFLUOROMETHYL-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in
30 Example 58 from methyl [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate (Example 153).

m.p.: 203.6 °C.

IR (KBr) ν : 1703, 1647, 1599, 1537, 1340, 1202, 1105, 1051 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.52 (1H, br s), 8.74 (1H, d, $J=5.1\text{Hz}$), 8.24 (1H, s), 8.00-7.99 (1H, m), 7.86 (1H, d, $J=8.7\text{Hz}$), 7.64-7.58 (2H, m), 4.16 (2H, s), 2.79 (2H, q, $J=7.6\text{Hz}$), 1.26 (3H, t, $J=7.6\text{Hz}$).

EXAMPLE 155

METHYL (2-BENZOYL-1H-INDOL-3-YL)ACETATE

STEP 1. Methyl *trans*-2-(phenylsulfonylamino)cinnamate

The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-2-aminocinnamate.

$^1\text{H-NMR}$ (CDCl_3) δ : 7.71-7.67 (2H, m), 7.58 (1H, d, $J=15.8\text{Hz}$), 7.52-7.32 (6H, m), 7.28-7.22 (1H, m), 7.08 (1H, br s), 6.15 (1H, d, $J=15.8\text{Hz}$), 3.78 (3H, s).

STEP 2. Methyl (2-benzoyl-1H-indol-3-yl)acetate

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-2-(phenylsulfonylamino)cinnamate (step 1).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.91 (1H, br s), 7.80-7.77 (2H, m), 7.67-7.58 (2H, m), 7.53-7.48 (2H, m), 7.43-7.34 (2H, m), 7.21-7.15 (1H, m), 3.86 (2H, s), 3.65 (3H, s).

EXAMPLE 156

(2-BENZOYL-1H-INDOL-3-YL)ACETIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl (2-benzoyl-1H-indol-3-yl)acetate (Example 155).

m.p.: 194-196 °C.

IR (KBr) ν : 1713, 1597, 1541, 1450, 1402, 1267, 1180, 729 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.62 (1H, br s), 7.77-7.74 (2H, m), 7.69-7.66 (2H, m), 7.60-7.55 (2H, m), 7.48-7.45 (1H, m), 7.34-7.28 (1H, m), 7.13-7.08 (1H, m), 3.80 (2H, s).

EXAMPLE 157

METHYL [2-(4-CHLOROBENZOYL)-6-METHYL-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-methyl-2-(phenylsulfonylamino)cinnamate (step 2 of Example 149) and 4-chlorophenacyl bromide.

¹H-NMR (CDCl₃) δ: 8.84 (1H, br s), 7.74-7.71 (2H, m), 7.53-7.45 (3H, m), 7.14 (1H, m), 7.03-7.00 (1H, m), 3.81 (2H, s), 3.65 (3H, s), 2.46 (3H, s).

EXAMPLE 158

[2-(4-CHLOROBENZOYL)-6-METHYL-1H-INDOL-3-YL]ACETIC ACID

5 The title compound was prepared according to the procedure described in Example 58 from methyl [2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetate (Example 157).

m.p.: 193-195 °C.

IR (KBr) ν: 3302, 1697, 1587, 1335, 1263, 1090, 999, 770 cm⁻¹.

10 ¹H-NMR (DMSO-d₆) δ: 11.47 (1H, br s), 7.76-7.73 (2H, m), 7.65-7.61 (2H, m), 7.57 (1H, d, J=8.4Hz), 7.23 (1H, s), 6.96-6.93 (1H, m), 3.79 (2H, s), 2.42 (3H, s).

EXAMPLE 159

[2-(4-CHLOROBENZOYL)-5-METHYL-1H-INDOL-3-YL]ACETIC ACID

STEP 1. 2-(4-Chlorobenzoyl)-5-methyl-1-(phenylsulfonyl)indole

15 The title compound was prepared according to the procedure described in step 2 of Example 2 (Method B) from 5-methyl-1-(phenylsulfonyl)indole (E.Wenkert, P.Moeller, and S.Piettre, *J.Am.Chem.Soc.*, **1988**, 110, 7188-7194) and 4-chlorobenzoyl chloride.

20 ¹H-NMR (CDCl₃) δ: 8.02-7.96 (3H, m), 7.96-7.88 (2H, m), 7.57-7.43 (5H, m), 7.33-7.25 (2H, m), 6.88 (1H, s), 2.41 (3H, s).

STEP 2. 2-(4-Chlorobenzoyl)-5-methylindole

 The title compound was prepared according to the procedure described in step 3 of Example 2 (Method B) from 2-(4-chlorobenzoyl)-5-methyl-1-(phenylsulfonyl)indole (step 1).

25 ¹H-NMR (CDCl₃) δ: 9.43 (1H, br s), 7.94-7.91 (2H, m), 7.51-7.47 (3H, m), 7.37 (1H, d, J=8.4Hz), 7.21 (1H, dd, J=8.4, 1.5Hz), 7.04-7.03 (1H, m), 2.44 (3H, s).

STEP 3. Diethyl α-acetoxy-[2-(4-chlorobenzoyl)-5-methyl-1H-indol-3-yl]malonate

 The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 2-(4-chlorobenzoyl)-5-methylindole (step 3).

30 ¹H-NMR (CDCl₃) δ: 8.74 (1H, br s), 7.79 (2H, d, J=8.4Hz), 7.67 (1H, s), 7.42 (2H, d,

J=8.4Hz), 7.27-7.24 (1H, m), 7.14-7.11 (1H, m), 4.24-4.16 (4H, m), 2.45 (3H, s), 1.28-1.18 (6H, m).

STEP 4. Diethyl [2-(4-chlorobenzoyl)-5-methyl-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetoxymethyl-2-(4-chlorobenzoyl)-5-methyl-1H-indol-3-ylmalonate (step 3).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.80 (1H, br s), 7.73 (2H, d, J=8.4Hz), 7.54 (1H, s), 7.47 (2H, d, J=8.4Hz), 7.21-7.18 (1H, m), 7.13-7.10 (1H, m), 4.25-4.13 (4H, m), 2.42 (3H, s), 1.28-1.21 (6H, m).

STEP 5. [2-(4-Chlorobenzoyl)-5-methyl-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [2-(4-chlorobenzoyl)-5-methyl-1H-indol-3-yl]malonate (step 4).

m.p.: 194-197 °C.

15 IR (KBr) ν : 3308, 1695, 1609, 1529, 1402, 1263, 1223, 1088, 1015 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.18 (1H, br s), 11.49 (1H, s), 7.77-7.73 (2H, m), 7.66-7.62 (2H, m), 7.45 (1H, s), 7.35 (1H, d, J=8.6Hz), 7.15 (1H, dd, J=8.6, 1.5Hz), 3.81 (2H, s), 2.39 (3H, s).

EXAMPLE 160

20 METHYL [6-METHOXY-2-(4-CHLOROBENZOYL)-1H-INDOL-3-YL] ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-methoxy-3-(*p*-toluenesulfonylamino)cinnamate (step 3 of Example 131) and 4-chlorophenacyl bromide.

25 $^1\text{H-NMR}$ (CDCl_3) δ : 8.89 (1H, br s), 7.74-7.43 (5H, m), 6.86-6.77 (m, 2H), 3.85 (3H, s), 3.79 (3H, s), 3.65 (3H, s).

EXAMPLE 161

[6-METHOXY-2-(4-CHLOROBENZOYL)-1H-INDOL-3-YL] ACETIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-methoxy-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate (Example 160).

30

m.p. : 193 °C.

IR(KBr) v: 3308, 1701, 1628, 1603, 1526, 1427, 1335, 1271, 1205, 1148, 1092, 1032, 999 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.43 (1H, brs), 7.75-7.56 (5H, m), 6.87 (1H, d, 2.13Hz), 7.76
5 (1H, dd, 8.88Hz, 2.13Hz), 3.80 (3H, s), 3.79 (2H, s).

EXAMPLE 162

[2-(4-CHLOROBENZOYL)-6-TRIFLUOROMETHYL-1H-INDOL-3-YL]ACETIC ACID

STEP 1. 1-Phenylsulfonyl-6-(trifluoromethyl)indole

10 To a stirred mixture of 6-(trifluoromethyl)indole (500 mg, 2.70 mmol), 50 % aqueous sodium hydroxide (5 ml), water (7 ml), and tetrabutylammonium bromide (87 mg, 0.27 mmol) was added a solution of phenylsulfonyl chloride (379 µl, 2.97 mmol) in toluene (5 ml) at room temperature. After stirring for 1 h, the organic layer was separated. The aqueous layer was extracted with ethyl acetate (50 ml). The
15 combined organic layers were washed with saturated sodium bicarbonate (30 ml), water (30 ml), brine (30 ml), dried (MgSO₄), and concentrated to give 835 mg (95%) of the title compound.

¹H-NMR (CDCl₃) δ: 8.29 (1H, s), 7.91-7.88 (2H, m), 7.71 (1H, d, J=3.6Hz), 7.64-7.54 (2H, m), 7.49-7.44 (3H, m), 6.72 (1H, d, J=3.6Hz).

20 **STEP 2. 2-(4-Chlorobenzoyl)-1-phenylsulfonyl-6-(trifluoromethyl)indole**

The title compound was prepared according to the procedure described in step 2 of Example 2 (Method B) from 1-phenylsulfonyl-6-(trifluoromethyl)indole (step 1).

¹H-NMR (CDCl₃) δ: 8.42 (1H, s), 8.05-8.01 (2H, m), 7.92-7.89 (2H, m), 7.71-7.47 (7H, m), 6.95 (1H, s).

25 **STEP 3. 2-(4-Chlorobenzoyl)-6-(trifluoromethyl)indole**

The title compound was prepared according to the procedure described in step 3 of Example 2 (Method B) from 2-(4-chlorobenzoyl)-1-phenylsulfonyl-6-(trifluoromethyl)indole (step 2).

¹H-NMR (CDCl₃) δ: 9.53 (1H, br s), 7.99-7.95 (2H, m), 7.85-7.79 (2H, m), 7.56-7.53
30 (2H, m), 7.42-7.39 (1H, m), 7.19-7.18 (1H, m).

STEP 4. Diethyl α -acetoxy-[2-(4-chlorobenzoyl)-6-(trifluoromethyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 2-(4-chlorobenzoyl)-6-(trifluoromethyl)indole (step 3).

$^1\text{H-NMR}$ (CDCl_3) δ : 9.27 (1H, br s), 8.00 (1H, d, $J=8.7\text{Hz}$), 7.78 (2H, d, $J=8.6\text{Hz}$), 7.68 (1H, s), 7.45-7.39 (3H, m), 4.24-4.12 (4H, m), 1.74 (3H, s), 1.21-1.14 (6H, m).

STEP 5. Diethyl [2-(4-chlorobenzoyl)-6-(trifluoromethyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetoxy-[2-(4-chlorobenzoyl)-6-trifluoromethyl-1H-indol-3-yl]malonate (step 4).

$^1\text{H-NMR}$ (CDCl_3) δ : 9.37 (1H, br s), 7.83 (1H, d, $J=8.6\text{Hz}$), 7.77-7.73 (2H, m), 7.51-7.48 (3H, m), 7.33-7.30 (1H, m), 5.27 (1H, s), 4.25-4.06 (4H, m), 1.31-1.15 (6H, m).

STEP 6. [2-(4-Chlorobenzoyl)-5-methyl-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [2-(4-chlorobenzoyl)-6-(trifluoromethyl)-1H-indol-3-yl]malonate (step 5).

m.p.: 155-160 °C.

IR (KBr) ν : 3379, 1705, 1611, 1589, 1516, 1337, 1229, 1119, 1092, 1055 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.09 (1H, br s), 7.96-7.93 (1H, m), 7.82-7.78 (3H, m), 7.68-7.64 (2H, m), 7.41-7.37 (1H, m), 3.86 (2H, s).

EXAMPLE 163

METHYL [2-(4-CHLOROBENZOYL)-5-ETHYL-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-ethyl-2-(phenylsulfonylamino)cinnamate (Example 133, step 2).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.86 (1H, br s), 7.74-7.71 (2H, m), 7.48-7.41 (3H, m), 7.32-7.21 (2H, m), 3.85 (2H, s), 3.66 (3H, s), 2.75 (2H, q, $J=7.6\text{Hz}$), 1.28 (3H, t, $J=7.6\text{Hz}$)

EXAMPLE 164

[2-(4-CHLOROBENZOYL)-5-ETHYL-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [2-(4-chlorobenzoyl)-5-ethyl-1H-indol-3-yl]acetate (Example 163).

m.p.: 165-168 °C.

5 IR (KBr) : 3321, 1693, 1605, 1531, 1221, 1088, 1011 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.49 (1H, br s), 7.76-7.72 (2H, m), 7.65-7.61 (2H, m), 7.46 (1H, m), 7.36 (1H, d, J=8.6Hz), 7.18 (1H, dd, J=8.6, 1.5Hz), 3.81 (2H, s), 2.67 (2H, q, J=7.6Hz), 1.21 (3H, t, J=7.6Hz).

EXAMPLE 165

10 METHYL [2-(4-CHLOROBENZOYL)-5-METHOXY-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-methoxy-2-(phenylsulfonylamino)cinnamate (Example 129, step 3).

¹H-NMR (CDCl₃) δ: 8.81 (1H, br s), 7.75-7.72 (2H, m), 7.49-7.46 (2H, m), 7.29 (1H, d, J=8.9Hz), 7.04 (1H, dd, J=8.9, 2.5Hz), 6.98 (1H, d, J=2.5Hz), 3.86 (3H, s), 3.85 (2H, s), 3.67 (3H, s)

EXAMPLE 166

[2-(4-CHLOROBENZOYL)-5-METHOXY-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [2-(4-chlorobenzoyl)-5-methoxy-1H-indol-3-yl]acetate (Example 165).

m.p.: 200-204 °C.

IR (KBr) ν: 3325, 1724, 1607, 1526, 1429, 1356, 1265, 1229, 1092, 1011 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.46 (1H, br s), 7.74 (2H, d, J=8.6Hz), 7.63 (2H, d, J=8.6Hz), 7.35 (1H, d, J=8.9Hz), 7.13 (1H, d, J=2.5Hz), 6.96 (1H, dd, J=8.9, 2.5Hz), 3.83 (2H, s), 3.77 (3H, s).

EXAMPLE 167

METHYL [2-(4-CHLOROBENZOYL)-5-ISOPROPYL-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-isopropyl-2-(phenylsulfonylamino)cinnamate (Example 139, step 2).

¹H-NMR (CDCl₃) δ: 8.87 (1H, br s), 7.74-7.70 (2H, m), 7.49-7.44 (3H, m), 7.33-7.25 (2H, m), 3.85 (2H, s), 3.66 (3H, s), 3.07-2.96 (1H, m), 1.32-1.29 (6H, m)

EXAMPLE 168

[2-(4-CHLOROBENZOYL)-5-ISOPROPYL-1H-INDOL-3-YL]ACETIC ACID

5 The title compound was prepared according to the procedure described in Example 58 from methyl [2-(4-chlorobenzoyl)-5-isopropyl-1H-indol-3-yl]acetate (Example 169).

m.p.: 197-200 °C.

IR (KBr) ν: 1697, 1609, 1533, 1429, 1348, 1265, 1090, 1011 cm⁻¹.

10 ¹H-NMR (DMSO-d₆) δ: 12.18 (1H, br s), 11.50 (1H, br s), 7.77-7.74 (2H, m), 7.65-7.62 (2H, m), 7.50 (1H, m), 7.40-7.37 (1H, m), 7.26-7.23 (1H, m), 3.83 (2H, s), 3.03-2.92 (1H, m), 1.26-1.24 (6H, m).

EXAMPLE 169

15 METHYL [2-(4-CHLOROBENZOYL)-5-TRIFLUOROMETHYL-1H-INDOL-3-YL]ACETATE

 The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-2-phenylsulfonylamino-5-(trifluoromethyl)cinnamate (Example 151, step 2).

20 ¹H-NMR (CDCl₃) δ: 9.34 (1H, br s), 7.90 (1H, s), 7.77-7.70 (2H, m), 7.51-7.43 (3H, m), 7.33 (1H, d, J=8.7Hz), 3.85 (2H, s), 3.66 (3H, s)

EXAMPLE 170

[2-(4-CHLOROBENZOYL)-5-TRIFLUOROMETHYL-1H-INDOL-3-YL]ACETIC ACID

25 The title compound was prepared according to the procedure described in Example 58 from methyl [2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetate (Example 169).

m.p.: 198-200 °C.

IR (KBr) ν: 3317, 1697, 1611, 1333, 1271, 1113, 1051, 1007 cm⁻¹.

30 ¹H-NMR (DMSO-d₆) δ: 12.08 (1H, br s), 8.18 (1H, s), 7.80-7.77 (2H, m), 7.68-7.57 (4H, m), 3.92 (2H, s).

EXAMPLE 171**METHYL [2-(4-CHLOROBENZOYL)-5-TRIFLUOROMETHOXY-1H-INDOL-3-YL] ACETATE**

The title compound was prepared according to the procedure described in
5 Example 57 from methyl *trans*-3-*p*-toluenesulfonylamino-5-(trifluoromethoxy)cinnamate (Example 145, step 2) and 4-chlorophenacyl bromide.

¹H-NMR (CDCl₃) δ: 8.93 (1H, br s), 7.76 (2H, d, 8.75Hz), 7.52-7.40 (5H, m), 3.82 (2H, s), 3.68 (3H, s).

EXAMPLE 172

10 **[2-(4-CHLOROBENZOYL)-5-TRIFLUOROMETHOXY-1H-INDOL-3-YL] ACETIC ACID**

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [2-(4-chlorobenzoyl)-5-trifluoromethoxy-1H-indol-3-yl]acetate (Example 171).

15 m.p.: 195 °C.

IR (KBr) v: 3339, 1705, 1622, 1589, 1533, 1435, 1342, 1256, 1225, 1177, 1092, 1013 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.91 (1H, br s), 7.80-7.64 (5H, m), 7.55 (1H, d, 8.91Hz), 7.32-7.24 (m, 1H), 3.87 (2H, s).

20 **EXAMPLE 173**

METHYL [6-CHLORO-2-(2-METHOXYBENZOYL)-1H-INDOL-3-YL] ACETATE**STEP 1. Methyl 2-[6-chloro-2-(2-methoxybenzoyl)-1H-indol-3-yl]acetate**

The title compound was prepared according to the procedure described in Example 57 from methyl *trans* 4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of
25 Example 8, Method A) and 2-methoxyphenacyl bromide.

¹H-NMR (CDCl₃) δ: 9.02 (1H, br s), 7.60-7.46 (2H, m), 7.42-7.34 (2H, m), 7.15-6.99 (3H, m), 3.78 (3H, s), 3.66 (2H, s), 3.60 (3H, m).

EXAMPLE 174**[6-CHLORO-2-(2-METHOXYBENZOYL)-1H-INDOL-3-YL] ACETIC ACID**

30 The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-(2-methoxybenzoyl)-1H-indol-3-

yl]acetate (Example 172).

m.p.: 214-217 °C.

IR (KBr) ν : 3398, 2939, 2642, 1711, 1680, 1624, 1537, 1461, 1315, 1230, 937 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.61 (1H, s), 7.67 (1H, d, $J=8.7\text{Hz}$), 7.60-7.51 (1H, m), 7.45
5 (1H, d, $J=2.0\text{Hz}$), 7.28 (1H, dd, $J=7.4, 1.6\text{Hz}$), 7.18 (1H, d, $J=8.2\text{Hz}$), 7.12-7.04 (2H, m), 3.70 (3H, s), 3.60 (2H, s).

EXAMPLE 175

METHYL [6-CHLORO-2-(3-METHOXYBENZOYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in
10 Example 57 from methyl *trans* 4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 3-methoxyphenacyl bromide.

$^1\text{H-NMR}$ (CDCl_3) δ : 8.86 (1H, s), 7.57 (1H, d, $J=8.7\text{Hz}$), 7.46-7.26 (4H, m), 7.19-7.12 (2H, m), 3.86 (3H, s), 3.83 (2H, s), 3.67 (3H, s).

EXAMPLE 176

15 [6-CHLORO-2-(3-METHOXYBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-(3-methoxybenzoyl)-1H-indol-3-yl]acetate (Example 175).

m.p.: 227-231 °C.

20 IR (KBr) ν : 3354, 2933, 2636, 1709, 1607, 1569, 1427, 1321, 1269, 1218, 1049 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.76 (1H, s), 7.72 (1H, d, $J=8.6\text{Hz}$), 7.54-7.45 (2H, m), 7.35-7.22 (3H, m), 7.16-7.10 (1H, m), 3.83 (3H, s), 3.79 (2H, s).

EXAMPLE 177

25 METHYL [6-CHLORO-2-(3-BENZYLOXYBENZOYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans* 4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 3-benzyloxyphenacyl bromide (A.Hernandez et al., *J.Org.Chem.*, 1994, 59, 1058).

30 $^1\text{H-NMR}$ (CDCl_3) δ : 8.79 (1H, s), 7.56 (1H, d, $J=8.6\text{Hz}$), 7.46-7.33 (9H, m), 7.26-7.19 (1H, m), 7.16 (1H, dd, $J=8.6, 1.8\text{Hz}$), 5.12 (2H, s), 3.82 (2H, s), 3.64 (3H, s).

EXAMPLE 178**[6-CHLORO-2-(3-BENZYLOXYBENZOYL)-1H-INDOL-3-YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl 2-[6-chloro-2-(3-benzyloxybenzoyl)-1H-indol-3-yl]acetate (Example 177).

m.p.: 174-177 °C.

IR (KBr) ν : 3308, 3028, 2897, 1697, 1612, 1566, 1444, 1328, 1269, 1223, 732 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.76 (1H, s), 7.71 (1H, d, $J=8.6\text{Hz}$), 7.54-7.30 (10H, m), 7.13 (1H, dd, $J=8.6, 1.6\text{Hz}$), 5.18 (2H, s), 3.78 (2H, s).

EXAMPLE 179**METHYL [6-CHLORO-2-(3-HYDROXYBENZOYL)-1H-INDOL-3-YL]ACETATE**

A mixture of methyl [6-chloro-2-(3-benzyloxybenzoyl)-1H-indol-3-yl]acetate (Example 177, 0.37g, 0.85mmol) and 10% palladium-charcoal (80 mg) in ethyl acetate-methanol (5:1, 30 ml) was stirred under hydrogen atmosphere for 2.5 h. The mixture was filtered thorough a pad of Celite and the filtrate was concentrated. The solids were washed with dichloromethane (10 ml) to afford 70mg (24%) of the title compound as white solids.

$^1\text{H-NMR}$ (CDCl_3) δ : 10.36 (1H, br s), 9.06 (1H, s), 7.55 (1H, d, $J=8.7\text{Hz}$), 7.50-7.46 (1H, m), 7.36-7.22 (3H, m), 7.15-7.05 (2H, m), 3.89 (2H, s), 3.65 (3H, s).

EXAMPLE 180**[6-CHLORO-2-(3-HYDROXYBENZOYL)-1H-INDOL-3-YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-(3-hydroxybenzoyl)-1H-indol-3-yl]acetate (Example 179).

m.p.: 213-215 °C.

IR (KBr) ν : 3311, 3069, 1715, 1624, 1583, 1529, 1448, 1325, 1278, 1220, 761 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.71 (1H, s), 9.86 (1H, br s), 7.71 (1H, d, $J=8.6\text{Hz}$), 7.47 (1H, d, $J=1.5\text{Hz}$), 7.41-7.32 (1H, m), 7.20-7.02 (4H, m), 3.82 (2H, s).

EXAMPLE 181**METHYL [6-CHLORO-2-(4-BENZYLOXYBENZOYL)-1H-INDOL-3-YL]ACETATE**

The title compound was prepared according to the procedure described in Example 57 from methyl *trans* 4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 4-benzyloxyphenacyl bromide (A.Brossi et al., *J.Heterocycl.Chem.*, 1965, 2, 310).

- 5 ¹H-NMR (CDCl₃) δ: 8.87 (1H, br s), 7.81 (2H, d, J=8.6Hz), 7.55 (1H, d, J=8.7Hz), 7.48-7.32 (6H, m), 7.14 (1H, dd, J=8.7,1.8Hz), 7.06 (2H, d, J=8.6Hz), 5.16 (2H, s), 3.86 (2H, s), 3.65 (3H, s).

EXAMPLE 182

[6-CHLORO-2-(4-BENZYLOXYBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

- 10 The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl 2-[6-chloro-2-(4-benzyloxybenzoyl)-1H-indol-3-yl]acetate (Example 181).

m.p.: 220-222 °C.

IR (KBr) ν: 3331, 3013, 2914, 1717, 1699, 1599, 1564, 1508, 1253, 1167, 941 cm⁻¹.

- 15 ¹H-NMR (DMSO-d₆) δ: 11.72 (1H, s), 7.76 (2H, d, J=8.9Hz), 7.69 (1H, d, J=8.4Hz), 7.52-7.32 (6H, m), 7.17 (2H, d, J=8.9Hz), 7.11 (1H, dd, J=8.4,2.2Hz), 5.23 (2H, s), 3.81 (2H, s).

EXAMPLE 183

METHYL [6-CHLORO-2-(4-HYDROXYBENZOYL)-1H-INDOL-3-YL]ACETATE

- 20 The title compound was prepared according to the procedure described in Example 179 from methyl [6-chloro-2-(4-benzyloxybenzoyl)-1H-indol-3-yl]acetate (Example 181).

- ¹H-NMR (acetone-d₆) δ: 10.82 (1H, br s), 9.21 (1H, br s), 7.75 (2H, d, J=8.7Hz), 7.71 (1H, d, J=8.7Hz), 7.56 (1H, d, J=2.0Hz), 7.13 (1H, dd, J=8.7, 2.0Hz), 6.97 (2H, d, J=8.7Hz), 3.95 (2H, s), 3.59 (3H, s).
- 25

EXAMPLE 184

[6-CHLORO-2-(4-HYDROXYBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

- The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-(4-hydroxybenzoyl)-1H-indol-3-yl]acetate (Example 183).
- 30

m.p.: 231-234 °C.

IR (KBr) ν : 3250, 3120, 2822, 1734, 1618, 1539, 1456, 1321, 1236, 1120, 1060 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.58 (1H, br s), 7.63-7.53 (3H, m), 7.35 (1H, d, $J=1.9\text{Hz}$), 7.00 (1H, dd, $J=8.6, 1.9\text{Hz}$), 6.80 (2H, d, $J=8.9\text{Hz}$), 3.69 (2H, s).

5 **EXAMPLE 185**

METHYL [6-CHLORO-2-(4-ISOPROPOXYBENZOYL)-1H-INDOL-3-YL]ACETATE

To a stirred solution of methyl [6-chloro-2-(4-hydroxybenzoyl)-1H-indol-3-yl]acetate (Example 183, 0.15 g, 0.44 mmol) in DMF (5 ml) was added sodium hydride (15 mg, 0.46 mmol) at room temperature under nitrogen atmosphere. After 5 min., 2-iodopropane (78 mg) was added and the mixture was stirred for 7 h. The mixture was quenched with 2N aqueous HCl (20 ml), and extracted with ethyl acetate (50 ml). The extract was washed with water (30 ml) and brine (30 ml), dried (MgSO_4), and concentrated. The residue was purified by flash column chromatography eluting with ethyl acetate-hexane (1:2) to afford 72 mg (43%) of the title compound as white solids.

$^1\text{H-NMR}$ (CDCl_3) δ : 8.81 (1H, br s), 7.80 (2H, d, $J=8.9\text{Hz}$), 7.56 (1H, d, $J=8.7\text{Hz}$), 7.41 (1H, d, $J=1.8\text{Hz}$), 7.15 (1H, dd, $J=8.7, 1.8\text{Hz}$), 6.95 (2H, d, $J=8.9\text{Hz}$), 4.67 (1H, heptet, $J=6.1\text{Hz}$), 3.87 (2H, s), 3.67 (3H, s), 1.39 (6H, d, $J=6.1\text{Hz}$).

EXAMPLE 186

20 [6-CHLORO-2-(4-ISOPROPOXYBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-(4-isopropoxybenzoyl)-1H-indol-3-yl]acetate (Example 185).

m.p.: 216-218 °C.

25 IR (KBr) ν : 3304, 2972, 2881, 1707, 1614, 1596, 1560, 1508, 1311, 1261, 1163 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.71 (1H, br s), 7.75 (2H, d, $J=8.9\text{Hz}$), 7.69 (1H, d, $J=8.6\text{Hz}$), 7.46 (1H, d, $J=2.0\text{Hz}$), 7.12 (1H, dd, $J=8.6, 2.0\text{Hz}$), 7.07 (2H, d, $J=8.9\text{Hz}$), 4.78 (1H, heptet, $J=5.9\text{Hz}$), 3.82 (2H, s), 1.32 (6H, d, $J=5.9\text{Hz}$).

EXAMPLE 187

30 METHYL [6-CHLORO-2-(4-PHENYLBENZOYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans* 4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 4-phenylphenacyl bromide.

¹H-NMR (CDCl₃) δ: 8.91 (1H, br s), 7.88 (2H, d, J=8.1Hz), 7.73 (2H, d, J=8.1Hz),
5 7.68-7.38 (7H, m), 7.16 (1H, dd, J=8.1,1.8Hz), 3.89 (2H, s), 3.66 (3H, s).

EXAMPLE 188

[6-CHLORO-2-(4-PHENYLBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-(4-phenylbenzoyl)-1H-indol-3-
10 yl]acetate (Example 187).

m.p.: 228-231 °C.

IR (KBr) ν: 3317, 3030, 2868, 1707, 1620, 1600, 1527, 1431, 1323, 1256, 1194 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.81 (1H, s), 7.89 (2H, d, J=8.6Hz), 7.87 (2H, d, J=8.6Hz),
7.82-7.72 (3H, m), 7.58-7.40 (4H, m), 7.14 (1H, dd, J=8.6, 1.8Hz), 3.87 (2H, s).

EXAMPLE 189

METHYL [6-CHLORO-2-(4-TRIFLUOROMETHOXYBENZOYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans* 4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of
20 Example 8, Method A) and 4-(trifluoromethoxy)phenacyl bromide*.

¹H-NMR (CDCl₃) δ: 8.85 (1H, br s), 7.85 (2H, d, J=8.9Hz), 7.57 (1H, d, J=8.6Hz),
7.40 (1H, d, J=1.8Hz), 7.35 (2H, d, J=8.9Hz), 7.16 (1H, dd, J=8.6,1.8Hz), 3.80 (2H, s),
3.65 (3H, s).

*4-(Trifluoromethoxy)phenacyl bromide was prepared as follows;

25 A mixture of 4-(trifluoromethoxy)acetophenone (0.52 g, 2.55 mmol) and tetrabutylammonium tribromide (1.35 , 2.80 mmol) in dichloromethane-methanol (1:1, 8 ml) was stirred for 18h and then concentrated. The residue was diluted with ethyl acetate (50 ml), washed with water (50 ml), brine (50 ml), and dried (MgSO₄). Removal of solvent gave 0.36 g (50 %) of the title compound as a yellow oil.

30 tlc: R_f = 0.47 (hexane-ethyl acetate=10:1).

EXAMPLE 190**[6-CHLORO-2-(4-TRIFLUOROMETHOXYBENZOYL)-1H-INDOL-3-YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-(4-trifluoromethoxybenzoyl)1H-indol-3-yl]acetate (Example 189).

m.p.: 166-168 °C.

IR (KBr) ν : 3315, 3219, 1719, 1699, 1616, 1527, 1508, 1254, 1221, 1167, 943 cm^{-1} .

$^1\text{H-NMR}$ (DMSO- d_6) δ : 11.79 (1H, s), 7.88 (2H, d, $J=8.6\text{Hz}$), 7.75 (1H, d, $J=8.9\text{Hz}$), 7.56 (2H, d, $J=8.6\text{Hz}$), 7.47 (1H, d, $J=1.8\text{Hz}$), 7.14 (1H, dd, $J=8.9, 1.8\text{Hz}$), 3.85 (2H, s).

EXAMPLE 191**METHYL [5-CHLORO-2-(4-TRIFLUOROMETHOXYBENZOYL)-1H-INDOL-3-YL]ACETATE**

The title compound was prepared according to the procedure described in Example 8 (Method B) from methyl *trans* 5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 4-(trifluoromethoxy)phenacyl bromide (Preparation is described in Example 189).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.92 (1H, br s), 7.85 (2H, d, $J=8.7\text{Hz}$), 7.61 (1H, br s), 7.38-7.28 (4H, m), 3.79 (2H, s), 3.67 (3H, s).

EXAMPLE 192**[5-CHLORO-2-(4-TRIFLUOROMETHOXYBENZOYL)-1H-INDOL-3-YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetate (Example 191).

m.p.: 186.7 °C.

IR (KBr) ν : 3332, 3086, 2925, 1697, 1610, 1408, 1259, 1217, 1161, 1007, 941 cm^{-1} .

$^1\text{H-NMR}$ (DMSO- d_6) δ : 11.85 (1H, br s), 7.89 (2H, d, $J=8.7\text{Hz}$), 7.81 (1H, d, $J=2.1\text{Hz}$), 7.60-7.53 (2H, m), 7.48 (1H, d, $J=8.7\text{Hz}$), 7.32 (1H, dd, $J=8.7, 2.1\text{Hz}$), 3.86 (2H, s).

EXAMPLE 193

METHYL [5-CHLORO-2-(4-METHOXYBENZOYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans* 5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 4-methoxyphenacyl bromide.

- 5 ¹H-NMR (CDCl₃) δ: 8.86 (1H, s), 7.82 (2H, d, J=8.9Hz), 7.63-7.60 (1H, m), 7.37-7.26 (2H, m), 6.99 (2H, d, J=8.9Hz), 3.90 (3H, s), 3.84 (2H, s), 3.69 (3H, s).

EXAMPLE 194**[5-CHLORO-2-(4-METHOXYBENZOYL)-1H-INDOL-3-YL]ACETIC ACID**

- The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate (Example 193).

m.p.: 232-235 °C.

IR (KBr) ν: 3312, 2833, 2621, 1701, 1599, 1510, 1454, 1263, 1167, 1001, 777 cm⁻¹.

- 15 ¹H-NMR (DMSO-d₆) δ: 11.78 (1H, s), 7.78 (2H, d, J=8.9Hz), 7.75 (1H, d, J=2.0Hz), 7.47 (1H, d, J=8.7Hz), 7.29 (1H, dd, J=8.7,2.0Hz), 7.11 (2H, d, J=8.9Hz), 3.88 (3H, s), 3.82 (2H, s).

EXAMPLE 195**METHYL [6-CHLORO-2-(4-NITROBENZOYL)-1H-INDOL-3-YL]ACETATE**

- The title compound was prepared according to the procedure described in Example 57 from methyl *trans* 4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 4-nitrophenacyl bromide.

¹H-NMR (CDCl₃) δ: 8.92 (1H, br s), 8.37 (2H, d, J=8.9Hz), 7.93 (2H, d, J=8.9Hz), 7.57 (1H, d, J=8.7Hz), 7.40 (1H, d, J=1.8Hz), 7.18 (1H, dd, J=8.7,1.8Hz), 3.73 (2H, s), 3.65 (3H, s).

- 25 **EXAMPLE 196**

[6-CHLORO-2-(4-NITROBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-(4-nitrobenzoyl)-1H-indol-3-yl]acetate (Example 195).

- 30 m.p.: 218.9 °C (decomposed).

IR (KBr) ν : 3365, 3101, 2846, 1718, 1699, 1647, 1537, 1348, 1255, 1227, 852 cm^{-1} .

$^1\text{H-NMR}$ ($\text{DMSO-}d_6$) δ : 11.84 (1H, s), 8.38 (2H, d, $J=8.9\text{Hz}$), 7.97 (2H, d, $J=8.9\text{Hz}$), 7.78 (1H, d, $J=8.6\text{Hz}$), 7.48 (1H, d, $J=2.0\text{Hz}$), 7.15 (1H, dd, $J=8.6, 2.0\text{Hz}$), 3.83 (2H, s).

EXAMPLE 197

5 METHYL [6-CHLORO-2-[(4-METHYLSULFONYL)BENZOYL]-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from 4-(methylsulfonyl)phenacyl bromide.

$^1\text{H-NMR}$ (CDCl_3) δ : 9.10 (1H, br s), 8.08 (2H, d, $J=8.6\text{Hz}$), 7.93 (2H, d, $J=8.3\text{Hz}$), 7.57
10 (1H, d, $J=8.7\text{Hz}$), 7.37 (1H, br), 7.20-7.10 m(1H, m), 3.78 (2H, s), 3.64 (3H, s), 3.12 (3H, s).

EXAMPLE 198

[6-CHLORO-2-[(4-METHYLSULFONYL)BENZOYL]-1H-INDOL-3-YL]ACETIC ACID

15 The title compound was prepared according to the procedure described in Example 9 (method B) from methyl [6-chloro-2-(4-methylsulfonyl)benzoyl-1H-indol-3-yl]acetate (Example 197).

m.p.: 241 $^{\circ}\text{C}$ (decomposed).

IR (KBr) ν : 3330, 1713, 1614, 1524, 1230, 1150, 941, 781 cm^{-1} .

20 $^1\text{H-NMR}$ ($\text{DMSO-}d_6$) δ : 11.83 (1H, s), 8.12 (2H, d, $J=8.3\text{Hz}$), 7.96 (2H, d, $J=8.3\text{Hz}$), 7.78 (1H, d, $J=8.7\text{Hz}$), 7.48 (1H, d, $J=1.8\text{Hz}$), 7.15 (1H, dd, $J=1.8, 8.7\text{Hz}$), 3.85 (2H, s), 3.33 (3H, s).

EXAMPLE 199

25 METHYL [6-CHLORO-2-[4-(METHYLSULFONYLAMINO)BENZOYL]-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 8 (Method B) from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 4-(methylsulfonylamino)phenacyl bromide*.

$^1\text{H-NMR}$ (CDCl_3) δ : 8.86 (1H, br s), 7.82 (2H, d, $J=8.72\text{Hz}$), 7.58 (1H, d, $J=8.72\text{Hz}$),
30 7.42- 7.15 (4H, m), 6.80 (1H, br s), 3.83 (2H, s), 3.67 (3H, s), 3.13 (3H, s).

*4-(Methylsulfonylamino)phenacyl bromide was prepared from (4-

acetylphenyl)methanesulfonamide (R. Lis, et al., *J. Org. Chem.*, **1987**, 52, 4377) according to the procedure for preparing 4-(trifluoromethoxy)phenacyl bromide described in Example 189.

¹H-NMR (CDCl₃) δ: 8.01 (2H, d, 8.72Hz), 7.28 (2H, d, 8.72Hz), 4.40 (2H, s), 3.13 (2H, s)

EXAMPLE 200

[6-CHLORO-2-[4-(METHYLSULFONYLAMINO)BENZOYL]-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-[4-(methylsulfonylamino)benzoyl]-1H-indol-3-yl]acetate (Example 199).

m.p.: 207-210 °C.

IR (KBr) ν: 3333, 3248, 1715, 1603, 1570, 1529, 1508, 1394, 1323, 1259, 1231, 1159, 1061 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.6 (1H, br s), 7.76 (2H, d, J=8.88Hz), 7.71 (1H, d, J=8.56Hz), 7.46 (1H, 1.65Hz), 7.12 (1H, dd, J=8.56Hz, 1.65Hz), 3.81 (2H, s), 3.13 (3H, s).

EXAMPLE 201

[6-CHLORO-2-(2-CHLOROBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

STEP 1. 6-Chloro-2-(2-chlorobenzoyl)-1-(phenylsulfonyl)indole

The title compound was prepared according to the procedure described in step 2 of Example 1 (Method B) from 6-chloro-1-(phenylsulfonyl)indole (step 1 of Example 1, Method B) and 2-chlorobenzoyl chloride. The crude product was used for the next step without further purification.

tlc: R_f = 0.25 (hexane-ethyl acetate=4:1).

STEP 2. 6-Chloro-2-(2-chlorobenzoyl)indole

The title compound was prepared according to the procedure described in step 3 of Example 1 (Method B) from 6-chloro-2-(2-chlorobenzoyl)-1-(phenylsulfonyl)indole (step 1). The crude product was used for the next step without further purification.

tlc: R_f = 0.37 (hexane-ethyl acetate=4:1)

STEP 3. Diethyl α -acetoxy-2-[6-chloro-2-(2-chlorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 1 (Method B) from 6-chloro-2-(2-chlorobenzoyl)indole (step 2).

¹H-NMR (CDCl₃) δ : 8.61 (1H, br s), 7.79 (1H, d, J=8.88Hz), 7.52-7.33 (5H, m), 7.15 (1H, dd, J=1.97Hz, 8.88Hz), 4.26 (4H, m), 2.02 (3H, s), 1.22 (6H, t, J=7.07Hz).

STEP 4. Diethyl [6-chloro-2-(2-chlorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 1 (Method B) from diethyl α -acetoxy-[6-chloro-2-(2-chlorobenzoyl)-1H-indol-3-yl]malonate (step 3).

¹H-NMR (CDCl₃) δ : 8.95 (1H, br s), 7.75 (1H, d, J=8.88Hz), 7.53-7.39 (5H, m), 4.83 (1H, s), 4.15 (4H, m), 1.20 (6H, t, J=7.26Hz).

STEP 5. 2-[6-Chloro-2-(2-chlorobenzoyl)-1H-indol-3-yl]acetic acid

The compound was prepared according to the procedure described in step 6 of Example 1 (Method B) from diethyl [6-chloro-2-(2-chlorobenzoyl)-1H-indol-3-yl]malonate (step 4).

m.p.: 138-140 °C.

IR (KBr) ν : 3315, 1713, 1632, 1564, 1526, 1435, 1325, 1254, 1215, 1151, 1061 cm⁻¹.

¹H-NMR (DMSO-d₆) δ : 11.67 (1H, s), 7.61 (1H, d, J=8.72), 7.51-7.34 (5H, m), 7.50 (1H, dd, J=1.97, 8.72Hz), 3.75 (2H, s).

EXAMPLE 202**[6-CHLORO-2-(2,4-DICHLOROBENZOYL)-1H-INDOL-3-YL]ACETIC ACID****STEP 1. 6-Chloro-2-(2,4-dichlorobenzoyl)-1-(phenylsulfonyl)indole**

The title compound was prepared according to the procedure described in step 2 of Example 1 (Method B) from 6-chloro-1-(phenylsulfonyl)indole (step 1 of Example 1, Method B) and 2,4-dichlorobenzoyl chloride. The crude product was used for the next step without further purification.

tlc: R_f = 0.34 (hexane-ethyl acetate = 4:1).

STEP 2. 6-Chloro-2-(2,4-dichlorobenzoyl)indole

The title compound was prepared according to the procedure described in step 3 of Example 1 (Method B) from 6-chloro-2-(2,4-dichlorobenzoyl)-1-(phenylsulfonyl)indole (step 1). The crude product was used for the next step without

further purification.

tlc: R_f = 0.45 (hexane-ethyl acetate =4:1)

STEP 3. Diethyl α -acetoxy-[6-chloro-2-(2,4-dichlorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 4 of Example 1 (Method B) from 6-chloro-2-(2,4-dichlorobenzoyl)indole (step 2).

¹H-NMR (CDCl₃) δ : 8.92 (1H, br s), 7.79-7.11 (6H, m), 4.24 (4H, m), 2.02 (3H, s), 1.22 (6H, t, J=7.10Hz).

STEP 4. Diethyl [6-chloro-2-(2,4-dichlorobenzoyl)-1H-indol-3-yl]malonate

The title compound was prepared according to the procedure described in step 5 of Example 1 from diethyl α -acetoxy-[6-chloro-2-(2,4-dichlorobenzoyl)-1H-indol-3-yl]malonate (step 3).

¹H-NMR (CDCl₃) δ : 8.92 (1H, br s), 7.75 (1H, d, J=8.92Hz), 7.55-7.36 (4H, m), 7.13 (1H, dd, 8.92Hz, 1.81Hz), 4.85 (1H, s), 4.17 (4H, m), 1.21 (6H, t, J=7.10Hz).

STEP 5. [6-Chloro-2-(2,4-dichlorobenzoyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 1 (method B) from diethyl 2-[6-chloro-2-(2,4-dichlorobenzoyl)-1H-indol-3-yl]malonate (step 4).

m.p.: 190-192 °C.

IR (KBr) ν : 3304, 1709, 1630, 1589, 1526, 1427, 1321, 1231, 1150, 1061 cm⁻¹.

¹H-NMR (DMSO-d₆) δ : 11.66 (1H, s), 7.72 (1H, d, J=2.00Hz), 7.64 (1H, d, 8.91Hz), 7.51-7.33 (3H, m), 7.01 (1H, dd, J=2.00, 8.91Hz), 3.61 (2H, s).

EXAMPLE 203

METHYL [6-CHLORO-2-(4-CHLORO-3-FLUOROBENZOYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 4-chloro-3-fluorophenacyl bromide* .

¹H-NMR (CDCl₃) δ : 8.87 (1H, br s), 7.65-7.36 (4H, m), 7.17 (1H, dd, J=8.72Hz, 1.81Hz), 3.80 (2H, s), 3.68 (3H, s).

4-chloro-3-fluorophenacylbromide was prepared as follows;

4-Acetyl-1-chloro-2-fluorobenzene:

To a solution of 1-bromo-4-chloro-3-fluorobenzene (2.09g, 10mmol) in dry diethyl ether (12.0 ml) was added a solution of n-BuLi (1.55M in hexane, 6.77ml, 10.5mmol) at -78 °C under nitrogen atmosphere. The mixture was allowed to warm to -20 °C stirred for 45 min. A solution of *N,N*-dimethylacetamide (1.04 ml, 11.2 mmol) in diethyl ether (1.5 ml) was added and the mixture stirred for an additional 1h. The mixture was then allowed to warm to room temperature. After stirring for 3 h, the reaction mixture was poured into saturated aqueous ammonium chloride (20 ml) and extracted with diethyl ether (30 ml x 3). The combined organic layers were washed with 2N aqueous HCl (20 ml), saturated aqueous sodium bicarbonate (20 ml), brine (20 ml), and dried (MgSO₄). Removal of solvent gave the title compound as a yellow oil (quant.).

¹H-NMR (CDCl₃) δ: 7.75-7.09 (3H, m), 2.59 (3H, s).

4-Chloro-3-fluorophenacyl bromide:

The title compound was prepared from 4-acetyl-1-chloro-2-fluorobenzene according to the procedure for preparing 4-(trifluoromethoxy)phenacyl bromide described in Example 189.

¹H-NMR (CDCl₃) δ: 7.79-7.71 (2H, m), 7.58-7.52 (1H, m), 4.38 (2H, s).

EXAMPLE 204

[6-CHLORO-2-(4-CHLORO-3-FLUOROBENZOYL)-1H-INDOL-3-YL]ACETIC

ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-(4-methylsulfonylaminophenacyl)-1H-indol-3-yl]acetate (Example 203).

m.p.: 179-182 °C.

MS (EI) m/z: 365 (M⁺).

¹H-NMR (DMSO-d₆) δ: 11.80 (1H, br s), 7.85-7.48 (5H, m), 7.14 (1H, dd, J=8.72Hz, 1.97Hz), 3.84 (2H, s).

EXAMPLE 205

METHYL [6-CHLORO-2-(4-CYANOBENZOYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 8 (Method B) from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate

(step 1 of Example 8, Method A) and 4-cyanophenacyl bromide.

¹H-NMR (CDCl₃) δ: 8.88 (1H, br s), 7.89-7.80 (4H, m), 7.58 (1H, d, J=8.75Hz), 7.41 (1H, d, 1.65Hz), 7.18 (1H, dd, 8.75Hz, 1.65Hz), 3.74 (2H, s), 3.36 (3H, s).

EXAMPLE 206

5 METHYL [6-CHLORO-2-[4-(BROMO)BENZOYL]-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans* 4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 4-bromophenacyl bromide.

¹H-NMR (CDCl₃) δ: 8.89 (1H, br s), 7.66 (4H, s), 7.55 (1H, d, J=8.6Hz), 7.36 (1H, d, J=1.6Hz), 7.15 (1H, dd, J=8.6,1.6Hz), 3.81 (2H, s), 3.66 (3H, s).

EXAMPLE 207

METHYL [6-CHLORO-2-[4-(2-THIENYL)BENZOYL]-1H-INDOL-3-YL]ACETATE

A mixture of methyl [6-chloro-2-(4-bromobenzoyl)-1H-indol-3-yl]acetate
15 (Example 206, 0.40 g, 0.98 mmol), thiophene-2-boronic acid (0.14g, 1.08mmol), saturated aqueous sodium bicarbonate (4 ml), and dichlorobis(triphenylphosphine)palladium(II) (70 mg, 0.098 mmol) in DME (15 ml) was refluxed for 3h. The mixture was poured into water (30 ml), and extracted with ethyl acetate (50 ml x2). The combined extracts were washed with brine (50 ml),
20 dried (MgSO₄), and concentrated. The residue was purified by flash column chromatography eluting with ethyl acetate-hexane (1:6) to afford 0.33g (83 %) of the title compound as yellow solids.

¹H-NMR (CDCl₃) δ: 8.88 (1H, br s), 7.83 (2H, d, J=8.2Hz), 7.75 (2H, d, J=8.2Hz), 7.58 (1H, d, J=8.6Hz), 7.46 (1H, dd, J=3.6,1.2Hz), 7.42-7.38 (2H, m), 7.19-7.12 (2H, m), 3.87 (2H, s), 3.67 (3H, s).

EXAMPLE 208

[6-CHLORO-2-[4-(2-THIENYL)BENZOYL]-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-[4-(2-thienyl)benzoyl]-1H-indol-3-yl]acetate (Example 207).

m.p.: 246-249 °C.

IR (KBr) ν : 3319, 3068, 2628, 1705, 1610, 1593, 1427, 1323, 1257, 1186, 941 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.79 (1H, s), 7.87 (2H, d, $J=8.6\text{Hz}$), 7.82 (2H, d, $J=8.6\text{Hz}$), 7.77-7.68 (3H, m), 7.48 (1H, d, $J=1.5\text{Hz}$), 7.26-7.20 (1H, m), 7.13 (1H, dd, $J=8.6, 2.0\text{Hz}$), 3.86 (2H, s).

5 **EXAMPLE 209**

METHYL [6-CHLORO-2-[4-(2-FURYL)BENZOYL]-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 207 from methyl [6-chloro-2-(4-bromobenzoyl)-1H-indol-3-yl]acetate (Example 206) and furan-2-boronic acid.

10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.90 (1H, br s), 7.83 (2H, d, $J=8.7\text{Hz}$), 7.80 (2H, d, $J=8.7\text{Hz}$), 7.60-7.54 (2H, m), 7.40 (1H, d, $J=1.8\text{Hz}$), 7.15 (1H, dd, $J=8.6, 1.8\text{Hz}$), 6.83 (1H, d, $J=3.5\text{Hz}$), 6.57-6.52 (1H, m), 3.86 (2H, s), 3.66 (3H, s).

EXAMPLE 210

[6-CHLORO-2-[4-(2-FURYL)BENZOYL]-1H-INDOL-3-YL]ACETIC ACID

15 The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetate (Example 209).

m.p.: 230-232 $^{\circ}\text{C}$.

IR (KBr) ν : 3315, 2873, 2630, 1709, 1616, 1597, 1527, 1431, 1321, 1257, 1232 cm^{-1} .

20 $^1\text{H-NMR}$ (DMSO-d_6) δ : 11.80 (1H, s), 7.93-7.79 (3H, m), 7.83 (2H, d, $J=8.1\text{Hz}$), 7.73 (1H, d, $J=8.6\text{Hz}$), 7.48 (1H, d, $J=1.8\text{Hz}$), 7.21 (1H, d, $J=3.5\text{Hz}$), 7.14 (1H, dd, $J=8.6, 1.8\text{Hz}$), 6.72-6.66 (1H, m), 3.84 (2H, s).

EXAMPLE 211

METHYL [6-CHLORO-2-[4-(3-PYRIDYL)BENZOYL]-1H-INDOL-3-YL]ACETATE

25 The title compound was prepared according to the procedure described in Example 207 from methyl [6-chloro-2-(4-bromobenzoyl)-1H-indol-3-yl]acetate (Example 206) and pyridine-3-boronic acid.

30 $^1\text{H-NMR}$ (CDCl_3) δ : 8.95 (1H, br s), 8.90 (1H, d, $J=2.5\text{Hz}$), 8.69-8.65 (1H, m), 7.98-7.92 (1H, m), 7.92 (2H, d, $J=8.6\text{Hz}$), 7.73 (2H, d, $J=8.6\text{Hz}$), 7.59 (1H, d, $J=8.7\text{Hz}$), 7.48-7.40 (2H, m), 7.17 (1H, dd, $J=8.7, 1.8\text{Hz}$), 3.88 (2H, s), 3.67 (3H, s).

EXAMPLE 212**[6-CHLORO-2-[4-(3-PYRIDYL)BENZOYL]-1H-INDOL-3-YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-[4-(3-pyridyl)benzoyl]-1H-indol-3-yl]acetate (Example 211).

m.p.: 194.8 °C.

IR (KBr) ν : 3224, 2960, 1604, 1568, 1527, 1382, 1321, 1259, 1004, 920 cm^{-1} .

$^1\text{H-NMR}$ (CD_3OD) δ : 8.93-8.89 (1H, m), 8.57 (1H, dd, $J=4.8, 1.5\text{Hz}$), 8.24-8.17 (1H, m), 7.98 (2H, d, $J=8.6\text{Hz}$), 7.85 (2H, d, $J=8.6\text{Hz}$), 7.68 (1H, d, $J=8.6\text{Hz}$), 7.60-7.53 (1H, m), 7.46-7.42 (1H, m), 7.04 (1H, dd, $J=8.6, 1.8\text{Hz}$), 3.67 (2H, s).

EXAMPLE 213**METHYL [6-CHLORO-2-[4-(2-THIAZOLYL)BENZOYL]-1H-INDOL-3-YL]ACETATE**

To a stirred solution of thiazole (0.11 g, 1.23 mmol) in diethyl ether (4 ml) was added n-BuLi (1.55M in hexane, 0.79ml) at -78°C under nitrogen atmosphere. After stirring for 30 min., zinc chloride (1.0M in diethyl ether, 3.7 ml, 3.7 mmol) was added, and the mixture stirred at 0°C for 30 min. To the resulting mixture was added palladium catalyst prepared in THF (5 ml) by the treatment of a suspension of dichlorobis(triphenylphosphine)palladium(II) (0.22 g, 0.31mmol) with n-BuLi (1.55M in hexane, 0.39 ml). Methyl [6-chloro-2-(4-bromobenzoyl)-1H-indol-3-yl]acetate (Example 206, 0.25 g, 0.61 mmol) was added to the mixture. The reaction mixture was heated at reflux temperature for 4h., poured into water (50 ml), and extracted with ethyl acetate (50 ml). The combined extracts were washed with brine (50 ml), dried (MgSO_4), and concentrated. This crude product was purified by flash column chromatography eluting with ethyl acetate-hexane (1:3) to afford 0.18g (72%) of the title compound as yellow solids.

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.87 (1H, br s), 8.16 (2H, d, $J=8.4\text{Hz}$), 8.04 (1H, d, $J=3.1\text{Hz}$), 7.93 (1H, d, $J=3.1\text{Hz}$), 7.86 (2H, d, $J=8.4\text{Hz}$), 7.77 (1H, d, $J=8.6\text{Hz}$), 7.49 (1H, d, $J=1.6\text{Hz}$), 7.15 (1H, dd, $J=8.6, 1.6\text{Hz}$), 3.97 (2H, s), 3.54 (3H, s).

EXAMPLE 214**[6-CHLORO-2-[4-(2-THIAZOLYL)BENZOYL]-1H-INDOL-3-YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-chloro-2-[4-(2-thiazolyl)benzoyl]-1H-indol-3-yl]acetate (Example 213).

m.p.: 230-233 °C.

5 IR (KBr) ν : 3331, 3126, 2546, 1693, 1635, 1535, 1350, 1313, 1213, 1150, 912 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.82 (1H, s), 8.15 (2H, d, $J=8.6\text{Hz}$), 8.04 (1H, d, $J=3.3\text{Hz}$), 7.93 (1H, d, $J=3.3\text{Hz}$), 7.88 (2H, d, $J=8.6\text{Hz}$), 7.75 (1H, d, $J=8.7\text{Hz}$), 7.49 (1H, d, $J=2.0\text{Hz}$), 7.14 (1H, dd, $J=8.7, 2.0\text{Hz}$), 3.86 (2H, s).

EXAMPLE 215

10 METHYL [6-CHLORO-2-(3-BROMOBENZOYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans* 4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 3-bromophenacyl bromide.

$^1\text{H-NMR}$ (CDCl_3) δ : 8.91 (1H, br s), 7.93-7.87 (1H, m), 7.77-7.67 (2H, m), 7.57 (1H, d, $J=8.7\text{Hz}$), 7.43-7.35 (2H, m), 7.19-7.13 (1H, m), 3.78 (2H, s), 3.69 (3H, s).

EXAMPLE 216

METHYL [6-CHLORO-2-[3-(2-FURYL)BENZOYL]-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 207 from methyl [6-chloro-2-(3-bromobenzoyl)-1H-indol-3-yl]acetate (Example 215) and furan-2-boronic acid.

$^1\text{H-NMR}$ (CDCl_3) δ : 8.93 (1H, br s), 8.05-8.01 (1H, m), 7.94-7.88 (1H, m), 7.68-7.62 (1H, m), 7.60-7.47 (3H, m), 7.45-7.41 (1H, m), 7.16 (1H, dd, $J=8.6, 1.6\text{Hz}$), 6.74 (1H, d, $J=3.3\text{Hz}$), 6.52-6.47 (1H, m), 3.80 (2H, s), 3.57 (3H, s).

EXAMPLE 217

25 [6-CHLORO-2-[3-(2-FURYL)BENZOYL]-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described Example 9 (Method B) from methyl [6-chloro-2-[3-(2-furyl)benzoyl]-1H-indol-3-yl]acetate (Example 216).

m.p.: 246-249 °C.

30 IR (KBr) ν : 3310, 2984, 2632, 1695, 1624, 1568, 1327, 1227, 1062, 806, 739 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.81 (1H, s), 8.05-7.96 (2H, m), 7.82-7.77 (1H, m), 7.74 (1H,

d, J=8.7Hz), 7.68-7.61 (2H, m), 7.49 (1H, d, J=2.0Hz), 7.14 (1H, dd, J=8.7, 2.0Hz), 7.09 (1H, d, J=3.3Hz), 6.66-6.61 (1H, m), 3.83 (2H, s).

EXAMPLE 218

METHYL *dl*-2-[6-CHLORO-2-(4-CHLOROBENZOYL)-1H-INDOL-3-
YL]PROPIONATE

STEP 1. Methyl [1-*tert*-butoxycarbonyl-2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-
yl]acetate

To a stirred suspension of methyl [2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetate (Example 8, Method B, 2.0 g, 5.5 mmol) in dichloromethane (20 ml) was
10 added di-*tert*-butyl dicarbonate (2.4 g, 11mmol) and 4-dimethylaminopyridine (670 mg, 5.5 mmol) at room temperature. After stirring for 5 min, the mixture was poured into 10% citric acid (200 ml) and extracted with dichloromethane (200 ml). The extract was washed with water (200 ml), brine (200 ml), dried (MgSO₄), and concentrated. The residue was purified by flash column chromatography eluting with
15 ethyl acetate-hexane (1:4) to afford 2.3 g (90%) of the title compound as a yellow solids.

¹H-NMR (CDCl₃) δ: 8.26 (1H, d, J=2.2Hz), 7.74 (2H, d, J=8.6Hz), 7.53 (1H, d, J=8.6Hz), 7.44 (2H, d, J=8.6Hz), 7.32 (1H, dd, J=1.6, 8.4Hz), 3.72 (2H, s), 3.56 (3H, s), 1.30 (9H, s).

20 STEP 2. Methyl [1-*tert*-butoxycarbonyl-2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-
yl]propionate

To a stirred solution of methyl [1-*tert*-butoxycarbonyl-2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetate (step 1, 200 mg, 0.43 mmol) in THF (3 ml) was added a solution of lithium bis(trimethylsilyl)amide in THF (1 M, 0.5 ml) at -78 °C. After
25 stirring for 0.5 h, iodomethane (0.14 ml, 2.2 mmol) was added at that temperature. The mixture was allowed to warm to -10 °C and stirred for an additional 0.5 h. The resulting mixture was poured into saturated aqueous ammonium chloride (50 ml) and extracted with diethyl ether (50 ml). The extract was washed with water (50 ml), brine (50 ml), dried (MgSO₄) and concentrated. The residue was purified by TLC
30 developing with ethyl acetate-hexane (1:4) to afford 146 mg (71 %) of the title compound as a colorless oil.

¹H-NMR (CDCl₃) δ: 8.28 (1H, d, J=1.9Hz), 7.75 (2H, d, J=8.4Hz), 7.59 (1H, d, J=8.4Hz), 7.44 (2H, d, J=8.6Hz), 7.28 (1H, dd, J=1.9, 8.4Hz), 3.86 (1H, q, J=7.3Hz), 3.54 (3H, s), 1.54 (3H, d, J=7.3Hz), 1.30 (9H, s).

STEP 3. Methyl *dl*-[2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]propionate

5 Methyl [1-*tert*-butoxycarbonyl-2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]propionate (step 2, 270 mg, 0.72 mmol) was dissolved in trifluoroacetic acid (5 ml). After stirring for 10 min, the mixture was concentrated. To the residue was added saturated sodium bicarbonate (30 ml), and then the mixture was extracted with ethyl acetate (100 ml). The extract was washed with water (50 ml) brine (50 ml), dried
10 (MgSO₄), and concentrated. The residue was purified by flash column chromatography eluting with ethyl acetate-hexane (1:3) to afford 210 mg (78%) of yellow solids.

¹H-NMR (CDCl₃) δ: 8.89 (1H, br), 7.78 (2H, d, J=8.7Hz), 7.67 (1H, d, J=8.7Hz), 7.50 (2H, d, J=8.6Hz), 7.40-7.37 (1H, m), 7.11 (1H, dd, J=1.8, 8.7Hz), 4.20 (1H, q, J=7.2Hz), 3.64 (3H, s), 1.54 (3H, d, J=7.2Hz).
15

EXAMPLE 219

dl-2-[2-(4-CHLOROBENZOYL)-6-CHLORO-1H-INDOL-3-YL]PROPIONIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl *dl*-[2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]propionate (Example 218).
20

¹H-NMR (CDCl₃) δ: 8.77 (1H, br s), 7.82 (2H, d, J=8.6Hz), 7.75 (1H, d, J=8.7Hz), 7.53 (2H, d, J=8.6Hz), 7.40 (1H, br), 7.13 (1H, dd, J=1.8, 8.7Hz), 4.29 (1H, q, J=7.2Hz), 1.61 (3H, d, J=7.2Hz).

EXAMPLE 220

25 METHYL [5-CHLORO-2-(ISOQUINOLINE-3-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 3-bromoacetylisoquinoline hydrobromide*.

30 ¹H-NMR (CDCl₃) δ: 12.72 (1H, br s), 9.37 (1H, s), 8.80 (1H, s), 8.15-8.02 (2H, m), 7.87-7.76 (2H, m), 7.69 (1H, d, J=2.0Hz), 7.48 (1H, d, J=8.9Hz), 7.32 (1H, dd, J=2.0,

8.7Hz), 4.33 (2H, s), 3.76 (3H, s).

* 3-Bromoacetylisoquinoline hydrobromide was prepared from 3-acetylisoquinoline (D. L. Klayman et al., *Arzneim. Forsch.*, 1986, 36, 10) according to the method of H.McKennis, Jr., L.B.Turnbull, E.R.Bowman, and E.Tamaki (in *J.Org.Chem.*, 1963, 5 28, 383-387).

¹H-NMR (DMSO-d₆) δ: 9.49 (1H, s), 8.68 (1H, s), 8.34-8.26 (2H, m), 7.99-7.88 (2H, m), 5.14 (2H, s).

EXAMPLE 221

[5-CHLORO-2-(ISOQUINOLINE-3-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

10 The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate (Example 219).

MS (EI) m/z: 364 (M⁺).

m.p.: 239-240 °C.

15 IR (KBr) ν: 3277, 1699, 1641, 1531, 1329, 1202, 1059, 961, 787 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.46 (1H, br s), 9.56 (1H, s), 8.69 (1H, s), 8.39-8.26 (2H, m), 7.99-7.90 (2H, m), 7.87 (1H, d, J=2.1Hz), 7.72 (1H, d, J=8.7Hz), 7.35 (1H, dd, J=2.1, 8.9Hz), 4.13 (2H, s)..

EXAMPLE 222

20 METHYL [6-CHLORO-2-(ISOQUINOLINE-3-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 3-bromoacetylisoquinoline (Preparation is described in 25 Example 220).

¹H-NMR (CDCl₃) δ: 12.69 (1H, br s), 9.38 (1H, s), 8.81 (1H, s), 8.16-8.04 (2H, m), 7.88-7.77 (2H, m), 7.65 (1H, d, J=8.9Hz), 7.57 (1H, d, J=2.0Hz), 7.15 (1H, dd, J=2.0 and 8.7Hz), 4.36 (2H, s), 3.75 (3H, s).

EXAMPLE 223

30 [6-CHLORO-2-(ISOQUINOLINE-3-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate (Example 222).

MS (EI) m/z: 364 (M^+).

5 m.p.: 236-237 °C.

IR (KBr) ν : 3229, 1709, 1641, 1618, 1531, 1198, 793 cm^{-1} .

$^1\text{H-NMR}$ (DMSO- d_6) δ : 12.43 (1H, br s), 9.56 (1H, s), 8.70 (1H, s), 8.41-8.26 (2H, m), 7.99-7.87 (2H, m), 7.82 (1H, d, $J=8.7\text{Hz}$), 7.77 (1H, d, $J=1.5\text{Hz}$), 7.14 (1H, dd, $J=2.0, 8.6\text{Hz}$), 4.14 (2H, s).

10 **EXAMPLE 224**

METHYL [5-CHLORO-2-(5-METHYLISOXAZOLE-3-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 15 36, step 3) and 3-bromoacetyl-5-methylisoxazole (M. D. Amici et al., *J. Org. Chem.*, 1989, 54, 2646).

$^1\text{H-NMR}$ (CDCl_3) δ : 10.75 (1H, br s), 7.68 (1H, br s), 7.42 (1H, d, $J=8.2\text{Hz}$), 7.34 (1H, dd, $J=1.8, 9.1\text{Hz}$), 6.60 (1H, s), 4.25 (2H, s), 3.74 (3H, s), 2.56 (3H, s).

EXAMPLE 225

20 **[5-CHLORO-2-(5-METHYLISOXAZOLE-3-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetate (Example 224).

25 MS (EI) m/z: 318 (M^+).

m.p.: 253-255 °C.

IR (KBr) ν : 3379, 1699, 1639, 1539, 1425, 1259, 1207, 1059, 804 cm^{-1} .

$^1\text{H-NMR}$ (DMSO- d_6) δ : 11.76 (1H, br s), 7.81 (1H, d, $J=2.0\text{Hz}$), 7.59 (1H, d, $J=8.9\text{Hz}$), 7.29 (1H, dd, $J=2.0, 8.7\text{Hz}$), 6.67 (1H, s), 4.00 (2H, s), 2.49 (3H, s).

30 **EXAMPLE 226**

METHYL [6-CHLORO-2-(5-METHYLISOXAZOLE-3-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 3-bromoacetyl-5-methylisoxazole (M. D. Amici et al., *J. Org. Chem.*, 1989, 54, 2646).

¹H-NMR (CDCl₃) δ: 10.70 (1H, br s), 7.62 (1H, d, J=8.7Hz), 7.49 (1H, d, J=1.8Hz), 7.15 (1H, dd, J=1.8 and 8.7Hz), 6.60 (1H, br s), 4.27 (2H, s), 3.72 (3H, s), 2.56 (3H, s).

EXAMPLE 227

[6-CHLORO-2-(5-METHYLISOXAZOLE-3-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetate (Example 226).

MS (EI) m/z: 318 (M⁺).

m.p.: 227-229°C.

IR (KBr) ν: 3331, 1713, 1645, 1543, 1404, 1259, 1202, 891, 804 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.70 (1H, br s), 7.74 (1H, d, J=8.7Hz), 7.62 (1H, d, J=1.3Hz), 7.07 (1H, dd, J=1.8, 8.6Hz), 6.67 (1H, br s), 4.01 (2H, s), 2.49 (3H, s).

EXAMPLE 228

METHYL [5-CHLORO-2-(4-METHYL-1,2,3-THIADIAZOLE-5-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 5-bromoacetyl-4-methyl-1,2,3-thiadiazole hydrobromide*.

¹H-NMR (CDCl₃) δ: 9.00 (1H, br s), 7.63 (1H, br s), 7.34 (1H, dd, J=1.8, 8.9Hz), 7.28 (1H, d, J=8.9Hz), 3.81 (2H, s), 3.66 (3H, s), 2.78 (3H, s).

* 5-Bromoacetyl-4-methyl-1,2,3-thiadiazole hydrobromide was prepared as follows;

N-Methoxy-*N*-methyl-4-methyl-1,2,3-thiadiazole-5-carboxamide:

To a solution of 1,2,3-thiadiazole-5-carbonyl chloride (10.00 g, 61.5 mmol) and *N,O*-dimethylhydroxylamine hydrochloride (7.20 g, 73.8 mmol) in

dichloromethane (200 ml) was added triethylamine (20.6 ml, 147.6 mmol) at 0°C. After stirring for 2 h at room temperature, the mixture was diluted with dichloromethane (300 ml), washed with water (200 ml x 2) and dried (MgSO₄). Removal of solvent gave 11.31 g (98%) of the title compound as brown crystals.

5 ¹H-NMR (CDCl₃) δ: 3.75 (3H, s), 3.40 (3H, s), 3.01 (3H, s).

5-Acetyl-4-methyl-1,2,3-thiadiazole:

To a solution of *N*-methoxy-*N*-methyl-4-methyl-1,2,3-thiadiazole-5-carboxamide (11.31 g, 60.4 mmol) in THF (100 ml) was added 2M diethyl ether solution of methylmagnesium iodide (45.3 ml, 90.6 mmol) over 0.5 h at 0°C. The mixture was allowed to warm to room temperature and stirred for 3 h.

The mixture was quenched with saturated aqueous ammonium chloride (100 ml) and then extracted with diethyl ether (200 ml x 2). The extracts were dried (MgSO₄) and concentrated to give 7.49 g (87%) of the title compound as a pale brown oil.

¹H-NMR (CDCl₃) δ: 2.96 (3H, s), 2.67 (3H, s).

15 5-Bromoacetyl-4-methyl-1,2,3-thiadiazole:

To a solution of 5-acetyl-4-methyl-1,2,3-thiadiazole (1.00 g, 7.03 mmol) in chloroform (20 ml) was added dropwise a solution of bromine (1.24 g, 7.73 mmol) in chloroform (10 ml) over 0.5 h at room temperature. The mixture was heated at reflux temperature for 2 h. After cooling to room temperature, the mixture was made basic with saturated aqueous sodium bicarbonate and extracted with dichloromethane (100 ml x 2). The extracts were dried (Na₂SO₄) and concentrated to give 1.55 g (100%) of the title compound as a brown oil.

¹H-NMR (CDCl₃) δ: 4.28 (2H, s), 2.98 (3H, s).

EXAMPLE 229

25 **[5-CHLORO-2-(4-METHYL-1,2,3-THIADIAZOLE-5-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID**

A mixture of methyl [5-chloro-2-(4-methyl-1,2,3-thiadiazole-5-carbonyl)-1H-indol-3-yl]acetate (Example 228, 190 mg, 0.54 mmol), 2N aqueous HCl (4 ml), and acetic acid (20 ml) was heated at reflux temperature for 2 h. After cooling to room temperature, the mixture was concentrated. The crystalline residue was diluted with THF (100 ml) and dried (MgSO₄) and concentrated. The residual solids were washed

with ethyl acetate gave 145 mg (79%) of the title compound as yellow solids.

MS (EI) m/z: 335 (M^+).

m.p.: 229-230 °C.

IR (KBr) ν : 3300, 1715, 1622, 1526, 1329, 1261, 1204, 1063, 1009, 822 cm^{-1} .

- 5 $^1\text{H-NMR}$ (DMSO-d_6) δ : 11.92 (1H, br s), 7.89 (1H, d, $J=2.0\text{Hz}$), 7.48 (1H, d, $J=8.9\text{Hz}$), 7.36 (1H, dd, $J=2.0, 8.9\text{Hz}$), 3.95 (2H, s), 2.63 (3H, s).

EXAMPLE 230

METHYL [6-CHLORO-2-(4-METHYL-1,2,3-THIADIAZOLE-5-CARBONYL)-1H-INDOL-3-YL]ACETATE

- 10 The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 5-bromoacetyl-4-methyl-1,2,3-thiadiazole (Preparation is described in Example 228).

tlc: $R_f=0.56$ (ethyl acetate/hexane=1:2)

- 15 **EXAMPLE 231**

[6-CHLORO-2-(4-METHYL-1,2,3-THIADIAZOLE-5-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

- 20 The title compound was prepared according to the procedure described in Example 229 from methyl [6-chloro-2-(4-methyl-1,2,3-thiadiazole-5-carbonyl)-1H-indol-3-yl]acetate (Example 230).

MS (EI) m/z: 335 (M^+).

m.p.: 215-216°C.

IR (KBr) ν : 3300, 1709, 1645, 1531, 1327, 1211, 1065, 922, 789 cm^{-1} .

- 25 $^1\text{H-NMR}$ (DMSO-d_6) δ : 12.35 (1H, br s), 11.86 (1H, br s), 7.82 (1H, d, $J=8.7\text{Hz}$), 7.47 (1H, d, $J=1.8\text{Hz}$), 7.17 (1H, dd, $J=1.8, 8.6\text{Hz}$), 3.93 (2H, s), 2.63 (3H, s).

EXAMPLE 232

METHYL [5-CHLORO-2-(5-METHYLTHIAZOLE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

- 30 The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetyl-5-methylthiazole hydrobromide*.

¹H-NMR (CDCl₃) δ: 11.74 (1H, br s), 7.76 (1H, d, J=1.1Hz), 7.67 (1H, d, J=1.8), 7.43 (1H, d, J=8.7Hz), 7.32 (1H, dd, J=2.0, 8.7Hz), 4.28 (2H, s), 3.73 (3H, s), 2.62 (3H, d, J=1.0Hz).

* 2-Bromoacetyl-5-methylthiazole was prepared from 2-acetyl-5-methylthiazole (Metzger et al., *Bull. Soc. Chim. Fr.*, **1953**, 702) according to the method of H.McKennis, Jr., L.B.Turnbull, E.R.Bowman, and E.Tamaki (in *J.Org.Chem.*, **1963**, 28, 383-387).

¹H-NMR (DMSO-d₆) δ: 7.91 (1H, d, J=1.2Hz), 4.87 (2H, s), 2.58 (3H, d, J=0.8Hz).

EXAMPLE 233

10 [5-CHLORO-2-(5-METHYLTHIAZOLE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate (Example 232).

15 MS (EI) m/z: 334 (M⁺).

m.p.: 231-233°C.

IR (KBr) v: 3348, 1699, 1630, 1541, 1404, 1333, 1271, 1057, 1003, 804 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.05 (1H, br s), 8.04 (1H, s), 7.88 (1H, br s), 7.77 (1H, d, J=8.7Hz), 7.35 (1H, br d, J=8.9Hz), 4.15 (2H, s), 2.63 (3H, s).

20 EXAMPLE 234

METHYL [6-CHLORO-2-(5-METHYLTHIAZOLE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of **25** Example 8, Method A) and 2-bromoacetyl-5-methylthiazole (Preparation is described in Example 232).

¹H-NMR (CDCl₃) δ: 11.73 (1H, br s), 7.77 (1H, s), 7.62 (1H, d, J=8.2Hz), 7.51 (1H, br s), 7.14 (1H, br d, J=8.7Hz), 4.31 (2H, s), 3.72 (3H, s), 2.62 (3H, s).

EXAMPLE 235

30 [6-CHLORO-2-(5-METHYLTHIAZOLE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate (Example 234).

MS (EI) m/z: 334 (M^+).

5 m.p.: 225-226°C.

IR (KBr) ν : 3277, 1707, 1630, 1541, 1398, 1350, 1231, 1219, 1138, 878, 800 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.25 (1H, br s), 11.99 (1H, br s), 8.04 (1H, s), 7.82 (1H, s), 7.81 (1H, d, $J=9.0\text{Hz}$), 7.13 (1H, br d, $J=8.7\text{Hz}$), 4.15 (2H, s), 2.63 (3H, s).

EXAMPLE 236

10 **[6-CHLORO-2-(2-THIENYL)CARBONYLINDOL-3-YL]ACETIC ACID**

STEP 1. 6-Chloro-2-(2-thienylcarbonyl)-1-(phenylsulfonyl)indole

The title compound was prepared according to the procedure described in step 2 of Example 2 (Method B) from 6-chloro-1-(phenylsulfonyl)indole (step 1 of Example 2, Method B) and 2-thienoyl chloride.

15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.13-8.16 (3H, m), 7.77-7.80 (2H, m), 7.50-7.63 (4H, m), 7.29 (1H, dd, $J=1.8, 8.4\text{Hz}$), 7.19 (1H, t, $J=5.4\text{Hz}$), 7.03 (1H, s).

STEP 2. 6-Chloro-2-(2-thienylcarbonyl)indole

20 The title compound was prepared according to the procedure described in step 3 of Example 2 (Method B) from 6-chloro-2-(2-thienylcarbonyl)-1-(phenylsulfonyl)indole (step 1).

$^1\text{H-NMR}$ (CDCl_3) δ : 10.96 (1H, br s), 8.01 (1H, dd, $J=1.2, 3.8\text{Hz}$), 7.73 (1H, dd, $J=1.2, 4.9\text{Hz}$), 7.64 (1H, d, $J=8.6\text{Hz}$), 7.54-7.55 (1H, m), 7.35-7.37 (1H, m), 7.23 (1H, dd, $J=3.8, 4.9\text{Hz}$), 7.10 (1H, dd, $J=1.8, 8.6\text{Hz}$).

STEP 3. Diethyl α -acetoxo-[6-chloro-2-(2-thienylcarbonyl)-1H-indol-3-yl]malonate

25 The title compound was prepared according to the procedure described in step 4 of Example 2 (Method B) from 6-chloro-2-(2-thienylcarbonyl)indole (step 2).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.93 (1H, br s), 7.82 (1H, d, $J=8.9\text{Hz}$), 7.76 (1H, d, $J=4.9\text{Hz}$), 7.55 (1H, d, $J=3.8\text{Hz}$), 7.39 (1H, d, $J=1.8\text{Hz}$), 7.11-7.18 (2H, m), 4.16-4.31 (4H, m), 1.87 (3H, s), 1.17-1.32 (6H, m).

30 **STEP 4. Diethyl [6-chloro-2-(2-thienylcarbonyl)-1H-indol-3-yl]malonate**

The title compound was prepared according to the procedure described in step 5 of Example 2 (Method B) from diethyl α -acetoxyl[6-chloro-2-(2-thienylcarbonyl)-1H-indol-3-yl]malonate (step 3).

$^1\text{H-NMR}$ (CDCl_3) δ : 9.27 (1H, br s), 7.71-7.75 (2H, m), 7.60-7.66 (1H, m), 7.05-7.17 (3H, m), 5.56 (1H, s), 4.07-4.26 (4H, m), 1.20-1.28 (6H, m).

STEP 5. [6-Chloro-2-(2-thienylcarbonyl)-1H-indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 6 of Example 2 (Method B) from diethyl [6-chloro-2-(2-thienylcarbonyl)-1H-indol-3-yl]malonate (step 4).

MS (EI) m/z : 319 (M^+).

m.p.: 177-178 °C.

IR (KBr) ν : 3323, 1701, 1593, 1568, 1524, 1435, 1412, 1323, 1258, 1229, 920 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.25 (1H, br s), 11.85 (1H, br s), 8.13 (1H, d, $J=4.9\text{Hz}$), 7.89 (1H, d, $J=3.6\text{Hz}$), 7.74 (1H, d, $J=8.7\text{Hz}$), 7.51 (1H, d, $J=1.8\text{Hz}$), 7.34 (1H, t, $J=4.8\text{Hz}$), 7.15 (1H, dd, $J=1.8, 8.7\text{Hz}$), 3.96 (2H, s).

EXAMPLE 237

METHYL [6-CHLORO-2-[3-(1-HYDROXY-1-METHYLETHYL)-2-FUROYL]-1H-INDOL-3-YL]ACETATE

The reaction was carried out according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(penylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 2-chloroacethyl-3-(1-hydroxy-1-methylethyl)furan*.

$^1\text{H-NMR}$ (CDCl_3) δ : 9.89 (1H, br s), 7.59 (1H, d, $J=1.6\text{Hz}$), 7.53 (1H, d, $J=8.7\text{Hz}$), 7.34 (1H, d, $J=1.5\text{Hz}$), 7.08 (1H, dd, $J=1.6, 8.7\text{Hz}$), 6.57 (1H, d, $J=1.8\text{Hz}$), 6.39 (1H, br s), 4.22 (2H, s), 3.73 (3H, s), 1.57 (6H, s).

*2-Chloroacethyl-3-(1-hydroxy-1-methylethyl)furan was prepared as follows;

To a solution of 2-(3-furyl)-2-propanol (T.M.Bargar et al., *J. Med. Chem.*, 1986, 29, 315., 2.0 g, 15.85 mmol) in THF (100 ml) was added a solution of *n*-butyllithium in hexane (1.55M, 30.7 ml, 47.55 mmol) at -78°C. After stirring for 1h, 2-chloro-*N*-methoxy-*N*-methylacetamide (6.54 g, 47.55 mmol) was added at 0°C. Saturated aqueous ammonium chloride (100 ml) was added to the mixture and the organic layer was separated. The organic layer was washed with water (100 ml x 2) and brine (50

ml) and dried (MgSO₄). After removal of the solvent, the residue was purified by flash chromatography eluting with ethyl acetate/hexane (1:4) to afford 1.03 g (32%) of the title compound as an oil.

¹H-NMR (CDCl₃) δ: 7.52 (1H, d, J=1.8Hz), 6.56 (1H, d, J=1.8Hz), 5.51 (1H, s), 4.74 (2H, s), 1.56 (6H, s).

EXAMPLE 238

[6-CHLORO-2-[3-(1-HYDROXY-1-METHYLETHYL)-2-FUROYL]-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-[3-(1-hydroxy-1-methylethyl)-2-furoyl]-1H-indol-3-yl]acetate (Example 237).

MS (EI) m/z: 361 (M⁺).

m.p.: 229-230°C.

IR(KBr)v: 3270, 2980, 1270, 1591, 1564, 1522, 1398, 1302, 1263, 1200, 785 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.78 (1H, br s), 7.97 (1H, d, J=1.6Hz), 7.77 (1H, d, J=8.6Hz), 7.56 (1H, d, J=1.5Hz), 7.13 (1H, dd, J=1.8, 8.6Hz), 6.87 (1H, d, J=1.6Hz), 5.69 (1H, br s), 3.96 (2H, s), 1.53 (6H, s).

EXAMPLE 239

METHYL [6-CHLORO-2-[3-METHOXYMETHYL-2-FUROYL]-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(penylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and 2-chloroacetyl-3-(methoxymethyl)furan*.

¹H-NMR (CDCl₃) δ: 9.81 (1H, br s), 7.60 (1H, d, J=1.6Hz), 7.56 (1H, d, J=8.7Hz), 7.40 (1H, d, J=1.5Hz), 7.10 (1H, dd, J=1.8, 8.7Hz), 6.79 (1H, d, J=1.6Hz), 4.81 (2H, s), 4.26 (2H, s), 3.74 (3H, s), 3.48 (3H, s).

*2-Chloroacetyl-3-(methoxymethyl)furan was prepared from 3-(methoxymethyl)furan (N.Greeves et al., *Synthesis*, 1993, 1109) according to the procedure for preparing 2-chloroacetyl-3-(1-hydroxy-1-methylethyl)furan described in Example 237.

¹H-NMR (CDCl₃) δ: 7.52 (1H, d, J=1.8Hz), 6.74 (1H, d, J=1.8Hz), 4.73 (2H, s), 4.62 (2H, s), 3.44 (3H, s).

EXAMPLE 240

[6-CHLORO-2-[3-METHOXYMETHYL-2-FUROYL]-1H-INDOL-3-YL]ACETIC

ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(3-methoxymethyl-2-furoyl)-1H-indol-3-yl]acetate (Example 239).

MS (EI) m/z: 347 (M⁺).

m.p.: 212-213°C.

IR (KBr) ν: 3373, 3221, 1720, 1601, 1576, 1529, 1205, 1173, 1115, 1088 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.72 (1H, br s), 8.04 (1H, d, J=1.6Hz), 7.76 (1H, d, J=8.7Hz), 7.59 (1H, d, J=1.5Hz), 7.12 (1H, dd, J=1.8, 8.7Hz), 6.86 (1H, d, J=1.6Hz), 4.70 (2H, s), 4.02 (2H, s), 3.36 (3H, s).

EXAMPLE 241

[6-CHLORO-2-(1-METHYLIMIDAZOLE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

STEP 1. Methyl [6-chloro-1-ethoxycarbonyl-2-(1-methylimidazole-2-carbonyl)indolin-3-yl]acetate

The title compound was prepared according to the procedure described in step 2 of Example 31 from methyl *trans*-4-chloro-2-(ethoxycarbonylamino)cinnamate (Example 31, step 1) and 2-bromoacetyl-1-methylimidazole hydrobromide*.

MS (EI) m/z: 405 (M⁺).

*2-bromoacetyl-1-methylimidazole hydrobromid was prepared from 2-acetyl-1-methylimidazole according to the procedure for preparing 2-bromoacetyl-4-methylpyridine hydrobromide described in step 2 of Example 31.

¹H-NMR (DMSO-d₆) δ: 7.69 (1H, s), 7.27 (1H, s), 4.68 (2H, s), 3.81 (3H, s).

STEP 2. [6-chloro-1-ethoxycarbonyl-2-(1-methylimidazole-2-carbonyl)indol-3-yl]acetic acid

The title compound was prepared according to the procedure described in step 3 of Example 31 from methyl [6-chloro-1-ethoxycarbonyl-2-(1-methylimidazole-2-

carbonyl]indolin-3-yl]acetate (step 1).

m.p.: 235.5 °C.

IR (KBr) ν : 3238, 1695, 1630, 1538, 1402, 1229, 1146 cm^{-1} .

$^1\text{H-NMR}$ (CDCl_3) δ : 12.3 (1H, br s), 7.64 (1H, d, $J=8.7\text{Hz}$), 7.50 (1H, d, $J=1.8\text{Hz}$),
5 7.41 (1H, s), 7.16 (1H, s), 7.09 (1H, dd, $J=1.8, 8.6\text{Hz}$), 4.25 (2H, s), 4.13 (3H, s).

EXAMPLE 242

METHYL [6-CHLORO-2-(1-METHYLMIDAZOLE-2-CARBONYL)-1H-INDOL-3- YL]ACETATE

To a stirred solution of [6-chloro-2-(1-methylimidazole-2-carbonyl)indol-3-
10 yl]acetic acid (Example 241, 65 mg, 0.21 mmol) in methanol (10 ml) was added
(trimethylsilyl)diazomethane (1.0 M solution in hexanes, 1.05 ml, 2.1 mmol) at room
temperature. After stirring for 19 h, the mixture was concentrated. The residue was
purified by TLC developing with ethyl acetate-hexane (1:2) to afford 20 mg (23 %) of
the title compound as yellow solids.

15 $^1\text{H-NMR}$ (CDCl_3) δ : 12.35 (1H, br s), 7.59 (1H, d, $J=8.0\text{Hz}$), 7.49 (1H, d, $J=1.5\text{Hz}$),
7.24 (1H, s), 7.11 (1H, dd, $J=1.7, 8.1\text{Hz}$), 7.08 (1H, s), 4.30 (2H, s), 4.12 (3H, s), 3.71
(3H, s).

EXAMPLE 243

METHYL [5-CHLORO-2-(1-METHYLMIDAZOLE-2-CARBONYL)-1H-INDOL-3- 20 YL]ACETATE

The title compound was prepared according to the procedure described in
Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example
36, step 3) and 2-bromoacetyl-1-methylimidazole hydrobromide (Preparation is
described in Example 241).

25 $^1\text{H-NMR}$ (DMSO-d_6) δ : 12.28 (1H, br s), 7.85 (1H, s), 7.77 (1H, d, $J=8.7\text{Hz}$), 7.66 (1H,
s), 7.33-7.29 (2H, m), 4.21 (2H, s), 4.03 (3H, s), 3.60 (3H, s).

EXAMPLE 244

[5-CHLORO-2-(1-METHYLMIDAZOLE-2-CARBONYL)-1H-INDOL-3- YL]ACETIC ACID

30 The title compound was prepared according to the procedure described in
Example 58 from methyl [5-chloro-2-(1-methylimidazol-2-carbonyl)-1H-indol-3-

yl]acetate (Example 243).

m.p.: 230-233°C.

¹H-NMR (DMSO-d₆) d: 12.50 (1H, br s), 7.84 (1H, s), 7.76 (1H, d, J=13.2Hz), 7.66 (1H, s), 7.35-7.7.28 (2H, m).4.15 (2H, s), 4.06 (3H, s).

5 **EXAMPLE 245**

**METHYL [5-CHLORO-2-(IMIDAZOLE-2-CARBONYL)-1H-INDOL-3-
YL]ACETATE**

**STEP 1. Methyl [5-chloro-2-[1-[2-(trimethylsilyl)ethoxymethyl]imidazole-2-carbonyl]-
1H-indol-3-yl]acetate**

10 The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 2-chloroacetyl-1-[2-(trimethylsilyl)ethoxymethyl]imidazole.*

¹H-NMR (CDCl₃) d: 12.38 (1H, br s), 7.66 (1H, s), 7.46-7.38 (2H, m), 7.34-7.29 (2H, m), 5.95 (2H, s), 4.29 (2H, s), 3.73 (3H, s), 3.66 (2H, t, J=8.0Hz), 0.98 (2H, t, J=8.0Hz),
15 0.03 (9H, s).

*2-chloroacetyl-1-[2-(trimethylsilyl)ethoxymethyl]imidazole was prepared as follows;

To a stirred solution of 1-[2-(trimethylsilyl)ethoxymethyl]imidazole (Jeffrey P. Whitten et al., *J. Org. Chem.*, **51**, 1891 (1986), 3.0 g, 15 mmol) in THF (30 ml) was added dropwise n-BuLi (1.55 M in n-hexane, 11.0 ml, 17 mmol) at -78 °C and the
20 mixture was stirred for 1 h. To the resulting mixture was added 2-chloro-*N*-methoxy-*N*-methylacetamide (2.4 g, 17 mmol) at that temperature. The mixture was allowed to warm to 0°C and stirred at for an additional 2 h. The mixture was poured into water (50 ml) and extracted with ethyl acetate (80 ml), dried (MgSO₄) and concentrated. The residual brown oil was purified by flash column chromatography eluting with ethyl
25 acetate/hexane (1:4) to afford 1.2 g (32 %) of the title compound as a yellow oil.

¹H-NMR (CDCl₃) d: 7.39 (1H, s), 7.25 (1H, s), 5.81 (2H, s), 4.96 (2H, s), 3.60 (2H, t, J=8.2Hz), 0.95 (2H, t, J=8.2Hz), 0.02 (9H, s).

STEP 2. Methyl [5-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetate

To a solution of methyl [5-chloro-2-[1-[2-(trimethylsilyl)ethoxymethyl]imidazole-2-carbonyl]-1H-indol-3-yl]acetate (step 1, 300
30 mg, 0.67 mmol) in methanol (10 ml) was added 2N aqueous HCl (7 ml) and the

mixture was refluxed for 1.5 h. After cooling to room temperature, the mixture was concentrated. To the residue was added saturated aqueous sodium bicarbonate (10 ml) and then the mixture was concentrated. The residual yellow solids were dissolved in THF (100 ml) and dried (MgSO₄). Removal of solvent afforded 220 mg (100%) of the title compound as yellow solids.

¹H-NMR (CDCl₃) δ: 12.16 (1H, br s), 10.80 (1H, br s), 7.68 (1H, s), 7.50-7.28 (5H, m), 4.29 (2H, s), 3.70 (3H, s).

EXAMPLE 246

[5-CHLORO-2-(IMIDAZOLE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetate (Example 245).

m.p.: 253-254 °C.

¹H-NMR (DMSO-d₆) δ: 12.28 (1H, br s), 7.84 (1H, d, J=2.0Hz), 7.80 (1H, d, J=9.1Hz), 7.32 (1H, dd, J=2.0, 9.1Hz), 4.17 (2H, s).

EXAMPLE 247

METHYL [6-CHLORO-2-(IMIDAZOLE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

STEP 1. Methyl [6-chloro-2-[1-[2-(trimethylsilyl)ethoxymethyl]imidazole-2-carbonyl]-1H-indol-3-yl]acetate

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (Step 1 of Example 8, Method A) and 2-chloroacetyl-1-[2-(trimethylsilyl)ethoxymethyl]imidazole (Preparation is described in step 1 of Example 245).

¹H-NMR (CDCl₃) δ: 12.34, (1H, br s), 7.61 (1H, d, J=8.6Hz), 7.52 (1H, d, J=1.6Hz), 7.40 (1H, d, J=1.4Hz), 7.31 (1H, d, J=1.4Hz), 7.13 (1H, dd, J=1.6, 8.6Hz), 5.95 (2H, s), 4.32 (2H, s), 3.72 (3H, s), 3.67 (2H, t, J=8.0Hz), 1.00 (2H, t, J=8.0Hz), 0.02 (9H, s).

STEP 2. Methyl [5-Chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetate

The title compound was prepared according to the procedure described in step 2 of Example 245 from methyl [6-chloro-2-[1-[2-(trimethylsilyl)ethoxymethyl]imidazole-2-carbonyl]-1H-indol-3-yl]acetate (step 1).

¹H-NMR (CDCl₃) δ: 12.50 (1H, br s), 7.68-7.48 (2H, m), 7.40-7.30 (1H, m), 7.19-7.06 (1H, m), 6.97-6.92 (1H, m), 4.30 (2H, s), 3.67 (3H, s).

EXAMPLE 248

[6-CHLORO-2-(IMIDAZOLE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

5 The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetate (Example 247).

m.p.: 252-253 °C.

¹H-NMR (DMSO-d₆) δ: 13.63 (1H, br s), 12.40-12.15 (2H, br), 7.88 (1H, d, J=1.8Hz),
10 7.78 (1H, d, J=8.6Hz), 7.59 (1H, s), 7.40 (1H, s), 7.11 (1H dd, J=1.8, 8.6Hz), 4.18 (2H, s).

EXAMPLE 249

METHYL [5-CHLORO-2-(4-METHYLTHIAZOLE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

15 The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 2-bromoacetyl-4-methylthiazole hydrobromide (Cowden, William B. et al., *Aust. J. Chem.*, 1985, 38, 1257).

¹H-NMR (DMSO-d₆) δ: 12.02 (1H, br s), 7.93-7.85 (2H, m), 7.75 (1H, d, J=8.9Hz),
20 7.38 (1H, dd, J=2.0, 8.9Hz), 4.23 (2H, s), 3.60 (3H, s), 2.61 (3H, s).

EXAMPLE 250

[5-CHLORO-2-(4-METHYLTHIAZOLE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

25 The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate (Example 249).

m.p.: 218-220 °C.

¹H-NMR (DMSO-d₆) δ: 11.97 (1H, br s), 7.89 (1H, d, J=2.0Hz), 7.75 (1H, d, J=8.7Hz),
30 7.37 (1H, dd, J=2.0, 8.7Hz), 4.13 (2H, s), 2.62 (3H, s).

EXAMPLE 251

METHYL [5-CHLORO-2-(1-METHYLPYRROLE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 8 (Method B) from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 2-chloroacetyl-1-methylpyrrole (P. D. Croce et al., *Synthesis*, **1990**, 212).

¹H-NMR (DMSO-d₆) δ: 11.8 (1H, br s), 7.80 (1H, br s), 7.55 (1H, d, J=8.6Hz), 7.40-7.32 (2H, m), 6.95-6.90 (1H, m), 6.35-6.26 (1H, m), 4.00 (2H, s), 3.43 (3H, s), 2.50 (3H, s).

10 EXAMPLE 252

[5-CHLORO-2-(1-METHYLPYRROLE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

A mixture of methyl [5-chloro-2-(1-methylpyrrole-2-carbonyl)-1H-indol-3-yl]acetate (Example 251, 250 mg, 0.79 mmol) and potassium carbonate (900 mg, 6.4 mmol) in methanol-water (1:1, 40 ml) was heated at reflux temperature for 1 h. After cooling to room temperature, the mixture was concentrated. The residue was neutralized with 2N aqueous HCl and extracted with ethyl acetate (50 ml x 2). The combined extracts were dried (MgSO₄) and concentrated. The residual solids were recrystallized from ethyl acetate/hexane to afford 40 mg (23%) of the title compound as pale yellow solids.

mp: 203-205°C.

¹H-NMR (DMSO-d₆) δ: 11.56 (1H, br s), 7.57 (1H, s), 7.33 (1H, d, J=8.6Hz), 7.20-7.10 (2H, m), 6.72 (1H, s), 6.10 (1H, s), 3.81 (2H, s), 2.52 (3H, s).

EXAMPLE 253

25 METHYL [5-CHLORO-2-(2-METHYLIMIDAZOLE-4-CARBONYL)-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 8 (Method B) from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 4-bromoacetyl-2-methylimidazole (Deady, Leslie W. et al., *Aust. J. Chem.*, **1981**, 34, 1295).

¹H-NMR (DMSO-d₆) δ: 12.04 (1H br s), 8.51 (1H, s), 7.86 (1H, d, J=2.0Hz), 7.71 (1H, d, J=8.9Hz), 7.34 (1H, dd, J=2.0, 8.9Hz), 4.17 (2H, s), 3.59 (3H, s), 2.71 (3H, s).

EXAMPLE 254

[5-CHLORO-2-(2-METHYLIMIDAZOLE-4-CARBONYL)-1H-INDOL-3-

5 **YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetate (Example 253).

m.p.: 237-238 °C.

10 ¹H-NMR (DMSO-d₆) δ: 12.03 (1H, br), 12.00 (1H, s), 8.49 (1H, s), 7.83 (1H, d, J=2.0Hz), 7.70 (1H, d, J=8.9Hz), 7.34 (1H, dd, J=2.0, 8.9Hz), 4.09 (2H, s), 2.87 (3H, s).

EXAMPLE 255

METHYL [5-CHLORO-2-(THIAZOLE-5-CARBONYL)-1H-INDOL-3-

15 **YL]ACETATE**

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and 5-bromoacetylthiazole hydrobromide*.

20 ¹H-NMR (CDCl₃) δ: 12.05 (1H, br s), 9.51 (1H, s), 8.62 (1H, s), 7.85 (1H, d, J=1.9Hz), 7.55 (1H, d, J=8.7Hz), 7.35 (1H, dd, J=1.9, 8.7Hz), 4.09 (2H, s), 3.59 (3H, s).

*5-bromoacetylthiazole hydrobromide was prepared from 5-acetylthiazole according to the procedure for preparing 2-bromoacetyl-4-methylpyridine hydrobromide described in step 2 of Example 31.

¹H-NMR (DMSO-d₆) δ: 9.49 (1H, s), 8.34 (1H, s), 4.91 (2H, s).

25 **EXAMPLE 256**

[5-CHLORO-2-(THIAZOLE-5-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [5-chloro-2-(thiazole-5-carbonyl)-1H-indol-3-yl]acetate (Example 255).

30 m.p.: 175-180 °C.

¹H-NMR (DMSO-d₆) δ: 11.80 (1H, br s), 9.48 (1H, s), 8.74 (1H, s), 7.78 (1H, s), 7.48 (1H, d, J=8.7Hz), 7.30 (1H, d, J=8.9Hz), 3.81 (2H, s).

EXAMPLE 257

METHYL [6-CHLORO-2-(4-METHYLTHIAZOLE-2-CARBONYL)-1H-INDOL-3-
YL]ACETATE

The title compound was prepared according to the procedure described in Example 8 (Method B) from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (Step 1 of Example 8, Method A) and 2-bromoacetyl-4-methylthiazole hydrobromide (Cowden, William B. et al., *Aust. J. Chem.*, **1985**, 38, 1257).

¹H-NMR (DMSO-d₆) δ: 11.95 (1H, br s), 7.90 (1H, s), 7.84 (1H, d, J=8.7Hz), 7.84 (1H, d, J=2.0Hz), 7.15 (1H, d, J=2.0Hz), 4.24 (2H, s), 3.60 (3H, s), 2.62 (3H, s).

EXAMPLE 258

[6-CHLORO-2-(4-METHYLTHIAZOLE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC
ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(4methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate (Example 257).

m.p.: 239-240 °C.

¹H-NMR (DMSO-d₆) δ: 11.80 (1H, br s), 7.75 (1H, s), 7.68 (1H, d, J=8.4Hz), 7.67 (1H, d, J=2.0Hz), 7.02 (1H, dd, J=2.0, 8.4Hz), 4.03 (2H, s), 2.49 (3H, s).

EXAMPLE 259

METHYL [5-CHLORO-2-[3-(ETHOXYCARONYL)ISOXAZOLE-5-CARBONYL]-
1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in Example 8 (Method B) from methyl *trans*-5-chloro-2-(phenylsulfonylamino)cinnamate (Example 36, step 3) and ethyl 5-(bromoacetyl)isoxazole-3-carboxylate.

¹H-NMR (DMSO-d₆) δ: 12.07 (1H, br s), 7.93 (1H, d, J=1.9Hz), 7.70 (1H, s), 7.39 (1H, dd, J=1.9, 12.8Hz), 7.62 (1H, d, J=12.8Hz), 4.44 (2H, q, J=7.1Hz), 4.15 (2H, s), 3.60 (3H, s), 1.37 (3H, t, J=7.1Hz).

EXAMPLE 260**[5-CHLORO-2-[3-(CARBOXY)ISOXAZOLE-5-CARBONYL]-1H-INDOL-3-YL]ACETIC ACID**

To a solution of methyl [5-chloro-2-[3-(ethoxycarbonyl)isoxazole-5-carbonyl]-1H-indol-3-yl]acetate (Example 259, 314 mg, 0.80 mmol) in acetic acid (20 ml) was added 2N aqueous HCl (6.0 ml) and the mixture was heated at 110 °C for 5 h. The mixture was then cooled to room temperature and concentrated. The residual yellow solids were washed with ethyl acetate and recrystallized from ethyl acetate/hexane to afford 120 mg (43 %) of the title compound as pale yellow solids.

10 m.p.: 200-205°C.

¹H-NMR (DMSO-d₆) δ: 12.01 (1H, s), 7.91 (1H, s), 7.65-7.56 (2H, m), 7.39 (1H, d, J=8.9Hz), 4.06 (2H, s).

EXAMPLE 261**METHYL [6-CHLORO-2-CYCLOPROPANECARBONYL-1H-INDOL-3-YL]ACETATE**

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and bromomethyl cyclopropyl ketone* .

20 ¹H-NMR (CDCl₃) δ: 9.55 (1H, br s), 7.58 (1H, d, J=8.56Hz), 7.26 (1H, d, J=1.97Hz) 7.10 (1H, dd, J=8.56Hz, 1.97Hz), 4.17 (2H, s), 3.73 (3H, s), 2.58-2.49 (1H, m), 1.30-1.25 (2H, m), 1.09-1.02 (2H, m).

*Bromomethylcyclopropyl ketone was prepared from cyclopropyl methyl ketone according to the procedure for preparing 4-(trifluoromethoxy)phenacyl bromide described in Example 189.

25 ¹H-NMR (CDCl₃) δ: 3.91 (2H, s), 2.65 (1H, t, 6.94Hz), 1.20-0.98 (4H, m).

EXAMPLE 262**[6-CHLORO-2-CYCLOPROPANECARBONYL-1H-INDOL-3-YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl 2-(6-chloro-2-cyclopropanecarbonyl-1H-indol-3-yl)acetate (Example 261).

30 m.p.: 207-210 °C.

IR (KBr) ν : 3304, 3013, 1709, 1624, 1566, 1443, 1414, 1387, 1340, 1286, 1248, 1217, 1200, 1157, 1057, 1045, 1022 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.98 (1H, br s), 7.75 (1H, d, $J=8.75\text{Hz}$), 7.48 (1H, d, $J=1.81\text{Hz}$) 7.11 (1H, dd, $J=8.75\text{Hz}$, 1.81Hz), 4.08 (2H, s), 2.73 (1H, quintet, 6.24Hz),
5 1.07 (4H, d, 6.24Hz).

EXAMPLE 263

METHYL [6-CHLORO-2-CYCLOBUTANECARBONYL-1H-INDOL-3-YL]ACETATE

The title compound was prepared according to the procedure described in
10 Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of Example 8, Method A) and bromomethyl cyclobutyl ketone*.

$^1\text{H-NMR}$ (CDCl_3) δ : 9.35 (1H, br s), 7.53 (1H, d, $J=8.72\text{Hz}$), 7.18 (1H, d, $J=1.65\text{Hz}$) 7.07 (1H, dd, $J=8.72\text{Hz}$, 1.65Hz), 4.10 (2H, s), 3.77 (3H, s), 3.72 (1H, m), 2.44-1.86 (6H, m).

15 *Bromomethylcyclobutyl ketone was prepared from cyclobutyl methyl ketone according to the procedure for preparing 4-(trifluoromethoxy)phenacyl bromide described in Example 189.

$^1\text{H-NMR}$ (CDCl_3) δ : 3.88 (2H, s), 3.60 (1H, m), 2.33-1.80 (6H, m)

EXAMPLE 264

20 [6-CHLORO-2-CYCLOBUTANECARBONYL-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl 2-(6-chloro-2-cyclobutanecarbonyl-1H-indol-3-yl)acetate (Example 263).

m.p.: 225-228 $^{\circ}\text{C}$.

25 IR (KBr) ν : 3303, 2954, 1705, 1632, 1564, 1529, 1437, 1412, 1335, 1242, 1213, 1188, 1157, 1056, 1024 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.63 (1H, br s), 7.71 (1H, d, $J=8.72\text{Hz}$), 7.46 (1H, d, $J=1.81\text{Hz}$), 7.09 (1H, dd, $J=8.72\text{Hz}$, 1.81Hz), 4.04 (2H, s), 2.30-1.78 (7H, m).

EXAMPLE 265

30 METHYL [5-(*tert*-BUTYL)-2-(4-CHLOROBENZOYL)-1H-INDOL-3-YL] ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-*tert*-butyl-2-(*p*-toluenesulfonylamino)cinnamate (Example 143, step 2) and 4-chlorophenacyl bromide.

¹H-NMR (CDCl₃) δ: 8.82 (1H, br s), 7.8-7.31 (7H, m), 3.87 (2H, s), 3.67 (3H, s), 1.38 (9H, s).

EXAMPLE 266

[5-(*tert*-BUTYL)-2-(4-CHLOROBENZOYL)-1H-INDOL-3-YL] ACETIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [5-*tert*-butyl-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate (Example 265).

m.p.: 171°C.

IR (KBr) v: 3241, 2963, 1699, 1634, 1589/ 1541, 1394, 1331, 1222, 1091, 1011 cm⁻¹

¹H-NMR (DMSO-d₆) δ: 11.48 (1H, br s), 7.78-7.37 (7H, m), 3.85 (2H, s), 1.34 (9H, s).

EXAMPLE 267

[6-CHLORO-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]-*N,N*-DIMETHYLACETAMIDE

The title compound was prepared according to the procedure described in Example 43 from [6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid (Example 31).

m.p.: 208 °C (decompose).

IR (KBr): 3233, 1655, 1638, 1524, 1398, 1200, 1134 cm⁻¹.

¹H-NMR (CDCl₃) δ: 12.54 (1H, br s), 8.62 (1H, d, J=5.0Hz), 8.15 (1H, br), 7.79 (1H, d, J=8.7Hz), 7.49 (1H, d, J=1.8Hz), 7.40-7.30 (1H, m), 7.09 (1H, dd, J=1.8, 8.7Hz), 4.43 (2H, s), 3.16 (3H, s), 2.98 (3H, s), 2.48 (3H, s).

EXAMPLE 268

[6-CHLORO-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]-*N*-METHYLACETAMIDE

The title compound was prepared according to the procedure described in Example 43 from [6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid (Example 31) and methylamine hydrochloride.

m.p.: 231 °C.

IR (KBr) ν : 3306, 1643, 1595, 1560, 1526, 1277, 1202, 797 cm^{-1} .

$^1\text{H-NMR}$ (CDCl_3) δ : 12.45 (1H, br), 8.64 (1H, d, $J=4.9\text{Hz}$), 8.19 (1H, br), 7.85 (1H, d, $J=8.6\text{Hz}$), 7.50 (1H, br), 7.40 (1H, br d, $J=4.6\text{Hz}$), 7.15 (1H, dd, $J=1.6, 8.9\text{Hz}$), 6.67 (1H, br), 4.14 (2H, s), 2.73 (3H, d, $J=4.8\text{Hz}$), 2.51 (3H, s).

EXAMPLE 269

[5-CHLORO-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]-N-(2-HYDROXYETHYL)ACETAMIDE

The title compound was prepared according to the procedure described in Example 43 from [5-chloro-2-(4-methylpyridine)-1H-indol-3-yl]acetic acid (Example 36) and 2-aminoethanol.

MS (EI) m/z : 371 (M^+).

m.p.: 195.9 °C.

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.28 (1H, br s), 8.69 (1H, d, $J=4.9\text{Hz}$), 7.94 (1H, br s), 7.78 (2H, m), 7.66 (1H, d, $J=8.7\text{Hz}$), 7.56 (1H, m), 7.31 (1H, dd, $J=2.0\text{Hz}, 8.7\text{Hz}$), 3.94 (2H, s), 3.09 (2H, dd, $J=5.93\text{Hz}, 11.86\text{Hz}$), 2.47 (3H, s).

EXAMPLE 270

[5-CHLORO-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]-N-METHOXYACETAMIDE

The title compound was prepared according to the procedure described in Example 43 from [5-chloro-2-(4-methylpyridine)-1H-indol-3-yl]acetic acid (Example 36) and *O*-methylhydroxylamine hydrochloride.

MS (EI) m/z : 357 (M^+).

$^1\text{H-NMR}$ (CDCl_3) δ : 12.53 (1H, br s), 9.54 (1H, br s), 8.65 (1H, d, $J=5.1\text{Hz}$), 8.19 (1H, br s), 7.92 (1H, br s), 7.42 (2H, m), 7.34 (1H, dd, $J=1.7\text{Hz}, 8.9\text{Hz}$), 4.01 (2H, s), 3.74 (3H, s), 2.52 (3H, s).

EXAMPLE 271

2-[5-CHLORO-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]-1-(1-PIPERAZINYL)-1-ETHANONE

STEP 1. **2-[5-Chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-1-(4-*tert*-butoxycarbonyl)-1-piperazinyl)-1-ethanone**

The title compound was prepared according to the procedure described in Example 43 from [5-chloro-2-(4-methylpyridine)-1H-indol-3-yl]acetic acid (Example 36) and *tert*-butyl 1-piperadinecarboxylate.

¹H-NMR (CDCl₃) δ: 12.61 (1H, br s), 8.62 (1H, d, J=5.1Hz), 8.14 (1H, br s), 7.84 (1H, s), 7.42 (1H, d, J=8.7Hz), 7.37 (br d, 1H, J=4.9Hz), 7.30 (1H, dd, J=2.0Hz, 8.9Hz), 4.42 (2H, s), 3.66 (4H, m), 2.50 (3H, s), 1.64 (4H, m), 1.46 (9H, s).

STEP2. 2-[5-Chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-1-(1-piperazinyl)-1-ethanone

To a solution of [5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-*N*-(4-*tert*-butoxycarbonylpiperadino)acetamide (step 1, 152.6 mg, 0.042 mmol) in THF (1 ml) was added dropwise trifluoroacetic acid (2 ml) at 0°C. The mixture was stirred at room temperature for 1.5h and then concentrated. The residue was diluted with dichloromethane (25 ml), washed with saturated aqueous sodium bicarbonate (25 ml). The aqueous layer was extracted with dichloromethane (25 ml x 2). The combined organic layers were dried (Na₂SO₄) and concentrated. The residue was purified by flash column chromatography on eluting with methanol/dichloromethane (1:10) to afford 80.8 mg of the title compound as yellow crystals.

MS (EI) m/z: 396(M⁺).

m.p.: 205.0 °C.

IR (KBr) ν: 3244, 1647, 1595, 1525, 1429, 1205 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.28 (1H, br s), 8.68 (1H, d, J=4.9Hz), 7.90 (1H, br s), 7.81 (1H, br s), 7.65 (1H, d, J=8.9Hz), 7.56 (br d, 1H, J=4.1Hz), 7.31 (1H, dd, J=1.8Hz, 8.9Hz), 4.15 (2H, br s), 3.50-3.15 (4H, m), 2.70-2.55 (4H, m), 2.46 (3H, s).

EXAMPLE 272

[5-CHLORO-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]-*N*-(2-AMINOETHYL)ACETAMIDE

STEP 1. [5-Chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-*N*-(2-*tert*-butoxycarbonylaminoethyl)acetamide

The title compound was prepared according to the procedure described in Example 43 from [5-chloro-2-(4-methylpyridine)-1H-indol-3-yl]acetic acid (Example 36) and *N*-(2-aminoethyl)carbamic acid *tert*-butyl ester.

¹H-NMR (CDCl₃) δ: 12.29 (1H, br s), 8.68 (1H, d, J=4.9Hz), 7.94 (1H, br s), 7.78 (2H, m), 7.67 (1H, d, J=8.9Hz), 7.57 (1H, br d, J=4.9Hz), 7.32 (dd, 1H, J=2.0Hz, 8.7Hz), 6.73 (1H, m), 3.94 (2H, s), 3.05-2.95 (4H, m), 2.47 (3H, s), 1.35 (9H, s).

STEP2.[5-Chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-N-(2-aminoethyl)acetamide

The title compound was prepared according to the procedure described in step 2 of Example 273 from [5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-N-(2-*tert*-butoxycarbonylaminoethyl)acetamide (step 1).

MS (EI) m/z: 370 (M⁺).

m.p.: 165.7 °C.

IR (KBr) ν: 3346, 2927, 1665, 1627, 1593, 1515, 1435, 1267, 1207 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.28 (1H, br s), 8.69 (1H, d, J=4.9Hz), 7.93 (1H, br s), 7.80-7.76 (2H, m), 7.66 (1H, d, J=8.9Hz), 7.57 (1H, m), 7.31 (dd, 1H, J=2.0Hz, 8.9Hz), 3.94 (2H, s), 3.01 (2H, q, J=5.77Hz), 2.55-2.45 (2H, m), 2.47 (3H, s).

EXAMPLE 273

2-[5-CHLORO-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]-1-(3-AMINO-1-PYRROLIDINYL)-1-ETHANONE

STEP1.2-[5-Chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-1-[3-(*tert*-butoxycarbonylamino)-1-pyrrolidinyl]-1-ethanone

The title compound was prepared according to the procedure described in Example 43 from [5-chloro-2-(4-methylpyridine)-1H-indol-3-yl]acetic acid (Example 36) and 3-(*tert*-butoxycarbonylamino)pyrrolidine.

¹H-NMR (CDCl₃) δ: 12.56 (1H, br s), 8.62 (1H, d, J=4.9Hz), 8.18 (1H, br s), 7.78 (1H, br s), 7.42 (1H, d, J=8.9Hz), 7.35 (1H, br d, J=4.9Hz), 7.29 (2H, m), 4.89 (1H, br d, J=25.05), 4.36-3.45 (7H, m), 2.48 (3H, s), 2.35-1.80 (2H, m), 1.46 (9H, s).

STEP2.2-[5-Chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-1-(3-amino-1-pyrrolidinyl)-1-ethanone

The title compound was prepared according to the procedure described in step 2 of Example 271 from 2-[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-1-[3-(*tert*-butoxycarbonylamino)-1-pyrrolidinyl]-1-ethanone (step 1).

MS (EI) m/z: 396(M⁺).

m.p.: 179.2 °C.

IR (KBr) ν : 3238, 2876, 1638, 1595, 1526, 1423, 1203 cm^{-1} .

$^1\text{H-NMR}$ (DMSO-d_6) δ : 12.26 (1H, br s), 8.68 (1H, d, $J=4.9\text{Hz}$), 7.89 (1H, br s), 7.81 (1H, d, $J=1.81\text{Hz}$), 7.65 (1H, d, $J=8.9\text{Hz}$), 7.55 (1H, br d, $J=5.1\text{Hz}$), 7.31 (dd, 1H, $J=2.0\text{Hz}$, 8.7Hz), 4.07 (1H, s), 4.05 (1H, s), 3.70-2.90 (5H, m), 2.46 (3H, s), 2.10-1.80 (1H, m), 1.75-1.45 (1H, m).

EXAMPLE 274

METHYL [6-CHLORO-2-(4-CHLOROBENZOYL)-5-FLUORO-1H-INDOL-3-YL]ACETATE

10 STEP 1. Methyl *trans*-2-amino-4-chloro-5-fluorocinnamate

The title compound was prepared according to the procedure described in step 1 of Example 133 from 2-bromo-5-chloro-4-fluoroaniline (JP 01311056 A2, Nippon Kayaku Co., Ltd., Japan).

$^1\text{H-NMR}$ (CDCl_3) δ : 7.69 (1H, d, $J=15.8\text{Hz}$), 7.15 (1H, d, $J=9.7\text{Hz}$), 6.74 (1H, d, $J=6.4\text{Hz}$), 6.31 (1H, d, $J=15.8\text{Hz}$), 3.81 (3H, s).

15 STEP 2. Methyl *trans*-4-chloro-5-fluoro-2-(phenylsulfonylamino)cinnamate

The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-2-amino-4-chloro-5-fluorocinnamate (step 1).

$^1\text{H-NMR}$ (CDCl_3) δ : 7.71-7.68 (1H, m), 7.57-7.40 (6H, m), 7.23 (1H, d, $J=9.4\text{Hz}$), 6.09 (1H, d, $J=15.8\text{Hz}$), 3.77 (3H, s).

STEP 3. Methyl [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetate

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-5-fluoro-2-(phenylsulfonylamino)cinnamate (step 2).

$^1\text{H-NMR}$ (CDCl_3) δ : 9.05 (1H, br.s), 7.74 (2H, d, $J=8.7\text{Hz}$), 7.49 (2H, d, $J=8.7\text{Hz}$), 7.35-7.32 (2H, m), 3.77 (2H, s), 3.66 (3H, s)

EXAMPLE 275

[6-CHLORO-2-(4-CHLOROBENZOYL)-5-FLUORO-1H-INDOL-3-YL]ACETIC

30 ACID

The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetate. m.p.: 215-220 °C.

IR(KBr)v: 1709, 1626, 1585, 1529, 1456, 1439, 1279, 1250 cm⁻¹.

- 5 ¹H-NMR (DMSO-d₆) δ: 11.85 (1H, br.s), 7.79-7.75 (3H, m), 7.67-7.63 (2H, m), 7.60-7.58 (1H, m), 3.83 (2H, s).

EXAMPLE 276

METHYL [6-CHLORO-5-FLUORO-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE

- 10 The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-5-fluoro-2-(phenylsulfonylamino)cinnamate (Example 274, step 2).

¹H-NMR (CDCl₃) δ: 12.54 (1H, br.s), 8.58 (1H, d, J=4.9Hz), 8.14 (1H, m), 7.56-7.53 (1H, m), 7.39-7.34 (2H, m), 4.25 (2H, s), 3.75 (3H, s), 2.48 (3H, s).

- 15 EXAMPLE 277

[6-CHLORO-5-FLUORO-2-(4-METHYLPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

- The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 276).

mp: 219.5 °C.

IR (KBr) v: 1732, 1709, 1647, 1597, 1529, 1279, 1252, 1204 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 12.35 (1H, br.s), 8.70 (1H, d, J=5.1Hz), 7.96 (1H, s), 7.87 (1H, d, J=6.6Hz), 7.81 (1H, d, J=10.1Hz), 7.59-7.58 (1H, m), 4.05 (2H, s), 2.47 (3H, s).

- 25 EXAMPLE 278

METHYL [6-CHLORO-2-[4-(1-HYDROXYETHYL)PYRIDINE-2-CARBONYL]-1H-INDOL-3-YL]ACETATE

STEP 1. methyl [6-chloro-2-[4-[1-(tert-butyldimethylsilyloxy)ethyl]pyridine-2-carbonyl]-1H-indol-3-yl]acetate

- 30 The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-chloro-2-(phenylsulfonylamino)cinnamate (step 1 of

Example 8, Method A) and 2-bromoacetyl-4-[1-(*tert*-butyldimethylsilyloxy)ethyl]pyridine*.

¹H-NMR (CDCl₃) δ: 12.49 (1H, br s), 8.71 (1H, d, J=4.9Hz), 8.24 (1H, d, J=1.5Hz), 7.61-7.63 (2H, m), 7.53 (1H, d, J=1.8Hz), 7.13 (1H, dd, J=1.8, 8.7Hz), 4.96 (1H, q, J=6.4Hz), 4.32 (2H, s), 3.73 (3H, s), 1.45 (3H, d, J=6.4Hz), 0.93 (9H, s), 0.10 (3H, s), 0.03 (3H, s).

*2-Bromoacetyl-4-[1-(*tert*-butyldimethylsilyloxy)ethyl]pyridine was prepared as follows;

4-[1-(Trimethylsilyloxy)ethyl]-2-pyridinecarbonitrile:

10 The title compound was prepared from 4-(1-hydroxyethyl)pyridine-*N*-oxide (C.W.Muth et al., *J. Heterocycl. Chem.*, 1972, 9, 1299) according to the procedure for preparing 4-chloro-2-pyridinecarbonitrile described in Example 33.

¹H-NMR (CDCl₃) δ: 8.64 (1H, d, J=5.1Hz), 7.69-7.70 (1H, m), 7.45-7.48 (1H, m), 4.88 (2H, q, J=6.4Hz), 1.43 (3H, d, J=6.6Hz), 0.14 (9H, s).

15 4-[1-(*tert*-Butyldimethylsilyloxy)ethyl]-2-pyridinecarbonitrile:

To a solution of 4-[1-(trimethylsilyloxy)ethyl]-2-pyridinecarbonitrile (39.04 g, 0.1624 mol) in THF (200 ml) was added a solution of tetrabutylammonium fluoride in THF (1M, 178.6 ml, 0.1786 mol) at room temperature. After stirring for 0.5 h, the mixture was concentrated. The residue was diluted with ethyl acetate (300 ml) and washed with water (200 ml). The aqueous layer was then extracted with dichloromethane (200 ml x 2). The combined organic layers were dried (MgSO₄) and concentrated. The residual oil was dissolved in DMF (200 ml). To the solution was added *tert*-butyldimethylsilylchloride (36.72 g, 0.2436 mol) and imidazole (22.11 g, 0.3248 mol) at room temperature. After stirring for 19h, diethyl ether (500 ml) and water (200 ml) were added to the mixture and the organic layer was separated. The organic layer was washed with water (100 ml x 2), dried (MgSO₄) and concentrated. The residue was purified by flash column chromatography eluting with ethyl acetate/hexane (1:20) to give 39.23 g (92 %) of the title compound as an oil.

25 ¹H-NMR (CDCl₃) δ: 8.64 (1H, d, J=5.1Hz), 7.67-7.68 (1H, m), 7.46-7.48 (1H, m), 4.89 (1H, q, J=6.4Hz), 1.42 (3H, d, J=6.4Hz), 0.92 (9H, s), 0.10 (6H, s).

2-Acetyl-4-[1-(*tert*-butyldimethylsilyloxy)ethyl]pyridine:

The title compound was prepared from 4-[1-(*tert*-butyldimethylsilyloxy)ethyl]-2-pyridinecarbonitrile according to the procedure for preparing 2-acetyl-4-chloropyridine described Example 33.

- 5 ¹H-NMR (CDCl₃) δ: 8.62 (1H, d, J=4.9Hz), 7.95-7.96 (1H, m), 7.49-7.52 (1H, m), 4.91 (1H, q, J=6.4Hz), 2.73 (3H, s), 1.41 (3H, d, J=6.4Hz), 0.91 (9H, s), 0.10 (6H, s).

2-Bromoacetyl-4-[1-(*tert*-butyldimethylsilyloxy)ethyl]pyridine

- The title compound was prepared according to the procedure for preparing 2-bromoacetyl-4-(*tert*-butyldimethylsilyloxymethyl)pyridine described in Example 95
10 from 2-acetyl-4-[1-(*tert*-butyldimethylsilyloxy)ethyl]pyridine.

STEP 2. Methyl [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate.

- The title compound was prepared according to the procedure described in step 2 of Example 95 from methyl [6-chloro-2-[4-[1-(*tert*-
15 butyldimethylsilyloxy)ethyl]pyridine-2-carbonyl]-1H-indol-3-yl]acetate (step 1).

¹H-NMR (DMSO-d₆) δ: 12.34 (1H, br s), 8.77 (1H, d, J=5.1Hz), 8.11 (1H, s), 7.81 (1H, d, J=8.7Hz), 7.75 (1H, d, J=1.6Hz), 7.70 (1H, dd, J=1.2, 4.9Hz), 7.13 (1H, dd, J=1.8, 8.6Hz), 5.61 (1H, d, J=4.6Hz), 4.84-4.93 (1H, m), 4.17 (2H, s), 3.59 (3H, s), 1.39 (3H, d, J=6.4Hz).

20 **EXAMPLE 279**

[6-CHLORO-2-[4-(1-HYDROXYETHYL)PYRIDINE-2-CARBONYL]-1H-INDOL-3-YL]ACETIC ACID

- The title compound was prepared according to the procedure described in Example 58 from methyl [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-
25 indol-3-yl]acetate (Example 278).

m.p.: 193-194°C.

IR (KBr) v: 3464, 1707, 1632, 1591, 1526, 1250, 1225, 1192, 1144, 914 cm⁻¹.

- ¹H-NMR (DMSO-d₆) δ: 12.30 (1H, br s), 8.78 (1H, d, J=5.1Hz), 8.11 (1H, s), 7.79 (1H, d, J=8.6Hz), 7.74 (1H, d, J=1.6Hz), 7.71 (1H, d, J=5.3Hz), 7.12 (1H, dd, J=1.8, 8.7Hz),
30 5.60 (1H, d, J=4.4Hz), 4.84-4.93 (1H, m), 4.09 (2H, s), 1.39 (3H, d, J=6.6Hz).

EXAMPLE 280**[6-CHLORO-2-(4-ETHYL-3-FLUOROPYRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 112 from methyl [6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 113).

m.p.; 193-194°C.

IR (KBr) ν : 3256, 1707, 1645, 1529, 1420, 1227, 1180, 1159, 1153 cm^{-1} .

MS (EI) m/z : 360 (M^+).

$^1\text{H-NMR}$ (DMSO-d_6) δ : 11.84 (1H, br s), 8.46 (1H, d, $J=5.3\text{Hz}$), 7.77 (1H, d, $J=8.7\text{Hz}$), 7.67 (1H, t, $J=5.1\text{Hz}$), 7.51 (1H, d, $J=1.8\text{Hz}$), 7.12 (1H, dd, $J=1.8, 8.6\text{Hz}$), 3.74 (2H, s), 2.76 (2H, q, $J=7.6\text{Hz}$), 1.25 (3H, t, $J=7.6\text{Hz}$).

EXAMPLE 281**[6-CHLORO-2-(2-NITROBENZOYL)-1H-INDOL-3-YL]ACETIC ACID****15 STEP 1. Polymer-bound *trans* 4-chloro-2-(phenylsulfonylamino)cinnamate**

To a mixture of Wang resin (200-400 mesh, 1.37 g, ca. 0.89 mmol) and *trans* 4-chloro-2-(phenylsulfonylamino)cinnamic acid (600 mg, 1.77 mmol) was added dichloromethane (10 ml) and *N,N*-diisopropylethylamine (1.86 ml, 10.7 mmol). The mixture was allowed to stand for 1h, and then 4-dimethylaminopyridine (22 mg, 0.18 mmol) and WSC (339 mg, 1.77 mmol) were added. The mixture was agitated for 18h and filtrated. The residual resin was washed with water (20 ml x 3), methanol (20 ml x 3), acetone (20 ml x 3), dichloromethane (20 ml x 3) and dried to give 1.65 g of the title compound.

STEP 2. [6-chloro-2-(2-nitrobenzoyl)-1H-indol-3-yl]acetic acid

25 To a mixture of polymer-bound *trans* 4-chloro-2-(phenylsulfonylamino)cinnamate (step 1, 100mg, 53 μmol) and 2-nitrophenacyl bromide (39 mg, 0.16 mmol) in acetone (3 ml) was added potassium carbonate (37 mg, 0.27 mmol). The mixture was agitated for 18 h. and filtrated. The residual resin was washed with water (20 ml x 3), acetone (20ml x 3), dichloromethane (20 ml x 3) and THF (20 ml x 2) and dried. The resin was diluted with THF (4 ml) and then DBU was added (40 μl , 0.27mmol). After agitating for 6 h, the resin was filtered off and

washed with THF (20 ml x 3), acetone (20 ml x 3) and dichloromethane (20 ml x 3). To the resin was added 95% trifluoroacetic acid in dichloromethane (5 ml) and the mixture was agitated for 3h. The mixture was filtered and the residue was washed with dichloromethane (20ml x 5). The filtrate was concentrated and the residue was
5 purified by HPLC (MeOH/AcONH₄ aqueous solution=60/40-90/10) to give 3.2 mg (17%) of the title compound.

MS (ESI) m/z : 359 (MH⁺).

EXAMPLE 282

[6-CHLORO-2-(2,4-DIMETHOXYBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

10 The title compound was prepared according to the procedure described in Example 281 from 4-chloro-2-(phenylsulfonylamino)cinnamic acid.

MS (ESI) m/z : 374 (MH⁺).

EXAMPLE 283

[6-CHLORO-2-(4-DIFLUOROMETHOXYBENZOYL)-1H-INDOL-3-YL]ACETIC

15 **ACID**

The title compound was prepared according to the procedure described in Example 289 from 4-chloro-2-(phenylsulfonylamino)cinnamic acid.

MS (ESI) m/z : 380 (MH⁺).

EXAMPLE 284

20 **[6-CHLORO-2-(2,5-DIMETHOXYBENZOYL)-1H-INDOL-3-YL]ACETIC ACID**

The title compound was prepared according to the procedure described in Example 281 from 4-chloro-2-(phenylsulfonylamino)cinnamic acid.

MS (ESI) m/z : 374 (MH⁺).

EXAMPLE 285

25 **METHYL [5-ACETYL-2-(4-CHLOROBENZOYL)-1H-INDOL-3-YL]ACETATE**

STEP 1. 4-ACETYL-2-BROMOANILINE

To a stirred suspension of 4-acetamido-3-bromoacetophenone in ethanol (12 ml) was added dropwise hydrochloric acid (3 ml) at 0 °C. The reaction mixture was stirred under reflux condition for 4.5 h. The mixture was cooled and concentrated.
30 The residual solids were partitioned between saturated aqueous sodium bicarbonate (50 ml) and diethyl ether (50 ml). The aqueous layer was extracted with ethyl acetate (50

ml x 2). The combined organic layers were dried (MgSO₄) and concentrated to afford 1.79g (quant.) of the title compound as a brown oil.

¹H-NMR (CDCl₃) δ: 8.06 (1H, d, 1.97Hz), 7.76-7.72 (1H, m), 6.74 (1H, d, 8.40Hz), 4.60 (1H, br s), 2.50 (s, 3H)

5 STEP 2. Methyl *trans*-(5-acetyl-2-amino)cinnamate

The title compound was prepared according to the procedure described in step 1 of Example 133 from 4-acetyl-2-bromoaniline (step 1) and methyl acrylate.

¹H-NMR (CDCl₃) δ: 8.02 (1H, d, 2.13Hz), 7.81-7.75 (2H, m), 6.70 (1H, d, 8.56Hz), 6.44 (1H, d, 15.8Hz), 4.55 (2H, br s), 3.81 (3H, s), 2.53 (3H, s).

10 STEP 3. Methyl *trans*-5-acetyl-2-(*p*-toluenesulfonylamino)cinnamate

The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-(5-acetyl-2-amino)cinnamate (step 2) and *p*-toluenesulfonyl chloride.

¹H-NMR (CDCl₃) δ: 8.01 (1H, d, 2.00Hz), 7.89 (1H, dd, 8.59Hz, 1.97Hz), 7.68-7.22 (7H, m), 6.30 (1H, s), 6.15 (1H, d, 15.8Hz), 3.81 (3H, s), 2.57 (3H, s), 2.38 (3H, s).

STEP 4. Methyl [5-acetyl-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-5-acetyl-3-(*p*-toluenesulfonylamino)cinnamate (step 3) and 4-chlorophenacyl bromide.

20 ¹H-NMR (CDCl₃) δ: 9.2 (1H, br s), 8.34-7.75 (4H, m), 7.52 - 7.45 (3H, m), 3.89 (2H, s), 3.69 (3H, s), 2.68 (3H, s).

EXAMPLE 286

[5-ACETYL-2-(4-CHLOROBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

25 The title compound was prepared according to the procedure described in Method B of Example 9 from methyl [5-acetyl-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate (Example 285).

m.p. : 225°C.

IR (KBr) v: 3281, 1703, 1666, 1643, 1614, 1574, 1539, 1452, 1425, 1402, 1364, 1263, 1240, 1178, 1092, 1011, 959 cm⁻¹.

30 ¹H-NMR (DMSO-d₆) δ: 12.01 (1H, br s), 8.49 (1H, br s), 7.93 - 7.89 (1H, m), 7.78 (2H, d, 8.56Hz), 7.65 (d, 2H, 8.56Hz), 7.52 (1H, d, 8.72Hz), 3.93 (2H, s), 2.63 (3H, s).

EXAMPLE 287**METHYL [6-FLUORO-2-(4-METHYLPRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETATE****STEP 1. Methyl *trans*-(4-fluoro-2-nitro)cinnamate**

5 The title compound was prepared according to the procedure described in step 1 of Example 133 from 3-fluoro-6-iodonitrobenzene and methyl acrylate .

¹H-NMR (CDCl₃) δ: 8.67 (1H, d, 15.8Hz), 7.78 (1H, dd, 8.07Hz, 2.65Hz), 7.68-7.63 (2H, m), 6.34 (1H, d, 15.8Hz), 3.84 (3H, s).

STEP 2. Methyl *trans*-(2-amino-4-fluoro)cinnamate

10 The title compound was prepared according to the procedure described in step 2 of Example 36 from methyl *trans*-(4-fluoro-2-nitro)cinnamate (step 1).

¹H-NMR (CDCl₃) δ: 7.75 (1H, d, 15.8Hz), 7.37-7.31 (1H, m), 6.50-6.37 (2H, m), 6.32-6.26 (1H, m), 4.13 (2H, br s), 3.80 (3H, s).

STEP 3. Methyl *trans*-4-fluoro-2-(*p*-toluenesulfonylamino)cinnamate

15 The title compound was prepared according to the procedure described in step 1 of Example 8 (Method A) from methyl *trans*-(2-amino-4-fluoro)cinnamate (step 2) .

¹H-NMR (CDCl₃) δ: 7.64 (1H, d, 8.40Hz), 7.46-7.19 (4H, m), 6.94-6.87 (1H, m), 6.77 (1H, s), 6.16-6.10 (1H, m), 3.79 (3H, s) , 2.38 (3H, s) .

STEP 4. Methyl [6-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-

20 **yl]acetate**

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-fluoro-2-(*p*-toluenesulfonylamino)cinnamate (step 3).

¹H-NMR (CDCl₃) δ: 12.49 (1H, br s), 8.61 (1H, d, 4.94Hz), 8.17-7.61 (2H, m), 7.36-7.14 (2H, m), 6.98-6.90 (1H, m), 4.31 (2H, s), 3.73 (3H, s), 2.47 (3H, s).

25 **EXAMPLE 288**

[6-FLUORO-2-(4-METHYLPRIDINE-2-CARBONYL)-1H-INDOL-3-YL]ACETIC ACID

30 The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate (Example 287).

m.p.: 208 °C.

IR (KBr) v: 3238, 1701, 1638, 1597, 1533, 1398, 1281, 1211, 1132, 1003 cm⁻¹.

¹H-NMR (CDCl₃) δ: 12.28 (1H, br s), 8.70 (1H, d, 4.94Hz), 7.96 (1H, s), 7.83-7.77 (1H, m), 7.57 (1H, d, 4.94Hz), 7.42 (1H, dd, 10.2Hz, 2.13Hz), 7.03-6.95 (1H, m), 4.09 (2H, s), 2.47 (3H, s).

EXAMPLE 289

METHYL [6-FLUORO-2-(4-CHLOROBENZOYL)-1H-INDOL-3-YL] ACETATE

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-4-fluoro-3-(*p*-toluenesulfonylamino)cinnamate (step 3 of Example 287) and 4-chlorophenacylbromide.

¹H-NMR (CDCl₃) δ: 9.10 (1H, br s), 7.76-7.71 (2H, m), 7.60-7.54 (2H, m), 7.50-7.45 (2H, m), 7.04-6.90 (2H, m), 3.81 (2H, s), 3.66 (3H, s).

EXAMPLE 290

[6-FLUORO-2-(4-CHLOROBENZOYL)-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Example 9 (Method B) from methyl [6-fluoro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate (Example 289).

m.p.: 214 °C.

IR(KBr)v: 3335, 1699, 1618, 1605, 1587, 1531, 1425, 1327, 1267, 1231, 1134, 1094, 1001 cm⁻¹.

¹H-NMR (DMSO-d₆) δ: 11.73 (1H, br s), 7.78-7.63 (5H, m), 7.17 (1H, dd, 9.72Hz, 2.13Hz), 7.04-6.96 (1H, m), 3.84 (2H, s).

EXAMPLE 291

[2-(4-METHYLPYRIDINE-2-CARBONYL)-5-METHYLTHIO-1H-INDOL-3-

YL]ACETIC ACID

STEP 1. Methyl *trans*-(2-amino-5-methylthio)cinnamate

The title compound was prepared according to the procedure described in step 1 of Example 133 from 2-bromo-4-methylthioxylaniline (JP80-122756) and methyl acrylate.

¹H-NMR (CDCl₃) δ: 7.77 (1H, d, 15.8Hz), 7.38 (1H, d, 2.13Hz), 7.19 (1H, dd, 8.40Hz, 2.13Hz), 6.66 (1H, d, 8.40Hz), 6.36 (1H, d, 15.8Hz), 3.97 (1H, br s), 3.80 (3H, s), 2.43

(3H, s).

STEP 2. Methyl *trans*-2-*p*-toluenesulfonylamino-5-methylthiocinnamate

The title compound was prepared according to the procedure described in step 1 of Example 8 from methyl *trans*-(2-amino-4-methylthio)cinnamate (step 1) and *p*-toluenesulfonyl chloride.

¹H-NMR (CDCl₃) δ: 7.81 (2H, d, 8.40Hz), 7.56-6.91 (8H, m), 6.13 (1H, d, 15.6Hz), 3.79 (3H, s), 2.46 (3H, s), 2.39 (3H, s).

STEP 3. Methyl [2-(4-methylpyridine-2-carbonyl)-5-methylthio-1H-indol-3-yl]acetate

The title compound was prepared according to the procedure described in Example 57 from methyl *trans*-2-*p*-toluenesulfonylamino-5-methylthiocinnamate (step 2) and 2-bromoacetyl-4-methylpyridine hydrobromide (Preparation is described in step 2 of Example 31).

¹H-NMR (CDCl₃) δ: 12.47 (1H, br s), 8.62 (1H, d, 4.78Hz), 8.19-8.17 (1H, m), 7.65-7.62 (1H, m), 7.47-7.30 (3H, m), 4.31 (2H, s), 3.73 (3H, s), 2.48 (3H, s).

EXAMPLE 292

[2-(4-METHYLPYRIDINE-2-CARBONYL)-5-METHYLTHIO-1H-INDOL-3-YL]ACETIC ACID

The title compound was prepared according to the procedure described in Method B of Example 9 from methyl [2-(4-methylpyridine-2-carbonyl)-5-methylthio-1H-indol-3-yl]acetate (Example).

¹H-NMR (DMSO-d₆) δ: 12.22 (1H, br s), 8.70 (1H, d, 4.62Hz), 7.95-7.17 (4H, m), 4.08 (2H, s), 2.47 (3H, s), 2.35 (3H, s).

MS (EI) m/z: 340 (M⁺).

Example 293

3-[4-Chloro-2-(toluene-4-sulfonylamino)-phenyl]-acrylic acid ethyl ester

To 3-(2-amino-4-chloro-phenyl)-acrylic acid ethyl ester (18.0g, 79.8 mmol) in dichloromethane (144 ml) was added pyridine (9.04 ml, 112 mmol) and *p*-toluenesulfonyl chloride (16.0g, 83.9 mmol). The reaction was stirred at room temperature for 18 hours and poured into 1N hydrochloric acid (150 ml). The layers were separated and the organic layer was dried over magnesium sulfate, filtered, and concentrated. The resulting solid was slurried in hexanes and filtered to afford 3-[4-

chloro-2-(toluene-4-sulfonylamino)-phenyl]-acrylic acid ethyl ester (28.3g, 93%). mp = 124-127 °C. ¹H NMR (300 MHz, CDCl₃) δ 1.35 (t, 3, J = 7.2), 2.40 (s, 3), 4.27 (q, 2, J = 7.2), 6.12 (dd, 1, J = 15.9, 0.9), 7.20-7.39 (m, 4), 7.39 (d, 1, J = 8.6), 7.48-7.53 (m, 2), 7.62 (d, 2, J = 8.3). ¹³C NMR (75 MHz, CDCl₃) δ 15.50, 22.79, 62.24, 122.48, 127.98, 128.49, 129.34, 129.50, 131.05, 136.82, 137.00, 137.73, 138.93, 145.55, 167.59. IR 3214, 1694, 1631, 1318, 1167 cm⁻¹. Analysis calculated for C₁₈H₁₈ClNO₄S: C, 56.91; H, 4.78; N, 3.69. Found: C, 57.10; H, 5.08; N, 3.70.

Example 294

[6-Chloro-2-(4-chloro-benzoyl)-1H-indol-3-yl]-acetic acid

To a solution of 3-[4-chloro-2-(toluene-4-sulfonylamino)-phenyl]-acrylic acid ethyl ester (13.0g, 34.2 mmol) in *N,N*-dimethylacetamide (120 ml) was added potassium carbonate (9.45g, 68.4 mmol) and 2-bromo-4'-chloroacetophenone (8.78g, 37.6 mmol) and the reaction was stirred at room temperature for 15 minutes. 1N Sodium hydroxide (130 ml) was added and the reaction mixture was heated to 100 °C for 8 hours. The reaction mixture was cooled to room temperature and poured into a separatory funnel and washed with methyl *t*-butyl ether (2 X 200 ml). The aqueous layer was acidified to pH 1 with 6N hydrochloric acid and was extracted with ethyl acetate (150 ml). The solvent was removed under reduced pressure and to the resulting oil was added isopropyl alcohol (24 ml) and water (48 ml). A solid precipitated and the slurry was stirred 12 hours. The precipitate was filtered, washed with water, and dried to provide [6-chloro-2-(4-chloro-benzoyl)-1H-indol-3-yl]-acetic acid (9.73g, 82%). mp 181-183 °C. ¹H NMR (400 MHz, *d*₆-DMSO) δ 3.79 (s, 2), 7.08 (dd, 1, J = 8.5, 1.9), 7.42 (d, 1, J = 1.9), 7.60 (d, 2, J = 8.5), 7.62-7.73 (m, 3), 11.74 (bs, 1), 12.22 (bs, 1). ¹³C NMR (75 MHz, *d*₆-DMSO) δ 31.74, 113.25, 118.00, 121.93, 123.96, 127.69, 130.10, 131.30, 132.02, 133.59, 138.02, 138.37, 138.58, 173.20, 188.29. IR 3314, 1710, 1700, 1618, 1522, 1323, 1227, 1093, 941 cm⁻¹. Analysis calculated for C₁₇H₁₁Cl₂NO₃: C, 58.64; H, 3.18; N, 4.02. Found: C, 58.58; H, 3.22; N, 3.93.

Example 295

3-{4-Chloro-2-[[2-(4-chloro-phenyl)-2-oxo-ethyl]-(toluene-4-sulfonyl)-amino]-phenyl}-acrylic acid ethyl ester

To a solution of 3-[4-chloro-2-(toluene-4-sulfonylamino)-phenyl]-acrylic acid ethyl ester (3.00g, 7.90 mmol) in *N,N*-dimethylacetamide (15.0 ml) was added potassium carbonate (2.18g, 15.8 mmol) and 2-bromo-4'-chloroacetophenone (2.03g, 8.69 mmol). The reaction was stirred for 30 minutes, poured into 1N hydrochloric acid (30 ml) and extracted with methyl *t*-butyl ether (2 X 30 ml). The organic extracts were dried over magnesium sulfate, filtered, concentrated to a low volume. Hexanes was added and a solid precipitated. The precipitate was filtered to provide 3-{4-chloro-2-[[2-(4-chloro-phenyl)-2-oxo-ethyl]-(toluene-4-sulfonyl)-amino]-phenyl}-acrylic acid ethyl ester (3.19g, 76%). mp = 162-165 °C. ¹H NMR (300 MHz, CDCl₃) d 1.38 (t, 3, J = 7.2), 2.47 (s, 3), 4.28 (q, 2, J = 7.2), 5.00 (bs, 2), 6.23 (d, 1, J = 16.0), 7.29-7.36 (m, 4), 7.47 (d, 2, J = 8.7), 7.54 (d, 1, J = 8.4), 7.59 (d, 2, J = 8.3), 7.74 (d, 1, J = 16.0), 7.88 (d, 2, J = 8.7). ¹³C NMR (75 MHz, CDCl₃) d 14.33, 21.62, 57.72, 60.66, 112.49, 120.69, 128.07, 129.21, 129.58, 129.70, 131.22, 132.80, 133.83, 134.80, 135.84, 138.55, 139.28, 140.38, 144.46, 166.02, 191.70. IR 1720, 1698, 1590, 1338, 1313, 1179, 1161, 1089 cm⁻¹. Analysis calculated for C₂₆H₂₃Cl₂NO₅S: C, 58.65; H, 4.35; N, 2.63. Found: C, 58.74; H, 4.56; N, 2.72.

Example 296

cis- and *trans*-[6-Chloro-2-(4-chloro-benzoyl)-1-(toluene-4-sulfonyl)-2,3-dihydro-1H-indol-3-yl]-acetic acid ethyl ester

To 3-{4-chloro-2-[[2-(4-chloro-phenyl)-2-oxo-ethyl]-(toluene-4-sulfonyl)-amino]-phenyl}-acrylic acid ethyl ester (1.00g, 1.88 mmol) in *N,N*-dimethylacetamide (5.0 ml) was added potassium carbonate (0.520g, 3.76 mmol). The reaction mixture was stirred for four hours, poured in 1N hydrochloric acid (30 ml) and extracted with methyl *t*-butyl ether (2 X 30 ml). The organic extracts were dried with magnesium sulfate, filtered, and concentrated. The resulting solid was purified by chromatography on silica gel (ethyl acetate/hexanes 20/80) to provide [6-chloro-2-(4-chloro-benzoyl)-1-(toluene-4-sulfonyl)-2,3-dihydro-1H-indol-3-yl]-acetic acid ethyl ester as a 1 to 9 mixture of *cis* and *trans* isomers (0.488g, 49%). Some of the ¹H NMR (300 MHz, CDCl₃) significant signals are: d 1.09 (t, J = 7.2), 1.19 (t, J = 7.2), 2.44 (s), 5.40 (d, J = 4.0), 5.99 (d, J = 9.7), 6.92 (dd, J = 8.1, 1.1), 7.04 (dd, J = 8.1, 1.9), 7.52 (d, J = 8.4), 7.57 (d, J = 1.9), 7.73 (d, J = 8.3), 7.99 (d, J = 8.6). Lc-MS

analysis was performed on the mixture of diastereoisomers and indicated to products with identical mass of 531 ($M+H^+$).

Example 297

[6-Chloro-2-(4-chloro-benzoyl)-1H-indol-3-yl]-acetic acid ethyl ester

5 To a solution of 3-[4-chloro-2-(toluene-4-sulfonylamino)-phenyl]-acrylic acid ethyl ester (3.00g, 7.90 mmol) in *N,N*-dimethylacetamide (15.0 ml) was added potassium carbonate (2.18g, 15.8 mmol) and 2-bromo-4'-chloroacetophenone (2.03g, 8.69 mmol). The reaction was stirred 30 minutes and 1,8-diazabicyclo[5.4.0]undec-7-ene (3.54 ml, 23.7 mmol) was added. The reaction mixture was stirred one hour,
10 poured into 1N hydrochloric acid (30 ml) and extracted with methyl *t*-butyl ether (2 X 30 ml). The organic extracts were dried with magnesium sulfate, filtered, and concentrated to provide a solid which was slurried in a mixture of methyl *t*-butyl ether and hexanes to afford [6-chloro-2-(4-chloro-benzoyl)-1H-indol-3-yl]-acetic acid ethyl ester (2.42g, 81%). mp = 186-188 °C. 1H NMR (300 MHz, $CDCl_3$) δ 1.27 (t, 2, J = 7.1), 3.80 (s, 2), 4.11 (q, 2, J = 7.1), 7.15 (ddd, 1, J = 8.5, 1.7, 0.5), 7.28-7.30 (m, 1), 7.48 (d, J = 8.3), 7.54-7.57 (m, 1), 7.77 (d, 2, J = 8.3), 9.16 (bs, 1). ^{13}C NMR (75 MHz, $CDCl_3$) δ 15.42, 32.29, 62.45, 113.27, 117.60, 122.99, 123.23, 127.85, 130.12, 131.77, 133.61, 137.86, 138.13, 140.10, 172.45, 188.25. IR 3305, 1732, 1618, 1323 cm^{-1} . Analysis calculated for $C_{19}H_{15}Cl_2NO_3$: C, 60.65; H, 4.02; N, 3.72. Found: C, 60.70;
15 H, 3.97; N, 3.71.
20


Example 298

[6-Chloro-2-(4-chloro-benzoyl)-1H-indol-3-yl]-acetic acid

To a solution of [6-chloro-2-(4-chloro-benzoyl)-1H-indol-3-yl]-acetic acid ethyl ester (200 mg, 0.532 mmol) in methanol (2 ml) and water (0.8 ml) was added
25 sodium hydroxide (137 mg, 3.43 mmol). The reaction mixture was stirred 24 hours, and was concentrated to a low volume. Water (4 ml) was added, the material was transferred to a separatory funnel and was washed with dichloromethane (5 ml). The aqueous layer was acidified to pH 1 with 1N hydrochloric acid and was extracted with ethyl acetate (15 ml). The organic layer was dried over magnesium sulfate, filtered,
30 and concentrate to afford [6-chloro-2-(4-chloro-benzoyl)-1H-indol-3-yl]-acetic acid (150 mg, 81%). The 1H NMR spectrum was identical with the one obtained for the

compound prepared by the method described in example 2.

The chemical structures of the compounds prepared in the Examples 1 to 292 are summarized in the following tables.

TABLE

5

	Ex.#	(X) _n	R ¹	Z	Q
10	1	6-Cl	H	ethoxy	phenyl
	2	6-Cl	H	OH	phenyl
	3	6-Cl	H	ONa	phenyl
	4	6-Cl	H	OH	2-methylphenyl
	5	6-Cl	H	OH	3-methylphenyl
	6	6-Cl	H	OH	4-methylphenyl
15	7	6-Cl	H	OH	3-chlorophenyl
	8	6-Cl	H	methoxy	4-chlorophenyl
	9	6-Cl	H	OH	4-chlorophenyl
	10	6-Cl	H	OH	3-fluorophenyl
20	11	6-Cl	H	OH	4-fluorophenyl
	12	6-Cl	H	OH	3-bromophenyl
	13	6-Cl	H	OH	4-bromophenyl
	14	6-Cl	H	OH	3-CF ₃ -phenyl
	15	6-Cl	H	OH	4-CF ₃ -phenyl
	16	6-Cl	H	OH	3,4-dichlorophenyl
25	17	4-Cl	H	OH	phenyl
	18	5-Cl	H	OH	3-methylphenyl
	19	5-Cl	H	OH	4-chlorophenyl
	20	5-Cl	H	OH	3-chlorophenyl
30	21	5-F	H	OH	4-chlorophenyl
	22	5-F	H	OH	3-chlorophenyl
	23	5-methoxy	H	OH	3-methylphenyl
	24	7-Cl	H	OH	phenyl
	25	4,5-di-Cl	H	OH	phenyl
35	26	4,6-di-Cl	H	OH	phenyl
	27	5,6-di-Cl	H	OH	phenyl

Ex.#	(X) _n	R ¹	Z	Q
5	28 6-Cl (racemic)	methyl	OH	phenyl
	29 6-Cl (less polar antipode)	methyl	OH	phenyl
	30 6-Cl (more polar antipode)	methyl	OH	phenyl
10	31 6-Cl	H	OH	4-methyl-2-pyridyl
	32 6-Cl	H	OH	5-methyl-2-pyridyl
	33 6-Cl	H	methoxy	4-chloro-2-pyridyl
	34 6-Cl	H	OH	4-chloro-2-pyridyl
	35 6-Cl	H	OH	2-pyridyl
15	36 5-Cl	H	OH	4-methyl-2-pyridyl
	37 5-Cl	H	methoxy	6-methyl-2-pyridyl
	38 5-Cl	H	OH	6-methyl-2-pyridyl
	39 6-Cl	H	OH	1-methyl-2-imidazolyl
	40 5-Cl	H	methoxy	2-thiazolyl
20	41 5-Cl	H	OH	2-thiazolyl
	42 6-Cl	H	methoxy	phenyl
	43 6-Cl	H	dimethylamino	phenyl
	44 6-Cl	H	methylamino	phenyl
	45 6-Cl	H	amino	phenyl
25	46 6-Cl	H	N-methoxy-N-methylamino	phenyl
30	47 6-Cl	H	piperidino	phenyl
	48 6-Cl	H	4-methyl-1-piperazinyl	phenyl
	49 6-Cl	H	2-cyanoethylamino	phenyl
	50 6-Cl	H	2-HO-ethylamino	phenyl
	51 6-Cl	H	morpholino	phenyl
	52 H	H	OH	4-chlorophenyl
	53 6-Cl	H	OH	2-furyl
35	54 6-Cl	H	OH	cyclohexyl
	55 6-Cl	H	OH	4-methoxyphenyl
	56 6-Cl	H	methoxy	4-methoxyphenyl
	57 6-Cl	H	methoxy	4-ethyl-2-pyridyl
	58 6-Cl	H	OH	4-ethyl-2-pyridyl
40	59 5-Cl	H	methoxy	4-ethyl-2-pyridyl
	60 5-Cl	H	OH	4-ethyl-2-pyridyl
	61 6-Cl	H	methoxy	4-isopropyl-2-pyridyl
	62 6-Cl	H	OH	4-isopropyl-2-pyridyl
	63 5-Cl	H	methoxy	4-isopropyl-2-pyridyl
45	64 5-Cl	H	OH	4-isopropyl-2-pyridyl
	65 6-Cl	H	methoxy	4-n-propyl-2-pyridyl

Ex.#	(X) _n	R ¹	Z	Q	
5	66	6-Cl	H	OH	4-n-propyl-2-pyridyl
	67	5-Cl	H	methoxy	4-n-propyl-2-pyridyl
	68	5-Cl	H	OH	4-n-propyl-2-pyridyl
	69	6-Cl	H	methoxy	4-tert-butyl-2-pyridyl
	70	6-Cl	H	OH	4-tert-butyl-2-pyridyl
10	71	5-Cl	H	methoxy	4-tert-butyl-2-pyridyl
	72	5-Cl	H	OH	4-tert-butyl-2-pyridyl
	73	6-Cl	H	methoxy	3-methyl-2-pyridyl
	74	6-Cl	H	OH	3-methyl-2-pyridyl
	75	5-Cl	H	methoxy	3-methyl-2-pyridyl
15	76	5-Cl	H	OH	3-methyl-2-pyridyl
	77	6-Cl	H	methoxy	6-methyl-2-pyridyl
	78	6-Cl	H	OH	6-methyl-2-pyridyl
	79	5-Cl	H	methoxy	5-methyl-2-pyridyl
	80	5-Cl	H	OH	5-methyl-2-pyridyl
20	81	6-Cl	H	methoxy	5-CF ₃ -2-pyridyl
	82	6-Cl	H	OH	5-CF ₃ -2-pyridyl
	83	5-Cl	H	methoxy	5-CF ₃ -2-pyridyl
	84	5-Cl	H	OH	5-CF ₃ -2-pyridyl
	85	5-Cl	H	methoxy	5-Cl-2-pyridyl
25	86	5-Cl	H	OH	5-Cl-2-pyridyl
	87	6-Cl	H	methoxy	5-Cl-2-pyridyl
	88	6-Cl	H	OH	5-Cl-2-pyridyl
	89	5-Cl	H	methoxy	4-Cl-2-pyridyl
	90	5-Cl	H	OH	4-Cl-2-pyridyl
30	91	6-Cl	H	methoxy	3-pyridyl
	92	6-Cl	H	OH	3-pyridyl
	93	6-Cl	H	methoxy	4-pyridyl
	94	6-Cl	H	OH	4-pyridyl
	95	6-Cl	H	methoxy	4-hydroxymethyl-2-pyridyl
35	96	6-Cl	H	OH	4-hydroxymethyl-2-pyridyl
	97	5-Cl	H	methoxy	4-hydroxymethyl-2-pyridyl
	98	5-Cl	H	OH	4-hydroxymethyl-2-pyridyl
	99	5-Cl	H	methoxy	3,4-dimethyl-2-pyridyl
	100	5-Cl	H	OH	3,4-dimethyl-2-pyridyl
40	101	5-Cl	H	methoxy	4,5-dimethyl-2-pyridyl
	102	5-Cl	H	OH	4,5-dimethyl-2-pyridyl
	103	6-Cl	H	methoxy	4,5-dimethyl-2-pyridyl
	104	6-Cl	H	OH	4,5-dimethyl-2-pyridyl
	105	6-Cl	H	methoxy	4-methoxy-2-pyridyl
45	106	6-Cl	H	OH	4-methoxy-2-pyridyl

Ex.#	(X) _n	R ¹	Z	Q
5	107 5-Cl	H	methoxy	4-methoxy-2-pyridyl
	108 5-Cl	H	OH	4-methoxy-2-pyridyl
	109 6-Cl	H	methoxy	3,5-dimethyl-2-pyridyl
	110 6-Cl	H	OH	3,5-dimethyl-2-pyridyl
	111 5-Cl	H	methoxy	3-F-4-ethyl-2-pyridyl
10	112 5-Cl	H	OH	3-F-4-ethyl-2-pyridyl
	113 6-Cl	H	methoxy	3-F-4-ethyl-2-pyridyl
	114 6-Cl	H	OH	3-C ₂ H ₅ O-4-ethyl-2-pyridyl
	115 6-Cl	H	methoxy	3-Cl-4-ethyl-2-pyridyl
	116 6-Cl	H	OH	3-Cl-4-ethyl-2-pyridyl
15	117 5-Cl	H	methoxy	3-Cl-4-ethyl-2-pyridyl
	118 5-Cl	H	OH	3-Cl-4-ethyl-2-pyridyl
	119 5-Cl	H	methoxy	4,6-dimethyl-2-pyridyl
	120 5-Cl	H	OH	4,6-dimethyl-2-pyridyl
	121 6-Cl	H	methoxy	4,6-dimethyl-2-pyridyl
20	122 6-Cl	H	OH	4,6-dimethyl-2-pyridyl
	123 5,6-di-Cl	H	methoxy	4-methyl-2-pyridyl
	124 5,6-di-Cl	H	OH	4-methyl-2-pyridyl
	125 5-methyl	H	methoxy	4-methyl-2-pyridyl
	126 5-methyl	H	OH	4-methyl-2-pyridyl
25	127 5-F	H	methoxy	4-methyl-2-pyridyl
	128 5-F	H	OH	4-methyl-2-pyridyl
	129 5-methoxy	H	methoxy	4-methyl-2-pyridyl
	130 5-methoxy	H	OH	4-methyl-2-pyridyl
	131 6-methoxy	H	methoxy	4-methyl-2-pyridyl
30	132 6-methoxy	H	OH	4-methyl-2-pyridyl
	133 5-ethyl	H	methoxy	4-methyl-2-pyridyl
	134 5-ethyl	H	OH	4-methyl-2-pyridyl
	135 5-ethyl	H	methoxy	4-ethyl-2-pyridyl
	136 5-ethyl	H	OH	4-ethyl-2-pyridyl
35	137 6-ethyl	H	methoxy	4-methyl-2-pyridyl
	138 6-ethyl	H	OH	4-methyl-2-pyridyl
	139 5-isopropyl	H	methoxy	4-ethyl-2-pyridyl
	140 5-isopropyl	H	OH	4-methyl-2-pyridyl
	141 6-CF ₃	H	methoxy	4-methyl-2-pyridyl
40	142 6-CF ₃	H	OH	4-methyl-2-pyridyl
	143 5-tert-butyl	H	methoxy	4-methyl-2-pyridyl
	144 5-tert-butyl	H	OH	4-methyl-2-pyridyl
	145 5-CF ₃ O	H	methoxy	4-methyl-2-pyridyl
	146 5-CF ₃ O	H	OH	4-methyl-2-pyridyl
45	147 5-CF ₃ O	H	methoxy	4-ethyl-2-pyridyl
	148 5-CF ₃ O	H	OH	4-ethyl-2-pyridyl

Ex.#	(X) _n	R ¹	Z	Q
5	149 6-methyl	H	methoxy	4-methyl-2-pyridyl
	150 6-methyl	H	OH	4-methyl-2-pyridyl
	151 5-CF ₃	H	methoxy	4-methyl-2-pyridyl
	152 5-CF ₃	H	OH	4-methyl-2-pyridyl
	153 5-CF ₃	H	methoxy	4-ethyl-2-pyridyl
10	154 5-CF ₃	H	OH	4-ethyl-2-pyridyl
	155 H	H	methoxy	phenyl
	156 H	H	OH	phenyl
	157 6-methyl	H	methoxy	4-chlorophenyl
	158 6-methyl	H	OH	4-chlorophenyl
15	159 5-methyl	H	OH	4-chlorophenyl
	160 6-methoxy	H	methoxy	4-chlorophenyl
	161 6-methoxy	H	OH	4-chlorophenyl
	162 6-CF ₃	H	OH	4-chlorophenyl
	163 5-ethyl	H	methoxy	4-chlorophenyl
20	164 5-ethyl	H	OH	4-chlorophenyl
	165 5-methoxy	H	methoxy	4-chlorophenyl
	166 5-methoxy	H	OH	4-chlorophenyl
	167 5-isopropyl	H	methoxy	4-chlorophenyl
	168 5-isopropyl	H	OH	4-chlorophenyl
25	169 5-CF ₃	H	methoxy	4-chlorophenyl
	170 5-CF ₃	H	OH	4-chlorophenyl
	171 5-CF ₃ O	H	methoxy	4-chlorophenyl
	172 5-CF ₃ O	H	OH	4-chlorophenyl
	173 6-Cl	H	methoxy	2-methoxyphenyl
30	174 6-Cl	H	OH	2-methoxyphenyl
	175 6-Cl	H	methoxy	3-methoxyphenyl
	176 6-Cl	H	OH	3-methoxyphenyl
	177 6-Cl	H	methoxy	3-benzyloxyphenyl
	178 6-Cl	H	OH	3-benzyloxyphenyl
35	179 6-Cl	H	methoxy	3-hydroxyphenyl
	180 6-Cl	H	OH	3-hydroxyphenyl
	181 6-Cl	H	methoxy	4-benzyloxyphenyl
	182 6-Cl	H	OH	4-benzyloxyphenyl
	183 6-Cl	H	methoxy	4-hydroxyphenyl
40	184 6-Cl	H	OH	4-hydroxyphenyl
	185 6-Cl	H	methoxy	4-isopropoxyphenyl
	186 6-Cl	H	OH	4-isopropoxyphenyl
	187 6-Cl	H	methoxy	4-biphenyl
	188 6-Cl	H	OH	4-biphenyl
45	189 6-Cl	H	methoxy	4-CF ₃ O-phenyl
	190 6-Cl	H	OH	4-CF ₃ O-phenyl


Ex.#	(X) _n	R ¹	Z	Q
5	191 5-Cl	H	methoxy	4-CF ₃ O-phenyl
	192 5-Cl	H	OH	4-CF ₃ O-phenyl
	193 5-Cl	H	methoxy	4-methoxyphenyl
	194 5-Cl	H	OH	4-methoxyphenyl
	195 6-Cl	H	methoxy	4-nitrophenyl
10	196 6-Cl	H	OH	4-nitrophenyl
	197 6-Cl	H	methoxy	4-methyl-S(O) ₂ -phenyl
	198 6-Cl	H	OH	4-methyl-S(O) ₂ -phenyl
	199 6-Cl	H	methoxy	4-methyl-S(O) ₂ -NH-phenyl
	200 6-Cl	H	OH	4-methyl-S(O) ₂ -NH-phenyl
15	201 6-Cl	H	OH	2-chlorophenyl
	202 6-Cl	H	OH	2,4-dichlorophenyl
	203 6-Cl	H	methoxy	3-F-4-Cl-phenyl
	204 6-Cl	H	OH	3-F-4-Cl-phenyl
	205 6-Cl	H	methoxy	4-cyanophenyl
20	206 6-Cl	H	methoxy	4-bromophenyl
	207 6-Cl	H	methoxy	4-(2-thienyl)phenyl
	208 6-Cl	H	OH	4-(2-thienyl)phenyl
	209 6-Cl	H	methoxy	4-(2-furyl)phenyl
	210 6-Cl	H	OH	4-(2-furyl)phenyl
25	211 6-Cl	H	methoxy	4-(3-pyridyl)phenyl
	212 6-Cl	H	OH	4-(3-pyridyl)phenyl
	213 6-Cl	H	methoxy	4-(2-thiazolyl)phenyl
	214 6-Cl	H	OH	4-(2-thiazolyl)phenyl
	215 6-Cl	H	methoxy	3-bromophenyl
30	216 6-Cl	H	methoxy	3-(2-furyl)phenyl
	217 6-Cl	H	OH	3-(2-furyl)phenyl
	218 6-Cl	methyl	methoxy	4-chlorophenyl
	219 6-Cl	methyl	OH	4-chlorophenyl
	220 5-Cl	H	methoxy	isoquinolin-3-yl
35	221 5-Cl	H	OH	isoquinolin-3-yl
	222 6-Cl	H	methoxy	isoquinolin-3-yl
	223 6-Cl	H	OH	isoquinolin-3-yl
	224 5-Cl	H	methoxy	5-methyl-3-isoxazolyl
	225 5-Cl	H	OH	5-methyl-3-isoxazolyl
40	226 6-Cl	H	methoxy	5-methyl-3-isoxazolyl
	227 6-Cl	H	OH	5-methyl-3-isoxazolyl
	228 5-Cl	H	methoxy	4-methyl-5-(1,2,3-thiadiazolyl)
	229 5-Cl	H	OH	4-methyl-5-(1,2,3-thiadiazolyl)
	230 6-Cl	H	methoxy	4-methyl-5-(1,2,3-thiadiazolyl)
45	231 6-Cl	H	OH	4-methyl-5-(1,2,3-thiadiazolyl)
	232 5-Cl	H	methoxy	5-methyl-2-thiazolyl

Ex.#	(X) _n	R ¹	Z	Q
5	233 5-Cl	H	OH	5-methyl-2-thiazolyl
	234 6-Cl	H	methoxy	5-methyl-2-thiazolyl
	235 6-Cl	H	OH	5-methyl-2-thiazolyl
	236 6-Cl	H	OH	2-thienyl
	237 6-Cl	H	methoxy	3-(HO)(CH ₃) ₂ C-2-furyl
10	238 6-Cl	H	OH	3-(HO)(CH ₃) ₂ C-2-furyl
	239 6-Cl	H	methoxy	3-methoxymethyl-2-furyl
	240 6-Cl	H	OH	3-methoxymethyl-2-furyl
	241 6-Cl	H	OH	1-methyl-2-imidazolyl
	242 6-Cl	H	methoxy	1-methyl-2-imidazolyl
15	243 5-Cl	H	methoxy	1-methyl-2-imidazolyl
	244 5-Cl	H	OH	1-methyl-2-imidazolyl
	245 5-Cl	H	methoxy	2-imidazolyl
	246 5-Cl	H	OH	2-imidazolyl
	247 6-Cl	H	methoxy	2-imidazolyl
20	248 6-Cl	H	OH	2-imidazolyl
	249 5-Cl	H	methoxy	4-methyl-2-thiazolyl
	250 5-Cl	H	OH	4-methyl-2-thiazolyl
	251 5-Cl	H	methoxy	1-methyl-2-pyrrolyl
	252 5-Cl	H	OH	1-methyl-2-pyrrolyl
25	253 5-Cl	H	methoxy	2-methyl-4-thiazolyl
	254 5-Cl	H	OH	2-methyl-4-thiazolyl
	255 5-Cl	H	methoxy	5-thiazolyl
	256 5-Cl	H	OH	5-thiazolyl
	257 6-Cl	H	methoxy	4-methyl-2-thiazolyl
30	258 6-Cl	H	OH	4-methyl-2-thiazolyl
	259 5-Cl	H	methoxy	3-carboxyl-5-isoxazolyl
	260 5-Cl	H	OH	3-carboxyl-5-isoxazolyl
	261 6-Cl	H	methoxy	cyclopropyl
	262 6-Cl	H	OH	cyclopropyl
35	263 6-Cl	H	methoxy	cyclobutyl
	264 6-Cl	H	OH	cyclobutyl
	265 5-tert-butyl	H	methoxy	4-chlorophenyl
	266 5-tert-butyl	H	OH	4-chlorophenyl
	267 6-Cl	H	dimethylamino	4-methyl-2-pyridyl
40	268 6-Cl	H	methylamino	4-methyl-2-pyridyl
	269 5-Cl	H	HO-(CH ₂) ₂ -NH-	4-methyl-2-pyridyl
	270 5-Cl	H	methoxyamino	4-methyl-2-pyridyl
	271 5-Cl	H	1-piperazinyl	4-methyl-2-pyridyl
	272 5-Cl	H	H ₂ N-(CH ₂) ₂ -NH-	4-methyl-2-pyridyl
	273 5-Cl	H	3-amino-1-pyrrolidinyl	4-methyl-2-pyridyl

Ex.#	(X) _n	R ¹	Z	Q	
5	274	5-F,6-Cl	H	methoxy	4-chlorophenyl
	275	5-F,6-Cl	H	OH	4-chlorophenyl
	276	5-F,6-Cl	H	methoxy	4-methyl-2-pyridyl
	277	5-F,6-Cl	H	OH	4-methyl-2-pyridyl
	278	6-Cl	H	methoxy	4-(HO)(H ₃ C)CH-2-pyridyl
10	279	6-Cl	H	OH	4-(HO)(H ₃ C)CH-2-pyridyl
	280	6-Cl	H	OH	4-ethyl-3-F-2-pyridyl
	281	6-Cl	H	OH	2-nitrophenyl
	282	6-Cl	H	OH	2,4-dimethoxyphenyl
	283	6-Cl	H	OH	4-CHF ₂ O-phenyl
15	284	6-Cl	H	OH	2,5-dimethoxyphenyl
	285	5-acetyl	H	methoxy	4-Cl-phenyl
	286	5-acetyl	H	OH	4-Cl-phenyl
	287	6-F	H	methoxy	4-methyl-2-pyridyl
	288	6-F	H	OH	4-methyl-2-pyridyl
20	289	6-F	H	methoxy	4-Cl-phenyl
	290	6-F	H	OH	4-Cl-phenyl
	291	5-CH ₃ S-	H	methoxy	4-methyl-2-pyridyl
	292	5-CH ₃ S-	H	OH	4-methyl-2-pyridyl

CLAIMS


1. A compound of the following formula:



(I)

- 5 or the pharmaceutically acceptable salts thereof wherein

Z is OH, C₁₋₆ alkoxy, -NR²R³ or a group of the formula (II) or (III):



(II)

(III)

wherein r is 1, 2, 3 or 4, Y is a direct bond, O, S or NR⁴, and W is OH or -NR²R³;

- 10 Q is selected from the following:

- (a) phenyl optionally substituted with one, two or three substituents independently selected from

(a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,

(a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋

4 alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),

(c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);

(d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and

(e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R¹ is hydrogen, C₁₋₄ alkyl or halo;

R² and R³ are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

R⁴ is hydrogen or C₁₋₄ alkyl;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino,

C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄

alkylsulfonylamino and C₃₋₇ cycloalkyl; and

n is 0, 1, 2, 3 or 4.

2. A compound according to claim 1, wherein

Z is OH, C₁₋₆ alkoxy, dimethylamino, methylamino, amino, N-methoxy-N-methylamino, 2-cyanoethylamino, 2-hydroxyethylamino, pyrrolidinyl, piperidino, piperazinyl, N-methyl-piperazinyl, morpholino, methoxyamino, piperazynyl, aminopyrrolidinyl or aminoethylamino.

3. A compound according to claim 2, wherein

Z is OH or C₁₋₆ alkoxy; and

10 Q is selected from the following:

(a) phenyl optionally substituted with one, two or three substituents independently selected from

15 (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,

20 (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

25 (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

30 (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-

substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

- 5 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- 10 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- 15 (d) C₃₋₇ cycloalkyl selected from cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl, and the said cycloalkyl being optionally substituted with one substituent selected from OH, methyl, ethyl, propyl, F, Cl and CF₃; and
- 20 (e) a benzo-fused heterocycle selected from quinolyl, isoquinolyl, cinnolyl, quinoxalyl, benzoimidazolyl, benzothiazolyl, benzoxazolyl, benzofuranyl, benzothiophenyl and indolyl, and the benzo-fused heterocycle being optionally substituted with one, two, or three substituents independently selected from the group (a-1).

4. A compound according to claim 3, wherein

Q is selected from the following:

- 25 (a) phenyl optionally substituted with one, two or three substituents independently selected from
- (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,
- 30 (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally

substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

5 (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

10 (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

15 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),

20 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);

25 (d) cyclopropyl, cyclobutyl and cyclohexyl; and

(e) quinolyl or isoquinolyl, and said quinolyl or isoquinolyl being optionally substituted with one substituent selected from the group halo, C₁₋₄ alkyl, NH₂, OH, C₁₋₄ alkoxy and C₁₋₄ halo-substituted alkyl.

5. A compound according to claim 4, wherein

30 Z is OH, C₁₋₆ alkoxy;

Q is selected from the following:

- (a) phenyl optionally substituted with one, two or three substituents independently selected from
- 5 (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,
- 10 (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 15 (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 20 (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 25 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- 30 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three

substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4); and

(e) isoquinolyl;

R^1 is hydrogen or C_{1-4} alkyl;

5 R^2 and R^3 are independently H or methyl;

X is independently selected from halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- (C_{1-4}) amino, C_{1-4} alkylamino, CN, HO- (C_{1-4}) alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $-NH_2S(O)_2NR^2NR^3$, acetyl, $-COOR^4$, C_{1-4} alkylsulfonylamino and C_{3-7}

10 cycloalkyl; and

n is 0, 1, 2, or 3.

6. A compound according to claim 5, wherein

Z is OH, methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, iso-butoxy or tert-butoxy;

15 Q is selected from the following:

(a) phenyl optionally substituted with one or two substituents independently selected from

(a-1) halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, HO- (C_{1-4}) alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, $-COOH$, C_{1-4} alkylsulfonylamino, nitro, C_{1-4} alkylsulfonyl and cyano,

(a-2) phenyl or benzyloxy, and the phenyl or phenyl moiety of benzyloxy being optionally substituted with one substituent selected from C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, halo, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy and NH_2 ,

(a-3) 5-membered monocyclic aromatic group selected from imidazolyl, thiazolyl, furyl, thienyl, pyrrolyl, tetrazolyl, triazolyl, oxazolyl, isoxazolyl, thiadiazolyl and pyrazolyl, and the 5-membered monocyclic aromatic group optionally being substituted with one substituent selected from C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, halo, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy and NH_2 ,

- (a-4) 6-membered monocyclic aromatic group selected from pyridyl, pyrazinyl, pyrimidinyl and pyridazinyl, and the 6-membered monocyclic aromatic group optionally being substituted with one substituent selected from C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, halo, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy and NH₂,
- (b) a 6-membered monocyclic aromatic group selected from pyridyl, pyrazinyl, pyrimidinyl and pyridazinyl, and said monocyclic aromatic group being optionally substituted with one or two substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- (c) a 5-membered monocyclic aromatic group selected from imidazolyl, thiazolyl, furyl, thienyl, pyrrolyl, tetrazolyl, triazolyl, oxazolyl, isoxazolyl, thiadiazolyl and pyrazolyl, and said monocyclic aromatic group being optionally substituted with one or two substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- R¹ is hydrogen, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl or tert-butyl;
- X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl and aminosulfonyl; and
- n is 0, 1, 2, or 3.

7. A compound according to claim 6, wherein

Z is OH, methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, iso-butoxy or tert-butoxy;

Q is selected from the following:

- (a) phenyl optionally substituted with one or two substituents independently selected from
- (a-1) fluoro, chloro, bromo, iodo, methyl, ethyl, propyl, butyl, CH₂F, CHF₂, CF₃, methoxy, ethoxy, n-propoxy, n-butoxy, isopropoxy, CH₂F-O-, CHF₂-O-, CF₃-O-, methylthio, ethylthio, hydroxymethyl, methoxymethyl, methoxyethyl, ethoxymethyl, hydroxy, nitro, methylsulfonyl, cyano, (HO)(H₃C)₂C-, acetyl and

methylsulfonylamino,

(a-2) phenyl or benzyloxy, and the phenyl or phenyl moiety of benzyloxy being optionally substituted with one substituent selected from methyl, ethyl, propyl, CF₃, F, Cl, OH, methoxy, ethoxy and NH₂,

5 (a-3) 5-membered monocyclic aromatic group selected from furyl, thienyl and pyrrolyl, and the 5-membered monocyclic aromatic group optionally being substituted with one substituent selected from methyl, ethyl, propyl, CF₃, F, Cl, OH, methoxy, ethoxy and NH₂,

10 (a-4) pyridyl optionally substituted with one substituent selected from methyl, ethyl, propyl, CF₃, F, Cl, OH, methoxy, ethoxy and NH₂,

(b) pyridyl optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-1), (a-2), (a-3) and (a-4),

15 (c) imidazolyl, thiazolyl, furyl, thienyl, isoxazolyl, 1,2,3-thiadiazolyl or pyrrolyl, and said imidazolyl, thiazolyl, furyl, thienyl, isoxazolyl, 1,2,3-thiadiazolyl or pyrrolyl being optionally substituted with one or two substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);

R¹ is hydrogen, methyl, ethyl, n-propyl, iso-propyl;

20 **X** is independently selected from fluoro, chloro, bromo, methyl, ethyl, propyl, butyl, CH₂F, CHF₂, CF₃, methoxy, CF₃O or ethoxy; and
n is 0, 1 or 2.

8. A compound according to claim 7, wherein

25 **Z** is OH, ethoxy or methoxy; **Q** is phenyl, chlorophenyl, fluorophenyl, bromophenyl, methylphenyl, methoxyphenyl, (furyl)phenyl, trifluoromethylphenyl, trifluoromethoxyphenyl, pyridyl, methylpyridyl, ethylpyridyl, propylpyridyl, dimethylpyridyl, chloropyridyl, fluoropyridyl, trifluoromethylpyridyl, methoxypyridyl, (ethyl)(ethoxy)pyridyl, (chloro)(ethyl)pyridyl, thiazolyl, methylthiazolyl, furyl, methoxymethylfuryl, isoquinolyl, cyclohexyl, methoxyphenyl, (fluoro)(ethyl)pyridyl,
30 dimethylpyridyl or (ethoxy)(ethyl)pyridyl;

R¹ is hydrogen; **X** is fluoro, chloro, methyl, ethyl, isopropyl, tert-butyl, CF₃ or

methoxy; and **n** is 1 or 2.

9. A compound according to claim 8, wherein

Z is OH, ethoxy or methoxy; **Q** is phenyl, chlorophenyl, pyridyl, methylpyridyl, ethylpyridyl, propylpyridyl or chloropyridyl; **R**¹ is hydrogen; **X** is fluoro, chloro,

5 methyl or CF₃; and **n** is 1 or 2.

10. A compound according to claim 1 selected from

ethyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;

(2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;

(2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid, sodium salt;

10 [6-chloro-2-(2-methylbenzoyl)-1H-indol-3-yl]acetic acid;

[6-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;

[6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]acetic acid;

[6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;

methyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate;

15 [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;

[6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;

[6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]acetic acid;

[2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;

[2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;

20 [6-chloro-2-(3-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;

[6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;

[6-chloro-2-(3,4-dichlorobenzoyl)-1H-indol-3-yl]acetic acid;

(2-benzoyl-4-chloro-1H-indol-3-yl)acetic acid;

[5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;

25 [5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;

[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;

[2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;

[2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;

[5-methoxy-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;

30 (2-benzoyl-7-chloro-1H-indol-3-yl)acetic acid;

(2-benzoyl-4,5-dichloro-1H-indol-3-yl)acetic acid;

- (2-benzoyl-4,6-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
dl-2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
less polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
5 more polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 [6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
15 methyl [5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
(2-benzoyl-6-chloro-1H-indol-3-yl)-*N,N*-dimethylacetamide;
(2-benzoyl-6-chloro-1H-indol-3-yl)-*N*-methylacetamide;
20 (2-benzoyl-6-chloro-1H-indol-3-yl)acetamide;
(2-benzoyl-6-chloro-1H-indol-3-yl)-*N*-methoxy-*N*-methylacetamide;
2-(2-benzoyl-6-chloro-1H-indol-3-yl)-1-piperidino-1-ethanone;
2-(2-benzoyl-6-chloro-1H-indol-3-yl)-1-(4-methyl-1-piperazinyl)-1-ethanone;
(2-benzoyl-6-chloro-1H-indol-3-yl)-*N*-(2-cyanoethyl)acetamide;
25 (2-benzoyl-6-chloro-1H-indol-3-yl)-*N*-(2-hydroxyethyl)acetamide;
2-(2-benzoyl-6-chloro-1H-indol-3-yl)-1-morpholino-1-ethanone;
[2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2-furylcarbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]acetic acid;
30 methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;

- methyl [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 5 methyl [6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- methyl [6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
- 10 [6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-*tert*-butylpyridine-2-carbonyl)-6-chloro-1H-indol-3-yl]acetate;
[2-(4-*tert*-butylpyridine-2-carbonyl)-6-chloro-1H-indol-3-yl]acetic acid;
- 15 methyl [2-(4-*tert*-butylpyridine-2-carbonyl)-5-chloro-1H-indol-3-yl]acetate;
[2-(4-*tert*-butylpyridine-2-carbonyl)-5-chloro-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
- 20 [5-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 25 methyl [6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
- methyl [5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
- 30 [5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;

- [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(pyridine-3-carbonyl)-1H-indol-3-yl]acetate;
5 [6-chloro-2-(pyridine-3-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(pyridine-4-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(pyridine-4-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
10 methyl [5-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[5-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(3,4-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(3,4-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
15 [5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 methyl [5-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
25 [5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-ethoxy-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 methyl [5-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- methyl [5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 5 methyl [5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- methyl [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
10 [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 15 methyl [5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-ethyl-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-ethyl-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
- 20 [6-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
- 25 methyl [5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetate;
[2-(4-methyl-2-pyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
methyl [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetate;
- 30 [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
methyl [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;

- [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
5 [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl (2-benzoyl-1H-indol-3-yl)acetate;
(2-benzoyl-1H-indol-3-yl)acetic acid;
methyl [2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetate;
[2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetic acid;
10 [2-(4-chlorobenzoyl)-5-methyl-1H-indol-3-yl]acetic acid;
methyl [6-methoxy-2-(4-chlorobenzoyl)-1H-indol-3-yl] acetate;
[6-methoxy-2-(4-chlorobenzoyl)-1H-indol-3-yl] acetic acid;
[2-(4-chlorobenzoyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-5-ethyl-1H-indol-3-yl]acetate;
15 [2-(4-chlorobenzoyl)-5-ethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-5-methoxy-1H-indol-3-yl]acetate;
[2-(4-chlorobenzoyl)-5-methoxy-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-5-isopropyl-1H-indol-3-yl]acetate;
[2-(4-chlorobenzoyl)-5-isopropyl-1H-indol-3-yl]acetic acid;
20 methyl [2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-5-trifluoromethoxy-1H-indol-3-yl] acetate;
[2-(4-chlorobenzoyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(2-methoxybenzoyl)-1H-indol-3-yl]acetate;
25 [6-chloro-2-(2-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-benzyloxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-benzyloxybenzoyl)-1H-indol-3-yl]acetic acid;
30 methyl [6-chloro-2-(3-hydroxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-hydroxybenzoyl)-1H-indol-3-yl]acetic acid;

- methyl [6-chloro-2-(4-benzyloxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-benzyloxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-hydroxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-hydroxybenzoyl)-1H-indol-3-yl]acetic acid;
- 5 methyl [6-chloro-2-(4-isopropoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-isopropoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-phenylbenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-phenylbenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetate;
- 10 [6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
- 15 methyl [6-chloro-2-(4-nitrobenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-nitrobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[(4-methylsulfonyl)benzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[(4-methylsulfonyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(methylsulfonylamino)benzoyl]-1H-indol-3-yl]acetate;
- 20 [6-chloro-2-[4-(methylsulfonylamino)benzoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2,4-dichlorobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chloro-3-fluorobenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chloro-3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
- 25 methyl [6-chloro-2-(4-cyanobenzoyl)-1H-indol-3-yl]acetate;
methyl [6-chloro-2-[4-bromobenzoyl]-1H-indol-3-yl]acetate;
methyl [6-chloro-2-[4-(2-thienyl)benzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(2-thienyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetate;
- 30 [6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(3-pyridyl)benzoyl]-1H-indol-3-yl]acetate;

- [6-chloro-2-[4-(3-pyridyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(2-thiazolyl)benzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(2-thiazolyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-bromobenzoyl)-1H-indol-3-yl]acetate;
5 methyl [6-chloro-2-[3-(2-furyl)benzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[3-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl *dl*-2-[6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]propionate;
dl-2-[2-(4-chlorobenzoyl)-6-chloro-1H-indol-3-yl]propionic acid;
methyl [5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
10 [5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetic acid;
15 methyl [6-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methyl-1,2,3-thiadiazole-5-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methyl-1,2,3-thiadiazole-5-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methyl-1,2,3-thiadiazole-5-carbonyl)-1H-indol-3-yl]acetate;
20 [6-chloro-2-(4-methyl-1,2,3-thiadiazole-5-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
25 [6-chloro-2-(2-thienyl)carbonylindol-3-yl]acetic acid;
methyl [6-chloro-2-[3-(1-hydroxy-1-methylethyl)-2-furoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[3-(1-hydroxy-1-methylethyl)-2-furoyl]-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetic acid;
30 [6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetate;

- methyl [5-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 5 methyl [6-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(1-methylpyrrole-2-carbonyl)-1H-indol-3-yl]acetate;
- 10 [5-chloro-2-(1-methylpyrrole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(thiazole-5-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(thiazole-5-carbonyl)-1H-indol-3-yl]acetic acid;
- 15 methyl [6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-[3-(ethoxycarbonyl)isoxazole-5-carbonyl]-1H-indol-3-yl]acetate;
[5-chloro-2-[3-(carboxy)isoxazole-5-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-cyclopropanecarbonyl-1H-indol-3-yl]acetate;
- 20 [6-chloro-2-cyclopropanecarbonyl-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-cyclobutanecarbonyl-1H-indol-3-yl]acetate;
[6-chloro-2-cyclobutanecarbonyl-1H-indol-3-yl]acetic acid;
methyl [5-(*tert*-butyl)-2-(4-chlorobenzoyl)-1H-indol-3-yl] acetate;
[5-(*tert*-butyl)-2-(4-chlorobenzoyl)-1H-indol-3-yl] acetic acid;
- 25 [6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-*N,N*-dimethylacetamide;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-*N*-methylacetamide;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-*N*-(2-hydroxyethyl)acetamide;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-*N*-methoxyacetamide;
- 30 2-[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-1-piperazinyl-1-ethanone;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-*N*-(2-aminoethyl)acetamide;

- 2-[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]-1-(3-amino-1-pyrrolidinyl)-1-ethanone;
[6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
methyl [6-chloro-5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
5 [6-chloro-5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2-nitrobenzoyl)-1H-indol-3-yl]acetic acid;
10 [6-chloro-2-(2,4-dimethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-difluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2,5-dimethoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [5-acetyl-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate;
[5-acetyl-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
15 methyl [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetate;
methyl [6-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-fluoro-2-(4-chlorobenzoyl)-1H-indol-3-yl] acetate;
[6-fluoro-2-(4-chlorobenzoyl) -1H-indol-3-yl]acetic acid;
20 [2-(4-methylpyridine-2-carbonyl)-5-methylthio-1H-indol-3-yl]acetic acid;
[2-(4-methylpyridine-2-carbonyl)-5-methylthio-1H-indol-3-yl]acetic acid, and a salt thereof.

11. A compound according to claim 10 selected from
ethyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
25 (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid, sodium salt;
[6-chloro-2-(2-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]acetic acid;
30 [6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate;

- [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
5 [2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3,4-dichlorobenzoyl)-1H-indol-3-yl]acetic acid;
(2-benzoyl-4-chloro-1H-indol-3-yl)acetic acid;
10 [5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
[2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
15 [5-methoxy-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
(2-benzoyl-7-chloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-4,5-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-4,6-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
20 *dl*-2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
less polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
more polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
25 methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
30 [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;

- methyl [5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
[2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
- 5 [6-chloro-2-(2-furylcarbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
- 10 [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 15 [5-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-*tert*-butylpyridine-2-carbonyl)-6-chloro-1H-indol-3-yl]acetic acid;
[2-(4-*tert*-butylpyridine-2-carbonyl)-5-chloro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 20 [6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
- 25 [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-4-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
- 30 [5-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3,4-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- [5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 [6-chloro-2-(3,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-ethoxy-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 [5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
15 [5-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-ethyl-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 [5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
[5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-methyl-2-pyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
[2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
25 [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
(2-benzoyl-1H-indol-3-yl)acetic acid;
[2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetic acid;
30 [2-(4-chlorobenzoyl)-5-methyl-1H-indol-3-yl]acetic acid;
[6-methoxy-2-(4-chlorobenzoyl)-1H-indol-3-yl] acetic acid;

- [2-(4-chlorobenzoyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-ethyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-methoxy-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-isopropyl-1H-indol-3-yl]acetic acid;
5 [2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-benzyloxybenzoyl)-1H-indol-3-yl]acetic acid;
10 [6-chloro-2-(3-hydroxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-benzyloxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-hydroxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-isopropoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-phenylbenzoyl)-1H-indol-3-yl]acetic acid;
15 [6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-nitrobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-[(4-methylsulfonyl)benzoyl]-1H-indol-3-yl]acetic acid;
20 [6-chloro-2-[4-(methylsulfonylamino)benzoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2,4-dichlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chloro-3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-cyanobenzoyl)-1H-indol-3-yl]acetate;
25 methyl [6-chloro-2-[4-bromobenzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(2-thienyl)benzoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-[4-(3-pyridyl)benzoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-[4-(2-thiazolyl)benzoyl]-1H-indol-3-yl]acetic acid;
30 methyl [6-chloro-2-(3-bromobenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-[3-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;

- dl*-2-[2-(4-chlorobenzoyl)-6-chloro-1H-indol-3-yl]propionic acid;
[5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetic acid;
5 [6-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methyl-1,2,3-thiadiazole-5-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methyl-1,2,3-thiadiazole-5-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 [6-chloro-2-(2-thienyl)carbonylindol-3-yl]acetic acid;
[6-chloro-2-[3-(1-hydroxy-1-methylethyl)-2-furoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetate;
15 [5-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(imidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(1-methylpyrrole-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 [5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(thiazole-5-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-[3-(carboxy)isoxazole-5-carbonyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-cyclopropanecarbonyl-1H-indol-3-yl]acetic acid;
25 [6-chloro-2-cyclobutanecarbonyl-1H-indol-3-yl]acetic acid;
[5-(*tert*-butyl)-2-(4-chlorobenzoyl)-1H-indol-3-yl] acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
[6-chloro-5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
30 [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-ethyl-3-fluoro-pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- [6-chloro-2-(2-nitrobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2,4-dimethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-difluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(2,5-dimethoxybenzoyl)-1H-indol-3-yl]acetic acid;
5 [5-acetyl-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid, and a salt thereof.

12. A compound according to claim 10 selected from

- ethyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
(2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
[6-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
10 [6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
15 [6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
20 (2-benzoyl-4-chloro-1H-indol-3-yl)acetic acid;
[5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
25 [2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
(2-benzoyl-4,5-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-4,6-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
dl-2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
30 less polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
more polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;

- [6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 [6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 methyl [5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
[6-chloro-2-(2-furylcarbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]acetic acid;
15 methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
20 [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
25 methyl [5-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-*tert*-butylpyridine-2-carbonyl)-6-chloro-1H-indol-3-yl]acetate;
[2-(4-*tert*-butylpyridine-2-carbonyl)-6-chloro-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
30 [6-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;

- [6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
5 [6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 methyl [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
15 [5-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 methyl [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
25 [6-chloro-2-(3,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-ethoxy-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 methyl [6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- methyl [5-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 methyl [6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
10 [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
15 methyl [6-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-ethyl-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
20 [5-ethyl-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
25 methyl [2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetate;
30 [2-(4-methyl-2-pyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
methyl [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;

- [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
5 [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetate;
[2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-methyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
10 [2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-chlorobenzoyl)-5-trifluoromethoxy-1H-indol-3-yl] acetate;
[2-(4-chlorobenzoyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
15 methyl [6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
20 [5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
25 methyl [6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
30 [5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;

- [6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 methyl [5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
10 [6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
15 methyl [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetate, and a salt thereof.

13. A compound according to claim 10 selected from
ethyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
(2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
[6-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
20 [6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
25 [6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
30 (2-benzoyl-4-chloro-1H-indol-3-yl)acetic acid;
[5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;

- [5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
[2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
5 (2-benzoyl-4,5-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-4,6-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
dl-2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
less polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
10 more polar antipode, 2-(2-benzoyl-6-chloro-1H-indol-3-yl)propanoic acid;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
15 [6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 methyl [5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
[6-chloro-2-(2-furylcarbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]acetic acid;
25 methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 [6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- [5-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-*tert*-butylpyridine-2-carbonyl)-6-chloro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 [5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 [5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-[4-(hydroxymethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
15 [5-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-ethoxy-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 [5-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
25 [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-methoxy-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-ethyl-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 [6-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- [2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
[5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-methyl-2-pyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
[6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-6-methyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-methyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
10 [2-(4-chlorobenzoyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
15 [6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylisoxazole-3-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 [6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(1-methylimidazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetic acid;
25 [6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
2-{6-chloro-2-[(4-ethyl-3-fluoro-2-pyridinyl)carbonyl]-1H-indol-3-yl}acetic acid;
30 methyl [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetate, and a salt thereof.

14. A compound according to claim 10 selected from

- (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
[6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate;
5 [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
10 [6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
15 [2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
(2-benzoyl-4,5-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-4,6-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 [6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
25 methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
30 [6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;

- [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
5 [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 methyl [6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
15 [6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 methyl [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
25 [5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 methyl [5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- methyl [6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-ethoxy-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 methyl [5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
10 [5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
15 methyl [5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetate;
20 [2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetate;
[2-(4-methyl-2-pyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
25 methyl [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
30 [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-methyl-1H-indol-3-yl]acetic acid;

- methyl [2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
5 methyl [5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetate;
10 [6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
15 methyl [5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetate;
20 [6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetic acid;
25 methyl [6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetate, and a salt thereof.
30 15. A compound according to claim 10 selected from
(2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;

- [6-chloro-2-(4-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
5 [6-chloro-2-(3-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-fluorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(3-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[2-(4-bromobenzoyl)-6-chloro-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-trifluoromethylbenzoyl)-1H-indol-3-yl]acetic acid;
10 [5-chloro-2-(3-methylbenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
[2-(3-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
15 (2-benzoyl-4,5-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-4,6-dichloro-1H-indol-3-yl)acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
25 [5-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(thiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl (2-benzoyl-6-chloro-1H-indol-3-yl)acetate;
[6-chloro-2-(cyclohexanecarbonyl)-1H-indol-3-yl]acetic acid;
30 methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;

- methyl [6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 [6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(6-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-[5-(trifluoromethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
10 [5-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4,5-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
15 [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-ethoxy-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(3-chloro-4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 [6-chloro-2-(4,6-dimethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5,6-dichloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-ethyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
25 [5-isopropyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-methylpyridine-2-carbonyl)-6-trifluoromethyl-1H-indol-3-yl]acetic acid;
[5-*tert*-butyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-methyl-2-pyridine-2-carbonyl)-5-trifluoromethoxy-1H-indol-3-yl]acetic acid;
[6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 [2-(4-methylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;

- [2-(4-chlorobenzoyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-trifluoromethoxybenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
5 [6-chloro-2-[4-(2-furyl)benzoyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 [6-chloro-2-[3-methoxymethyl-2-furoyl]-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(2-methylimidazole-4-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-methylthiazole-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
15 [6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetate;
[6-chloro-5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid,
20 and a salt thereof.

16. A compound according to claim 1 selected from
(2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
[6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
25 [5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 [6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;

- [6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
5 methyl [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
10 [6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetate;
[2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
15 methyl [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetate;
20 [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
25 methyl [5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
30 methyl [6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetate;
[6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;

- [6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetate;
[6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate;
5 [6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
methyl [5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetate;
[5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid (cj-020,099);
methyl [6-chloro-5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetate,
10 and a salt thereof.

17. A compound according to claim 10 selected from

- (2-benzoyl-6-chloro-1H-indol-3-yl)acetic acid;
[6-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
15 [5-chloro-2-(4-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(3-chlorobenzoyl)-1H-indol-3-yl]acetic acid;
[2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid;
(2-benzoyl-5,6-dichloro-1H-indol-3-yl)acetic acid;
[6-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
20 [6-chloro-2-(5-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(pyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
25 [5-chloro-2-(4-ethylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-isopropylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-propylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[2-(4-ethylpyridine-2-carbonyl)-5-trifluoromethyl-1H-indol-3-yl]acetic acid;
30 [6-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;

- [6-chloro-2-(4-methoxypyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-fluoro-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[5-chloro-2-(4-methoxybenzoyl)-1H-indol-3-yl]acetic acid;
5 [6-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-[4-(1-hydroxyethyl)pyridine-2-carbonyl]-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-ethyl-3-fluoropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(5-chloropyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
[6-methyl-2-(4-methylpyridine-2-carbonyl)-1H-indol-3-yl]acetic acid;
10 [5-chloro-2-(isoquinoline-3-carbonyl)-1H-indol-3-yl]acetic acid;
[6-chloro-2-(4-chlorobenzoyl)-5-fluoro-1H-indol-3-yl]acetic acid,
and a salt thereof.

18. A pharmaceutical composition useful for the treatment of a medical condition in which prostaglandins are implicated as pathogens, which comprises a
15 compound of the formula (I) of claim 1, and a pharmaceutically inert carrier.

19. A pharmaceutical composition according to claim 18, wherein the compound of the formula (I) is as defined in Claim 2.

20. A pharmaceutical composition according to claim 19, wherein the compound of the formula (I) is as defined in Claim 3.

20 21. A pharmaceutical composition according to claim 20, wherein the compound of the formula (I) is as defined in Claim 4.

22. A pharmaceutical composition according to claim 21, wherein the compound of the formula (I) is as defined in Claim 5.

25 23. A pharmaceutical composition according to claim 22, wherein the compound of the formula (I) is as defined in Claim 6.

24. A pharmaceutical composition according to claim 23, wherein the compound is as defined in Claim 7.

25. A pharmaceutical composition according to claim 24, wherein the compound is as defined in Claim 8.

30 26. A pharmaceutical composition according to claim 25, wherein the compound is as defined in Claim 9.

27. A pharmaceutical composition according to claim 26, wherein the compound is as defined in Claim 10.

28. A pharmaceutical composition according to claim 27, wherein the compound is as defined in Claim 11.

5 29. A pharmaceutical composition according to claim 28, wherein the compound is as defined in Claim 12.

30. A pharmaceutical composition according to claim 29, wherein the compound is as defined in Claim 13.

10 31. A pharmaceutical composition according to claim 30, wherein the compound is as defined in Claim 14.


32. A pharmaceutical composition according to claim 31, wherein the compound is as defined in Claim 15.

33. A pharmaceutical composition according to claim 32, wherein the compound is as defined in Claim 16.

15 34. A pharmaceutical composition according to claim 33, wherein the compound is as defined in Claim 17.

35. A method for the treatment of a medical condition in which prostaglandins are implicated as pathogens, in a mammalian subject, which comprises administering to said pharmaceutical composition according to claim 1.

20 36 A compound of the formula:



7-VI

wherein **B** is a suitable protecting group;

Q is selected from the following:

25 (a) phenyl optionally substituted with one, two or three substituents independently selected from

- 5 (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,
- 10 (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 15 (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 20 (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 25 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- 30 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);

(d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and


(e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R² and R³ are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; **R⁵** is C₁₋₆ alkyl; and

n is 0, 1, 2, 3 or 4.

37. A compound of the formula:



7-V

wherein **B** is a suitable protecting group;

Q is selected from the following:

(a) phenyl optionally substituted with one, two or three substituents independently selected from

(a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,


- 5 (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 10 (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 15 (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 20 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- 25 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- 30 (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and
- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R² and R³ are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted

with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; R⁵ is C₁₋₆ alkyl; and n is 0, 1, 2, 3 or 4.

38. A process for preparing a compound of the formula:



7-IV

10 wherein

Q is selected from the following:

(a) phenyl optionally substituted with one, two or three substituents independently selected from

(a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,

(a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-

substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),

(c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);

(d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and

(e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);


R² and R³ are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; and

n is 0, 1, 2, 3 or 4,


which process comprises the steps of:

- i) reacting a compound of the formula:



7-II

wherein B is a suitable protecting group; R⁵ is C₁₋₆ alkyl; X and n are as defined above, with a compound of the formula:




5

wherein E is halo and Q are as defined above, with a first base and a suitable solvent;

- ii) reacting the product of step i) with a second base.
 - iii) reacting the product of step ii) with an acid.
- 10 39. The process according to claim 38, wherein said first base is potassium carbonate, potassium bicarbonate, sodium bicarbonate, sodium carbonate or cesium carbonate.
40. The process according to claim 38, wherein said first base is potassium carbonate.
- 15 41. The process according to claim 38, wherein said second base is aqueous sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium pentoxide (followed by water), sodium methoxide (followed by water) or potassium *t*-butoxide (followed by water).
42. The process according to claim 38, wherein said second base is sodium
- 20 hydroxide.
43. The process according to claim 38, wherein said acid is aqueous hydrochloric acid, hydrobromic acid, sulfuric acid or ammonium chloride.
44. The process according to claim 38, wherein said acid is aqueous hydrochloric acid.
- 25 45. The process according to claim 38, wherein said solvent is *N,N*-dimethylacetamide, *N,N*-dimethylformamide, methyl ethyl ketone, acetone, or tetrahydrofuran.

46. The process according to claim 38, wherein said solvent is *N,N*-dimethylethylacetamide.

47. A process for preparing a compound of the formula:



7-IV

5 wherein

Q is selected from the following:

- (a) phenyl optionally substituted with one, two or three substituents independently selected from
 - (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,
 - (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
 - (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
 - (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-

substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino and CN,


- (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);
- (d) C_{3-7} cycloalkyl optionally substituted with one or two substituents independently selected from OH, C_{1-4} alkyl, halo and halo-substituted C_{1-4} alkyl; and
- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R^2 and R^3 are independently H, OH, C_{1-4} alkoxy, C_{1-4} alkyl or C_{1-4} alkyl substituted with halo, OH, C_{1-4} alkoxy, NH_2 or CN;

- X is independently selected from halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino, CN, HO- (C_{1-4}) alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $-NH_2S(O)_2NR^2NR^3$, acetyl, $-COOH$, $-C(O)O-C_{1-4}$ alkyl, C_{1-4} alkylsulfonylamino and C_{3-7} cycloalkyl; and

- n is 0, 1, 2, 3 or 4,

which process comprises reacting a compound of the formula:



7-VII

wherein R^5 is C_{1-6} alkyl; Q, X and n are as defined as before, with a base in a suitable solvent.


48. The process according to claim 47, wherein said base is sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate, sodium *t*-pentoxide, sodium methoxide, sodium ethoxide or potassium *t*-butoxide.

49. The process according to claim 47, wherein said base is sodium hydroxide.

50. The process according to claim 47, wherein said solvent is an aqueous mixture of methanol, ethanol, isopropyl alcohol or tetrahydrofuran.

51. The process according to claim 47, wherein said solvent is methanol containing water.

52. A process of preparing a compound of the formula:



7-VII

wherein

Q is selected from the following:

(a) phenyl optionally substituted with one, two or three substituents independently selected from

(a-1) halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino, CN, HO- (C_{1-4}) alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $-NH_2S(O)_2NR^2R^3$, acetyl, $-COOH$, $-C(O)O-C_{1-4}$ alkyl, C_{1-4} alkylsulfonylamino and C_{3-7} cycloalkyl,

(a-2) aryl or $-O-(CH_2)_n$ -aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently

selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

5 (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

10 (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

15 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),

20 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);

25 (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and

(e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);


30 **R² and R³** are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄

alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino, CN, $HO-(C_{1-4})$ alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $-NH_2S(O)_2NR^2NR^3$, acetyl, $-COOH$, $-C(O)O-C_{1-4}$ alkyl, C_{1-4} alkylsulfonylamino and C_{3-7} cycloalkyl; R^5 is C_{1-6} alkyl; and

5 n is 0, 1, 2, 3 or 4,

which process comprises reacting a compound of the formula:



7-VI

wherein B, Q, X, n and R^5 are as defined above with a base in a suitable solvent.


10 53. The process according to claim 52, wherein said base is 1,8-diazabicyclo[5.4.0]undec-7-ene, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,1,3,3-tetramethylguanidine, sodium *t*-pentoxide, sodium methoxide, or potassium *t*-butoxide.

54. The process according to claim 52, wherein said base is 1,8-diazabicyclo[5.4.0]undec-7-ene or potassium *t*-butoxide.

15 55. The process according to claim 52, wherein said solvent is *N,N*-dimethylacetamide, *N,N*-dimethylformamide, methyl ethyl ketone, acetone, or tetrahydrofuran.

56. The process according to claim 52, wherein said solvent is *N,N*-dimethylacetamide.

20 57. A process for preparing a compound of the formula:



7-VI

wherein B is a suitable protecting group;

Q is selected from the following:

- (a) phenyl optionally substituted with one, two or three substituents independently selected from
- 5 (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,
- 10 (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 15 (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 20 (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,
- 25 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),
- 30 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic

aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);


- (d) C_{3-7} cycloalkyl optionally substituted with one or two substituents independently selected from OH, C_{1-4} alkyl, halo and halo-substituted C_{1-4} alkyl; and
- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R^2 and R^3 are independently H, OH, C_{1-4} alkoxy, C_{1-4} alkyl or C_{1-4} alkyl substituted with halo, OH, C_{1-4} alkoxy, NH_2 or CN;

X is independently selected from halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino, CN, HO- (C_{1-4}) alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $-NH_2S(O)_2NR^2NR^3$, acetyl, $-COOH$, $-C(O)O-C_{1-4}$ alkyl, C_{1-4} alkylsulfonylamino and C_{3-7} cycloalkyl; R^5 is C_{1-6} alkyl; and

n is 0, 1, 2, 3 or 4,

which process comprises reacting a compound of the formula:



wherein B, Q, X, n and R^5 are as defined above, with a base in the presence of a solvent.


58. The process according to claim 57, wherein said base is potassium carbonate, potassium bicarbonate, sodium bicarbonate, sodium carbonate, or cesium carbonate.

59. The process according to claim 57, wherein said base is potassium carbonate.

60. The process according to claim 57, wherein said solvent is *N,N*-dimethylacetamide, *N,N*-dimethylformamide, methyl ethyl ketone, acetone, or tetrahydrofuran.

61. The process according to claim 57, wherein said solvent is *N,N*-dimethylacetamide.

62. A process for preparing a compound of the formula:



wherein **B** is a suitable protecting group;

10 **Q** is selected from the following:

(a) phenyl optionally substituted with one, two or three substituents independently selected from

(a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,

(a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-

substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),

(c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);

(d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and


(e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R² and R³ are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; R⁵ is C₁₋₆ alkyl; and


n is 0, 1, 2, 3 or 4,

which comprises reacting a compound of the formula:



7-II

wherein B, X, n and R⁵ are as defined above, with a compound of the formula:



wherein E is halo and Q are as defined above, with a base in the presence of a solvent.


63. The process according to claim 62, wherein said base is potassium carbonate, potassium bicarbonate, sodium bicarbonate, sodium carbonate, or cesium carbonate.

64. The process according to claim 62, wherein said base is potassium carbonate.

65. The process according to claim 62, wherein said solvent is *N,N*-dimethylacetamide, *N,N*-dimethylformamide, methyl ethyl ketone, acetone or tetrahydrofuran.

66. The process according to claim 62, wherein said solvent is *N,N*-dimethylacetamide.

67. A process for preparing a compound of the formula (XII):



wherein

Q is selected from the following:

- 20 (a) phenyl optionally substituted with one, two or three substituents independently selected from
- (a-1) halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋

₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,

5 (a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

10 (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

15 (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

20 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),

25 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);

30 (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and

- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R^1 is hydrogen, C_{1-4} alkyl or halo;


- R^2 and R^3 are independently H, OH, C_{1-4} alkoxy, C_{1-4} alkyl or C_{1-4} alkyl substituted with halo, OH, C_{1-4} alkoxy, NH_2 or CN;

R^5 is C_{1-6} alkyl;

- X is independently selected from halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino, CN, $HO-(C_{1-4})$ alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $-NH_2S(O)_2NR^2NR^3$, acetyl, $-COOH$, $-C(O)O-C_{1-4}$ alkyl, C_{1-4} alkylsulfonylamino and C_{3-7} cycloalkyl; and

n is 0, 1, 2, 3 or 4,


which process comprises treating a compound of the formula (X):



(X)

- wherein R^1 , R^5 , X , Q and n are as defined herein before, and B is a suitable protecting group, in the presence of a suitable base to obtain a compound of the formula (XII).

68. A process for preparing a compound of the formula (XII):



(XII)

wherein

- Q is selected from the following:

- (a) phenyl optionally substituted with one, two or three substituents independently selected from

(a-1) halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-

substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²R³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl,

(a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

(b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),

(c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);

(d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents

independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and

- (e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

5 **R¹** is hydrogen, C₁₋₄ alkyl or halo;

R² and **R³** are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;


R⁵ is C₁₋₆ alkyl;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄

10 alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; and


n is 0, 1, 2, 3 or 4,

15 which process comprises reacting a compound of the formula (IX):



(IX)

wherein **R¹**, **R⁵**, **X**, and **n** are as defined above, and **B** is a suitable protecting group, with a compound of the formula (XI):



(XI)

20 wherein **E** is halo and **Q** is as defined as before,

in the presence of a suitable base at a temperature of -40 °C to 200 °C to obtain a compound of the formula (XII).

69. A process according to Claim 68, wherein the reaction is carried out at a temperature of 0 °C to 100 °C

70. A process according to Claim 68, wherein the suitable base is potassium carbonate, cesium carbonate, sodium carbonate, sodium *tert*-butoxide, potassium *tert*-butoxide, sodium hydride, potassium hydride or potassium fluoride.

71. A process according to Claim 68, wherein the reaction is firstly carried out in the presence of a base for 2 minutes to a day; and then, another base is added to the reaction mixture.

72. A process according to Claim 71, wherein the reaction is firstly carried out for 30 minutes to 8 hours.

73. A process according to claims 67 and 68, wherein the suitable protecting group is methoxycarbonyl, ethoxycarbonyl, *tert*-butoxycarbonyl, benzyloxycarbonyl, phenylsulfonyl, *p*-toluenesulfonyl, methanesulfonyl or trifluoromethanesulfonyl.

74. A process according to claim 73, wherein the suitable protecting group is phenylsulfonyl, *p*-toluenesulfonyl, methanesulfonyl or trifluoromethanesulfonyl.

75. A process according to claim 71, wherein the first base is selected from sodium *tert*-butoxide, potassium *tert*-butoxide, sodium carbonate, potassium carbonate, cesium carbonate, sodium hydride, potassium hydride, sodium carbonate, potassium carbonate, cesium carbonate, potassium fluoride, 1,8-diazabicyclo[5.4.0]undec-7-ene, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,4-diazabicyclo[2.2.2]octane, pyridine, pyrrolidine, triethylamine, diisopropylamine, diisopropylethylamine and diethylisopropylamine; and the second base is selected from sodium *tert*-butoxide, potassium *tert*-butoxide, sodium carbonate, potassium carbonate, cesium carbonate, sodium hydride, potassium hydride, sodium carbonate, potassium carbonate, cesium carbonate, potassium fluoride, 1,8-diazabicyclo[5.4.0]undec-7-ene, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,4-diazabicyclo[2.2.2]octane, pyridine, pyrrolidine, triethylamine, diisopropylamine, diisopropylethylamine and diethylisopropylamine.

76. A process according to claim 75, wherein the first base is selected from potassium carbonate, cesium carbonate, sodium hydride and potassium fluoride; and the second base is selected from 1,8-diazabicyclo[5.4.0]undec-7-ene, cesium carbonate, pyrrolidine, diisopropylamine, triethylamine, diethylisopropylamine and diisopropylethylamine.

alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di-
(C_{1-4} alkyl)amino, C_{1-4} alkylamino and CN,

(a-3) 5-membered monocyclic aromatic group optionally substituted with
one, two or three substituents independently selected from halo, C_1 -
4 alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-
substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di-(C_{1-4}
alkyl)amino, C_{1-4} alkylamino and CN,

(a-4) 6-membered monocyclic aromatic group optionally substituted with
one, two or three substituents independently selected from halo, C_1 -
4 alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-
substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di-(C_{1-4}
alkyl)amino, C_{1-4} alkylamino and CN,

(b) a 6-membered monocyclic aromatic group containing one, two, three or
four nitrogen atom(s), and said monocyclic aromatic group being
optionally substituted with one, two or three substituents independently
selected from the above group (a-1), (a-2), (a-3) and (a-4),

(c) a 5-membered monocyclic aromatic group containing one heteroatom
selected from O, S and N and optionally containing one, two or three
nitrogen atom(s) in addition to said heteroatom, and said monocyclic
aromatic group being optionally substituted with one, two or three
substituents independently selected from the above group (a-1), (a-2), (a-
3) and (a-4);

(d) C_{3-7} cycloalkyl optionally substituted with one or two substituents
independently selected from OH, C_{1-4} alkyl, halo and halo-substituted C_1 -
4 alkyl; and

(e) a benzo-fused heterocycle optionally substituted with one, two or three
substituents independently selected from the group (a-1);

R^1 is hydrogen, C_{1-4} alkyl or halo;

R^2 and R^3 are independently H, OH, C_{1-4} alkoxy, C_{1-4} alkyl or C_{1-4} alkyl substituted
with halo, OH, C_{1-4} alkoxy, NH_2 or CN;


X is independently selected from halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4}

77. A process according to claim 75, wherein the first base is potassium
carbonate, cesium carbonate or potassium fluoride; and the second base is 1,8-
diazabicyclo[5.4.0]undec-7-ene, potassium tert-butoxide or cesium carbonate.

78. A process according to claim 77, wherein the combination of the first base
and the second base (first base/second base) is selected from potassium carbonate/1,8-
diazabicyclo[5.4.0]undec-7-ene, potassium carbonate/cesium carbonate, cesium
carbonate/potassium tert-butoxide, cesium carbonate/1,8-diazabicyclo[5.4.0]undec-7-
ene and potassium fluoride/1,8-diazabicyclo[5.4.0]undec-7-ene and potassium
fluoride/cesium carbonate.

79. A process according to claim 78, wherein the combination of the first base
and the second base (first base/second base) is selected from potassium carbonate/1,8-
diazabicyclo[5.4.0]undec-7-ene, potassium carbonate/cesium carbonate and cesium
carbonate/potassium tert-butoxide.

80. A process for preparing a compound of the formula (VIII):



wherein

Q is selected from the following:

(a) phenyl optionally substituted with one, two or three substituents
independently selected from


(a-1) halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-
substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di-(C_{1-4}
alkyl)amino, C_{1-4} alkylamino, CN, HO-(C_{1-4}) alkyl, C_{1-4} alkoxy- C_1 -
4 alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $-NH_2S(O)_2NR^2R^3$, acetyl,
-COOH, -C(O)O- C_{1-4} alkyl, C_{1-4} alkylsulfonylamino and C_{3-7}
cycloalkyl,

(a-2) aryl or -O-(CH₂)_n-aryl, and the aryl or aryl moiety being optionally
substituted with one, two or three substituents independently
selected from halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4}

alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino, CN, $HO-(C_{1-4})$ alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $-NH_2S(O)_2NR^2NR^3$, acetyl, $-COOH$, $-C(O)O-C_{1-4}$ alkyl, C_{1-4} alkylsulfonylamino and C_{3-7} cycloalkyl; and

5 n is 0, 1, 2, 3 or 4,


which process comprises treating a compound of the formula (X):



(X)

wherein R^1 , R^5 , X, Q and n are as defined here before, with a suitable base under hydrolyzing conditions to obtain the compound of formula (VIII).

10 81. A process for preparing a compound of the formula (VIII):



(VIII)

wherein

Q is selected from the following:

15 (a) phenyl optionally substituted with one, two or three substituents independently selected from

(a-1) halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino, CN, $HO-(C_{1-4})$ alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $-NH_2S(O)_2NR^2R^3$, acetyl, $-COOH$, $-C(O)O-C_{1-4}$ alkyl, C_{1-4} alkylsulfonylamino and C_{3-7} cycloalkyl,


20

(a-2) aryl or $-O-(CH_2)_n$ -aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently

alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino, CN, $HO-(C_{1-4})$ alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $-NH_2S(O)_2NR^2NR^3$, acetyl, $-COOH$, $-C(O)O-C_{1-4}$ alkyl, C_{1-4} alkylsulfonylamino and C_{3-7} cycloalkyl; and

5 n is 0, 1, 2, 3 or 4,


which process comprises treating a compound of the formula (X):



(X)

wherein R^1 , R^5 , X, Q and n are as defined here before, with a suitable base under hydrolyzing conditions to obtain the compound of formula (VIII).

10 81. A process for preparing a compound of the formula (VIII):



(VIII)

wherein

Q is selected from the following:

(a) phenyl optionally substituted with one, two or three substituents independently selected from

(a-1) halo, C_{1-4} alkyl, halo-substituted C_{1-4} alkyl, OH, C_{1-4} alkoxy, halo-substituted C_{1-4} alkoxy, C_{1-4} alkylthio, NO_2 , NH_2 , di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkylamino, CN, $HO-(C_{1-4})$ alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{1-4} alkylsulfonyl, aminosulfonyl, $-NH_2S(O)_2NR^2R^3$, acetyl, $-COOH$, $-C(O)O-C_{1-4}$ alkyl, C_{1-4} alkylsulfonylamino and C_{3-7} cycloalkyl,

(a-2) aryl or $-O-(CH_2)_n$ -aryl, and the aryl or aryl moiety being optionally substituted with one, two or three substituents independently

selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

5 (a-3) 5-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

10 (a-4) 6-membered monocyclic aromatic group optionally substituted with one, two or three substituents independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino and CN,

15 (b) a 6-membered monocyclic aromatic group containing one, two, three or four nitrogen atom(s), and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4),

20 (c) a 5-membered monocyclic aromatic group containing one heteroatom selected from O, S and N and optionally containing one, two or three nitrogen atom(s) in addition to said heteroatom, and said monocyclic aromatic group being optionally substituted with one, two or three substituents independently selected from the above group (a-1), (a-2), (a-3) and (a-4);

25 (d) C₃₋₇ cycloalkyl optionally substituted with one or two substituents independently selected from OH, C₁₋₄ alkyl, halo and halo-substituted C₁₋₄ alkyl; and


(e) a benzo-fused heterocycle optionally substituted with one, two or three substituents independently selected from the group (a-1);

R¹ is hydrogen, C₁₋₄ alkyl or halo;

30 R² and R³ are independently H, OH, C₁₋₄ alkoxy, C₁₋₄ alkyl or C₁₋₄ alkyl substituted with halo, OH, C₁₋₄ alkoxy, NH₂ or CN;

X is independently selected from halo, C₁₋₄ alkyl, halo-substituted C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, halo-substituted C₁₋₄ alkoxy, C₁₋₄ alkylthio, NO₂, NH₂, di-(C₁₋₄ alkyl)amino, C₁₋₄ alkylamino, CN, HO-(C₁₋₄) alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₁₋₄ alkylsulfonyl, aminosulfonyl, -NH₂S(O)₂NR²NR³, acetyl, -COOH, -C(O)O-C₁₋₄ alkyl, C₁₋₄ alkylsulfonylamino and C₃₋₇ cycloalkyl; and
 5 n is 0, 1, 2, 3 or 4,

which process comprises hydrolyzing a compound of the formula (XII):



wherein R⁵ is C₁₋₆ alkyl, R¹, X, Q and n are as defined herein before.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 98/02065

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D209/18 A61K31/40 C07D401/06 C07D403/06 C07D417/06
 C07D405/06 C07D409/10 C07D405/10 C07D401/10 C07D417/10
 C07D413/06 C07D409/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 510 368 A (MERCK FROSST CANADA INC.) 23 April 1996 cited in the application see claims	1,18
A	WO 96 37469 A (MERCK FROSST CANADA INC.) 28 November 1996 cited in the application see claims	1,18

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

19 February 1999

Date of mailing of the international search report

26/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3018

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB 98/02065

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 35
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim 35 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 98/02065

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5510368 A	23-04-1996	AU 5683196 A	11-12-1996
		CA 2219115 A	28-11-1996
		WO 9637468 A	28-11-1996
WO 9637469 A	28-11-1996	US 5604253 A	18-02-1997
		AU 5683296 A	11-12-1996
		CA 2219111 A	28-11-1996